

Delayed Cutaneous Reactions to Iodinated Contrast

Daniel R. Mazori, MD; Arielle R. Nagler, MD; Miriam Keltz Pomeranz, MD

PRACTICE POINTS

- Delayed cutaneous reactions to iodinated contrast (IC) are common, but patients frequently are misdiagnosed and inadvertently readministered the offending agent.
- The most common IC-induced delayed reactions are self-limited exanthematous eruptions that develop within 1 week of exposure.
- Risk factors for delayed reactions to IC include atopy, contrast exposure during high pollen season, use of the agent iodixanol, a history of other cutaneous drug eruptions, elevated serum creatinine levels, and treatment with recombinant interleukin 2.
- Dermatologists can help prevent recurrent reactions in patients who require repeated exposure to IC by recommending gadolinium-based contrast agents and/or premedication.

Iodinated contrast (IC) often is overlooked as a cause of delayed drug reactions. As a result, patients may continue to be readministered these agents, experiencing preventable morbidity. We report an illustrative case of a delayed cutaneous reaction to IC and review the literature on the epidemiology and management of this under-recognized phenomenon.

Cutis. 2018;101:433-435.

Case Report

A 67-year-old woman with a history of allergic rhinitis presented in the spring with a pruritic eruption of 2 days' duration that began on the abdomen and spread to the chest, back, and bilateral arms. Six days prior to the onset of the symptoms she underwent computed tomography (CT) of the abdomen and pelvis to evaluate abdominal pain and peripheral eosinophilia. Two iodinated contrast (IC)

agents were used: intravenous iohexol and oral diatrizoate meglumine–diatrizoate sodium. The eruption was not preceded by fever, malaise, sore throat, rhinorrhea, cough, arthralgia, headache, diarrhea, or new medication or supplement use. The patient denied any history of drug allergy or cutaneous eruptions. Her current medications, which she had been taking long-term, were aspirin, lisinopril, diltiazem, levothyroxine, esomeprazole, paroxetine, gabapentin, and diphenhydramine.

Physical examination was notable for erythematous, blanchable, nontender macules coalescing into patches on the trunk and bilateral arms (Figure). There was slight erythema on the nasolabial folds and ears. The mucosal surfaces and distal legs were clear. The patient was afebrile. Her white blood cell count was $12.5 \times 10^9/L$ with 32.3% eosinophils (baseline: white blood cell count, $14.8 \times 10^9/L$; 22% eosinophils) (reference range, $4.8\text{--}10.8 \times 10^9/L$; 1%–4% eosinophils). Her comprehensive metabolic panel was within reference range. The human immunodeficiency virus 1/2 antibody immunoassay was nonreactive.

The patient was diagnosed with an exanthematous eruption caused by IC and was treated with oral hydroxyzine and triamcinolone acetonide cream 0.1%. The eruption resolved within 2 weeks without recurrence at 3-month follow-up.

Comment

Delayed cutaneous eruptions caused by IC are under-recognized in medicine and are infrequently described

RELATED CONTENT ONLINE

Nephrogenic Systemic Fibrosis: Is Gadolinium the Missing Piece to the Puzzle?

>> <http://bit.ly/2sn0F2G>

From the Ronald O. Perelman Department of Dermatology, New York University School of Medicine, New York. The authors report no conflict of interest.

Correspondence: Miriam Keltz Pomeranz, MD, 240 E 38th St, 12th Floor, New York, NY 10016 (miriam.pomeranz@nyumc.org).



Erythematous, blanchable, nontender macules coalescing into patches on the abdomen (A) and left arm (B).

in the dermatology literature.¹ Unlike urticaria and other well-known immediate reactions to IC, delayed reactions develop when patients are less likely to be under medical supervision.² Moreover, only 12% to 33% of patients with delayed reactions to IC seek medical attention.³⁻⁶ As a result, these delayed reactions often are attributed to other causes.¹ Patients may then be unknowingly reexposed to the offending contrast agent and experience recurrent eruptions, such as in one fatal case of toxic epidermal necrolysis (TEN).⁷⁻¹¹ Given the role of dermatologists in the diagnosis and prevention of cutaneous drug reactions, it is important to be mindful of delayed cutaneous eruptions caused by IC.

Clinical Presentation of Delayed Reactions—Most delayed cutaneous reactions to IC present as exanthematous eruptions in the week following a contrast-enhanced CT scan or coronary angiogram.^{2,12} The reactions tend to resolve within 2 weeks of onset, and the treatment is largely supportive with antihistamines and/or corticosteroids for the associated pruritus.^{2,5,6} In a study of 98 patients with a history of delayed reactions to IC, delayed-onset urticaria and angioedema also have been reported with incidence rates of 19% and 24%, respectively.² Other reactions are less common. In the same study, 7% of patients developed palpable purpura; acute generalized exanthematous pustulosis; bullous, flexural, or psoriasislake exanthema;

exfoliative eruptions; or purpura and a maculopapular eruption combined with eosinophilia.² There also have been reports of IC-induced erythema multiforme,³ fixed drug eruptions,^{10,11} symmetrical drug-related intertriginous and flexural exanthema,¹³ cutaneous vasculitis,¹⁴ drug reactions with eosinophilia and systemic symptoms,¹⁵ Stevens-Johnson syndrome/TEN,^{7,8,16,17} and iododerma.¹⁸

IC Agents—Virtually all delayed cutaneous reactions to IC reportedly are due to intravascular rather than oral agents,^{1,2,19} with the exception of iododerma¹⁸ and 1 reported case of TEN.¹⁷ Intravenous iohexol was most likely the offending drug in our case. In a prospective cohort study of 539 patients undergoing CT scans, the absolute risk for developing a delayed cutaneous reaction (defined as rash, itching, or skin redness or swelling) to intravascular iohexol was 9.4%.²⁰ Randomized, double-blind studies have found that the risk for delayed cutaneous eruptions is similar among various types of IC, except for iodixanol, which confers a higher risk.^{5,6,21}

Risk Factors—Interestingly, analyses have shown that delayed reactions to IC are more common in atopic patients and during high pollen season.²² Our patient displayed these risk factors, as she had allergic rhinitis and presented for evaluation in late spring when local pollen counts were high. Additionally, patients who develop delayed reactions to IC are notably more likely than controls to have a history of other cutaneous drug reactions, serum creatinine levels greater than 2.0 mg/dL (reference range, 0.6–1.2 mg/dL),³ or history of treatment with recombinant interleukin 2.¹⁹

Patients with a history of delayed reactions to IC are not at increased risk for immediate reactions and vice versa.^{2,3} This finding is consistent with the evidence that delayed and immediate reactions to IC are mechanistically unrelated.²³ Additionally, seafood allergy is not a risk factor for either immediate or delayed reactions to IC, despite a common misconception among physicians and patients because seafood is iodinated.^{24,25}

Reexposure to IC—Patients who have had delayed cutaneous reactions to IC are at risk for similar eruptions upon reexposure. Although the reactions are believed to be cell mediated, skin testing with IC is not sensitive enough to reliably identify tolerable alternatives.¹² Consequently, gadolinium-based agents have been recommended in patients with a history of reactions to IC if additional contrast-enhanced studies are needed.^{13,26} Iodinated and gadolinium-based contrast agents do not cross-react, and gadolinium is compatible with studies other than magnetic resonance imaging.^{1,27}

Premedication—Despite the absence of cross-reactivity, the American College of Radiology considers patients with hypersensitivity reactions to IC to be at increased risk for reactions to gadolinium but does not make specific recommendations regarding premedication given the dearth of data.¹ As a result, premedication may be considered prior to gadolinium administration

depending on the severity of the delayed reaction to IC. Additionally, premedication may be beneficial in cases in which gadolinium is contraindicated and IC must be reused. In a retrospective study, all patients with suspected delayed reactions to IC tolerated IC or gadolinium contrast when pretreated with corticosteroids with or without antihistamines.²⁸ Regimens with corticosteroids and either cyclosporine or intravenous immunoglobulin also have prevented the recurrence of IC-induced exanthematous eruptions and Stevens-Johnson syndrome.^{29,30} Despite these reports, delayed cutaneous reactions to IC have recurred in other patients receiving corticosteroids, antihistamines, and/or cyclosporine for premedication or concurrent treatment of an underlying condition.^{16,29-31}

Conclusion

It is important for dermatologists to recognize IC as a cause of delayed drug reactions. Current awareness is limited, and as a result, patients often are reexposed to the offending contrast agents unsuspectingly. In addition to diagnosing these eruptions, dermatologists may help prevent their recurrence if future contrast-enhanced studies are required by recommending gadolinium-based agents and/or premedication.

REFERENCES

- Cohan RH, Davenport MS, Dillman JR, et al; ACR Committee on Drugs and Contrast Media. *ACR Manual on Contrast Media*. 9th ed. Reston, VA: American College of Radiology; 2013.
- Brockow K, Romano A, Aberer W, et al; European Network of Drug Allergy and the EAACI Interest Group on Drug Hypersensitivity. Skin testing in patients with hypersensitivity reactions to iodinated contrast media—a European multicenter study. *Allergy*. 2009;64:234-241.
- Hosoya T, Yamaguchi K, Akutsu T, et al. Delayed adverse reactions to iodinated contrast media and their risk factors. *Radiat Med*. 2000;18:39-45.
- Rydberg J, Charles J, Aspelin P. Frequency of late allergy-like adverse reactions following injection of intravascular non-ionic contrast media: a retrospective study comparing a non-ionic monomeric contrast medium with a non-ionic dimeric contrast medium. *Acta Radiol*. 1998;39:219-222.
- Sutton AG, Finn P, Grech ED, et al. Early and late reactions after the use of iopamidol 340, ioxaglate 320, and iodixanol 320 in cardiac catheterization. *Am Heart J*. 2001;141:677-683.
- Sutton AG, Finn P, Campbell PG, et al. Early and late reactions following the use of iopamidol 340, iomeprol 350 and iodixanol 320 in cardiac catheterization. *J Invasive Cardiol*. 2003;15:133-138.
- Brown M, Yowler C, Brandt C. Recurrent toxic epidermal necrolysis secondary to iopromide contrast. *J Burn Care Res*. 2013;34:E53-E56.
- Rosado A, Canto G, Veleiro B, et al. Toxic epidermal necrolysis after repeated injections of iohexol. *AJR Am J Roentgenol*. 2001;176:262-263.
- Peterson A, Katzberg RW, Fung MA, et al. Acute generalized exanthematous pustulosis as a delayed dermatotoxic reaction to IV-administered nonionic contrast media. *AJR Am J Roentgenol*. 2006;187:W198-W201.
- Good AE, Novak E, Sonda LP III. Fixed eruption and fever after urography. *South Med J*. 1980;73:948-949.
- Benson PM, Giblin WJ, Douglas DM. Transient, nonpigmenting fixed drug eruption caused by radiopaque contrast media. *J Am Acad Dermatol*. 1990;23(2, pt 2):379-381.
- Torres MJ, Gomez F, Doña I, et al. Diagnostic evaluation of patients with nonimmediate cutaneous hypersensitivity reactions to iodinated contrast media. *Allergy*. 2012;67:929-935.
- Scherer K, Harr T, Bach S, et al. The role of iodine in hypersensitivity reactions to radio contrast media. *Clin Exp Allergy*. 2010;40:468-475.
- Reynolds NJ, Wallington TB, Burton JL. Hydralazine predisposes to acute cutaneous vasculitis following urography with iopamidol. *Br J Dermatol*. 1993;129:82-85.
- Belhadjali H, Bouzgarrou L, Youssef M, et al. DRESS syndrome induced by sodium meglumine ioxitalamate. *Allergy*. 2008;63:786-787.
- Baldwin BT, Lien MH, Khan H, et al. Case of fatal toxic epidermal necrolysis due to cardiac catheterization dye. *J Drugs Dermatol*. 2010;9:837-840.
- Schmidt BJ, Foley WD, Bohorfoush AG. Toxic epidermal necrolysis related to oral administration of diluted diatrizoate meglumine and diatrizoate sodium. *AJR Am J Roentgenol*. 1998;171:1215-1216.
- Young AL, Grossman ME. Acute iododerma secondary to iodinated contrast media. *Br J Dermatol*. 2014;170:1377-1379.
- Choyke PL, Miller DL, Lotze MT, et al. Delayed reactions to contrast media after interleukin-2 immunotherapy. *Radiology*. 1992;183:111-114.
- Loh S, Bagheri S, Katzberg RW, et al. Delayed adverse reaction to contrast-enhanced CT: a prospective single-center study comparison to control group without enhancement. *Radiology*. 2010;255:764-771.
- Bertrand P, Delhommais A, Alison D, et al. Immediate and delayed tolerance of iohexol and ioxaglate in lower limb phlebography: a double-blind comparative study in humans. *Acad Radiol*. 1995;2:683-686.
- Munehchika H, Hiramatsu Y, Kudo S, et al. A prospective survey of delayed adverse reactions to iohexol in urography and computed tomography. *Eur Radiol*. 2003;13:185-194.
- Guéant-Rodriguez RM, Romano A, Barbaud A, et al. Hypersensitivity reactions to iodinated contrast media. *Curr Pharm Des*. 2006;12:3359-3372.
- Huang SW. Seafood and iodine: an analysis of a medical myth. *Allergy Asthma Proc*. 2005;26:468-469.
- Baig M, Farag A, Sajid J, et al. Shellfish allergy and relation to iodinated contrast media: United Kingdom survey. *World J Cardiol*. 2014;6:107-111.
- Böhm I, Schild HH. A practical guide to diagnose lesser-known immediate and delayed contrast media-induced adverse cutaneous reactions. *Eur Radiol*. 2006;16:1570-1579.
- Ose K, Doue T, Zen K, et al. "Gadolinium" as an alternative to iodinated contrast media for X-ray angiography in patients with severe allergy. *Circ J*. 2005;69:507-509.
- Jingu A, Fukuda J, Taketomi-Takahashi A, et al. Breakthrough reactions of iodinated and gadolinium contrast media after oral steroid premedication protocol. *BMC Med Imaging*. 2014;14:34.
- Romano A, Artesani MC, Andriolo M, et al. Effective prophylactic protocol in delayed hypersensitivity to contrast media: report of a case involving lymphocyte transformation studies with different compounds. *Radiology*. 2002;225:466-470.
- Hebert AA, Bogle MA. Intravenous immunoglobulin prophylaxis for recurrent Stevens-Johnson syndrome. *J Am Acad Dermatol*. 2004;50:286-288.
- Hasdenteufel F, Waton J, Cordebar V, et al. Delayed hypersensitivity reactions caused by iodixanol: an assessment of cross-reactivity in 22 patients. *J Allergy Clin Immunol*. 2011;128:1356-1357.