

## New 'legal' highs: Kratom and methoxetamine

Julianna D. Troy, MD, MPH

Dr. Troy is a Psychiatrist with Counseling and Psychological Services, Purdue University, West Lafayette, Indiana.

### Disclosure

Dr. Troy reports no financial relationship with any company whose products are mentioned in this article or with manufacturers of competing products.

The demand for “legal highs”—intoxicating natural or synthetic substances that are not prohibited by law—continues to increase. Young adults may use these substances, which are widely available on the internet, at “head shops,” and at gas stations. Such substances frequently cause adverse medical and psychiatric effects, exemplified by recent reports concerning the dangers of using synthetic cannabinoids (eg, “Spice,” “K2”) and synthetic cathinones (“bath salts”). Although these 2 substances now are illegal in many jurisdictions, other novel substances of misuse remain legal and widely available, including Kratom and methoxetamine.

Because these substances usually are not detectable on standard urine toxicology screens, clinicians need to be aware of them to be able to take an accurate substance use history, consider possible dangerous interactions with prescribed psychotropics, and address medical and psychiatric complications.

**Kratom** is an herbal product derived from *Mitragyna speciosa*, a plant native to Southeast Asia. Traditionally used as a medicinal herb, it increasingly is being used for recreational purposes and remains legal and widely available in the United States. Kratom’s leaves contain multiple alkaloids, including mitragynine and 7-hydroxymitragynine, which are believed to act as agonists at the  $\mu$ -opioid receptor. Mitragynine also may have agonist activity at post-synaptic  $\alpha$ 2-adrenergic receptors, as well as antagonist activity at 5-HT2A receptors.<sup>1</sup> Mitragynine is 13 times more potent than morphine, and 7-hydroxymitragynine is 4 times more potent than mitragynine.<sup>2</sup>

Kratom is available as leaves, powdered leaves, or gum. It can be smoked, brewed into tea, or mixed with liquid and ingested. Effects are dose-dependent; lower doses tend to produce a stimulant effect and higher doses produce an opioid effect. A typical dose is 1 to 8 g.<sup>3</sup> Users may take Kratom to experience euphoria or analgesia, or to self-treat opioid withdrawal symptoms.<sup>3</sup> Kratom withdrawal syndrome shares many features of classic opioid withdrawal—diarrhea, rhinorrhea, cravings, anxiety, tremor, myalgia, sweating, and irritability—but has been reported to be less severe and shorter-lasting.<sup>1</sup> Kratom withdrawal, like opioid withdrawal, may respond to supportive care in combination with opioid-replacement therapy. Airway management and naloxone treatment may be needed on an emergent basis if a user develops respiratory depression.<sup>2</sup> There have been case reports of seizures occurring following Kratom use.<sup>2</sup>

**Methoxetamine** is a ketamine analog originally developed as an alternative to ketamine. It isn’t classified as a controlled substance in the United States and is available on the internet.<sup>2</sup> Methoxetamine is a white powder typically snorted or taken sublingually, although it can be injected intramuscularly. Because methoxetamine’s structure is similar to ketamine, its mechanism of action is assumed to involve glutamate N-methyl-D-aspartate receptor antagonism and dopamine reuptake inhibition. Doses range from 20 to 100 mg orally and 10 to 50 mg when injected. Effects may not be apparent for 30 to 90 minutes after the drug is snorted, which may cause users to take another dose or ingest a different substance, possibly leading



Discuss this article at [www.facebook.com/CurrentPsychiatry](http://www.facebook.com/CurrentPsychiatry)

to synergistic adverse effects. Effects may emerge within 5 minutes when injected. The duration of effect generally is 5 to 7 hours—notably longer than ketamine—but as little as 1 hour when injected.

No clinical human or animal studies have been conducted on methoxetamine, which makes it difficult to ascertain the drug's true clinical and toxic effects; instead, these effects must be surmised from user reports and case studies. Desired effects described by users are similar to those of ketamine: dissociation, short-term mood elevation, visual hallucinations, and alteration of sensory experiences. Reported adverse effects include catatonia, confusion, agitation, and depression.<sup>4</sup> In addition, methoxetamine may induce sympathomimetic toxicity as evidenced by tachycardia and hypertension. Researchers have suggested that patients who experience methoxetamine toxicity and require emergency treatment be managed with supportive care and benzodiazepines.<sup>5</sup>

## Staying current is key

New and potentially dangerous substances are being produced so quickly distributors are able to stay ahead of regulatory efforts. When one substance is declared illegal, another related substance quickly is available to take its place. To provide the best care for our patients, it is essential for psychiatrists to stay up-to-date about these novel substances.

### References

1. McWhirter L, Morris S. A case report of inpatient detoxification after kratom (*Mitragyna speciosa*) dependence. *Eur Addict Res.* 2010;16(4):229-231.
2. Rosenbaum CD, Carreiro SP, Babu KM. Here today, gone tomorrow...and back again? A review of herbal marijuana alternatives (K2, Spice), synthetic cathinones (bath salts), Kratom, *Salvia divinorum*, methoxetamine, and piperazines. *J Med Toxicol.* 2012;8(1):15-32.
3. Boyer EW, Babu KM, Macalino GE. Self-treatment of opioid withdrawal with a dietary supplement, Kratom. *Am J Addict.* 2007;16(5):352-356.
4. Corazza O, Schifano F, Simonato P, et al. Phenomenon of new drugs on the Internet: the case of ketamine derivative methoxetamine. *Hum Psychopharmacol.* 2012;27(2):145-149.
5. Wood DM, Davies S, Puchnarewicz M, et al. Acute toxicity associated with the recreational use of the ketamine derivative methoxetamine. *Eur J Clin Pharmacol.* 2012; 68(5):853-856.

**Kratom and methoxetamine usually are undetectable on toxicology screens**

## Want to better understand challenges and opportunities in schizophrenia treatment?

Don't miss this new supplement to CURRENT PSYCHIATRY!

This supplement reviews the safety and efficacy challenges that clinicians and their adult patients with schizophrenia face when considering treatment options. The supplement also presents data from clinical trials of Latuda® (lurasidone HCl), an atypical antipsychotic indicated for the treatment of adult patients with schizophrenia.



### INDICATIONS AND USAGE

LATUDA is an atypical antipsychotic indicated for the treatment of patients with schizophrenia. Efficacy was established in five 6-week controlled studies of adult patients with schizophrenia. The effectiveness of LATUDA for longer-term use, that is, for more than 6 weeks, has not been established in controlled studies. Therefore, the physician who elects to use LATUDA for extended periods should periodically re-evaluate the long-term usefulness of the drug for the individual patient.

### IMPORTANT SAFETY INFORMATION FOR LATUDA

#### WARNING: INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

See full prescribing information for complete boxed warning.

- Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death.
- LATUDA is not approved for the treatment of patients with dementia-related psychosis.

Please see additional Important Safety Information, including **Boxed Warning**, and Brief Summary of Prescribing Information on adjacent pages.

This supplement is sponsored by



is a registered trademark of Dainippon Sumitomo Pharma Co., Ltd. Sunovion Pharmaceuticals Inc. is a U. S. subsidiary of Dainippon Sumitomo Pharma Co., Ltd. ©2013 Sunovion Pharmaceuticals Inc. All rights reserved. 2/13 LAT021-13

**Current**  
PSYCHIATRY

View this important update today at  
[CurrentPsychiatry.com](http://CurrentPsychiatry.com).  
Click on **Supplements**.