

Due to a submission error, the article, "Safety and Efficacy of a New Extended-Release Formulation of Minocycline," in the October 2006 supplement (*Cutis*. 2006;78[suppl 4]:21-31.) listed the incorrect available tablet strength of extended-release minocycline hydrochloride in subjects weighing 132 to 199 lb (60.00–90.45 kg)(Table 2). The correct tablet strength is 90 mg. Consequently, the table containing the weight-based dosing schedule in its corrected form is reprinted below.

Table 2.

### Extended-Release Minocycline Hydrochloride Weight-Based Dosing Schedule

Subject's Weight, lb (kg)	Available Tablet Strength, mg	Actual mg/kg Dose
99–131 (45.00–59.54)	45	1.00–0.76
132–199 (60.00–90.45)	90	1.50–1.00
200–300 (90.91–136.36)	135	1.48–0.99

In addition, a corrected acknowledgment is listed below.

*Acknowledgments*—The Solodyn Phase 2 Study Group: Alicia Bucko, DO, Albuquerque, New Mexico; Scott M. Dinehart, MD, Little Rock, Arkansas; Frank E. Dunlap, MD, Tucson, Arizona; Michael Jarratt, MD, Austin, Texas; Terry M. Jones, MD, PA, Bryan, Texas; Lewis Kaminester, MD, North Palm Beach, Florida; Steven Kempers, MD, Fridley, Minnesota; Robert W. Loss, MD, Rochester, New York; David Pariser, MD, Norfolk, Virginia; Ronald Savin, MD, New Haven, Connecticut; Daniel Stewart, DO, Clinton Township, Michigan; Dow B. Stough, MD, Hot Springs, Arkansas; Helen Mary Torok, MD, Medina, Ohio; Jonathan Weiss, MD, Snellville, Georgia; Donna Zeide, MD, Palm Beach Gardens, Florida.

The Solodyn Phase 3 Study Group: Carole C. Aponte, MD, Dallas, Texas; Elizabeth Arthur, MD, Rochester, New York; Karl R. Beutner, MD, Vallejo, California; Alicia Bucko, DO; Jeffrey A. Carmel, MD, Fremont, California; Scott M. Dinehart, MD; Frank E. Dunlap, MD; Alan B. Fleischer, Jr, MD, Winston-Salem, North Carolina; Michael Gold, MD, Nashville, Tennessee; Paul Harrison, MD, Sandy, Utah; James H. Herndon, Jr, MD, Dallas, Texas; Jo Lynne Herzog, MD, Birmingham, Alabama; Michael Jarratt, MD; Terry M. Jones, MD, PA; Steven Kempers, MD; Patricia C. Lee, MD, Houston, Texas; Robert W. Loss, MD; Keith Loven, MD, Goodlettsville, Tennessee; Larry Parish, MD, Philadelphia, Pennsylvania; Toivo E. Rist, MD, Knoxville, Tennessee; Stacy R. Smith, MD, La Jolla, California; Daniel Stewart, DO; Dow B. Stough, MD; Leonard Swinyer, MD, Salt Lake City, Utah; Helen Mary Torok, MD; Stephen K. Tying, MD, PhD, Houston, Texas; Jonathan Weiss, MD; Hector Wiltz, MD, Miami, Florida; Paul S. Yamauchi, MD, PhD, Santa Monica, California; Donna Zeide, MD.

*Cutis*® makes every possible effort to ensure accuracy in its articles and apologizes for the mistakes.