

# Inherited thrombophilia and adverse pregnancy outcomes: What the evidence shows

Recent findings suggest that inherited thrombophilias lead to habitual fetal loss, preeclampsia, and placental abruption. Here, an expert examines the role of gene polymorphism-related clotting abnormalities in these outcomes and reviews studies of promising interventions.

**W**hen obstetric histories of habitual or late fetal loss, preeclampsia, intra-uterine growth retardation, and placental abruption or infarct suggest placental compromise, a search for thrombophilia is warranted. This article examines:

- evidence implicating inherited thrombophil-

- ia polymorphisms in these adverse outcomes
- promising findings of 2 interventional studies of pregnancy outcomes in thrombophilia carriers: enoxaparin treatment for women experiencing habitual late fetal loss, and folic acid treatment for severe preeclampsia.

## KEY POINTS

- Because polymorphisms affecting thrombotic potential are frequent (2% to 20% of the general population), even a slightly increased risk of thrombosis affects a far greater number of pregnancies than complete deficiencies of clotting factor proteins S and C or antithrombin III.
- Observational studies suggest a role for heparin or folate supplementation in women with thrombophilia polymorphisms and prior adverse pregnancy outcomes.
- Hyperhomocysteinemia, which sometimes develops from the influences of a polymorphism, has been linked to preeclampsia and may contribute to placental abruption and infarct.
- The Leiden polymorphism of the factor V, with its greater propensity for thrombosis, occurs at increased frequency in women who experience habitual miscarriages.

## Polymorphisms linked to thrombotic risk

**T**he adverse pregnancy outcomes described above represent a spectrum of disease with considerable overlap in etiology. The risk of each is increased in the presence of even 1 of the other complications.

A similar placental pathology is common to these outcomes, suggesting aberrant maternal/fetal perfusion as a unifying feature. The causes of such perfusion abnormalities are diverse, although placental infarct and

• *Dr. Wilkins-Haug is chief, division of reproductive genetics, department of obstetrics and gynecology; and director, Center for Fetal Medicine and Prenatal Genetics at Brigham and Women's Hospital, Boston, Mass. She also is associate professor of obstetrics, gynecology, and reproductive biology, Harvard Medical School.*

microthrombi suggest an underlying perturbation of the clotting cascade.

Although clotting factor deficiencies including protein S, protein C, and antithrombin III have long been known to increase thrombotic disease risk, these hereditary deficiencies occur in less than 1% of the general population, and thus contribute minimally to placental compromise and adverse outcomes.<sup>1</sup>

Of greater clinical significance are changes in DNA sequences collectively known as gene polymorphisms. These gene sequence changes occur with variable population frequency and impact on protein function.

Now recognized are several polymorphic genes which modulate the balance between the opposing prothrombotic and antithrombotic actions of the clotting cascade. Polymorphisms of clotting-cascade proteins occur in 1% to 4% of the population, with variable impact on thrombotic risk.

### **Polymorphisms linked to endothelial damage**

**G**ene polymorphisms outside the clotting cascade also can increase the risk of thrombosis. Inflammation of the endothelial lining due to elevated homocysteine is associated with an increased risk of both venous and arterial thrombosis. In the conversion of homocysteine to methionine, highly polymorphic gene sequences of the methylene tetrahydrofolate reductase enzyme (MTHFR) can slow the pathway, resulting in hyperhomocysteinemia. Polymorphisms influencing homocysteine metabolism occur in up to 20% of individuals in some populations.

### **The power of polymorphisms**

**G**enetic polymorphisms do not necessarily imply a disease state. These DNA changes are preserved in populations, and their effects on the proteins they code are variable, even of benefit in some populations. (That may explain the continued presence of polymorphisms despite their negative clinical impact in other

populations.) The effect of a genetic polymorphism on the quantitative or qualitative function of a protein is variable, and often occurs in combination with other genetic, physiologic, or environmental changes.

The frequency of a specific polymorphism varies from population to population, reflecting ancient adaptation to specific environments. Given the complex interactions required for modulation of the opposing thrombotic and antithrombotic processes of the clotting cascade, genetic polymorphisms affecting key receptors, enzymes, and cofactors all contribute to functional control of clotting.

Without this precise control, 1 mL of blood can convert the total body volume of fibrinogen to fibrin—and clot formation—in 10 to 15 seconds.<sup>2</sup> The process leading to the fibrin plugs of pathologic thrombi differs little from the appropriate hemostasis of a laceration, leading to the description of thrombosis as “fibrin plugs in the wrong place or at the wrong time.”<sup>2</sup>

The major inhibitors of thrombosis are antithrombin III (ATIII), protein C, and protein S, which form a complex to prevent excessive clotting. Typically, this complex binds with factors V and VIII, rendering them inactive and thus limiting the progression of clot formation. Loss of interaction between this antithrombotic complex (ATIII-protein S-protein C) and the clotting-cascade factors leads to unregulated progression of the clotting cascade and excessive thrombosis formation.

### **Polymorphisms identified in clotting abnormalities**

**Resistance to activated protein C.** Occurring in 2% to 5% of the general population, resistance to activated protein C (APC) results in loss of clot inhibition and increased thrombotic potential of the clotting cascade, as reflected in the striking 20% to 60% incidence of APC resistance among individuals with thrombotic disease. While APC resistance can be acquired, such as through antiphospholipid antibody syndrome, in the

CONTINUED

majority of cases an inherited polymorphism of factor V is responsible.

**Factor V Leiden.** Polymorphism factor V Leiden, a substitution of glutamine for arginine at amino acid 506 of factor V, exemplifies the role of DNA polymorphisms in clotting abnormalities. The Leiden polymorphism renders factor V resistant to interaction with protein C. Even with adequate qualitative and quantitative function of protein S, protein C, and ATIII, the antithrombotic regulatory complex is rendered ineffective by factor V resistance to protein C binding.

The relative risk of thrombosis in the presence of factor V Leiden follows a dose-response curve. In some families, heterozygotic carriers of factor V Leiden with 1 copy of the polymorphism have a 10-fold increased risk of venous thrombosis. Homozygotes with 2 aberrant copies face a 91-fold increase.


**Prothrombin 20210A.** Recognition of polymorphisms at other sites along the clotting cascade is emerging. The clotting-cascade step of conversion of prothrombin to thrombin is central to thrombus formation. A change of G to A at position 20210 in prothrombin (prothrombin 20210A) elevates baseline prothrombin levels and thrombin formation.

Heterozygotic carriers of prothrombin 20210A are 3 times more prevalent among individuals with venous thrombosis and 9 times more prevalent in individuals with familial thrombosis.

**Hyperhomocysteinemia.** A frequent condition in the general population, hyperhomocysteinemia is also an important contributor to the overall risk of thrombotic disease.<sup>3</sup> Although the exact mechanism for this is unclear, mild to moderate increases in homocysteine levels are associated with an increased relative risk of thrombosis (2.5).

**FIGURE**

**Factor V Leiden among women with habitual pregnancy loss**



Dietary restriction of folate and vitamin B<sub>12</sub> remains the most common cause.

However, in at least a quarter of cases, DNA polymorphisms of the homocysteine-methionine pathway are associated with hyperhomocysteinemia and a mild to moderate increase in thrombosis. The most common inherited cause, a C to T change at position 677 of MTHFR, produces a thermolabile enzyme with reduced catalytic capacity and lowered conversion of homocysteine. The diminished conversion of homocysteine to methionine is further slowed by folate deficiency.<sup>4</sup> Other MTHFR polymorphisms also may contribute to diminished catalytic capacity of this pathway, with resultant increased levels of homocysteine.<sup>5</sup>

**Polymorphisms identified in obstetric complications**

**Fetal loss and habitual miscarriage.** Among women who experience habitual miscarriage, factor V Leiden contributes primarily to the subgroup of losses that occur during the second trimester. The association between factor V Leiden and early first-trimester fetal loss is not as robust (FIGURE).<sup>6</sup>

CONTINUED

In the first prospective analysis of obstetric outcomes in carriers of factor V Leiden, infertility or miscarriage was 1.5 times greater than controls (95% confidence interval [CI], 1.2–2.7). In carriers who had experienced 2 or more fetal losses, factor V Leiden was 2.5 times greater than expected (95% CI, 1.2–5.13).<sup>7</sup> Conversely, however, among families with multiple individuals with thrombotic disease, habitual early losses are not more common.<sup>8</sup>

Limited evaluations of the prothrombin gene polymorphism likewise both support and refute an association with early or late fetal loss.<sup>9</sup> Given the relatively low incidence of late fetal loss with either polymorphism, the majority of women with factor V Leiden or a prothrombin polymorphism will have successful pregnancies.<sup>10</sup>

### A variety of studies support a link between mild to moderate hyperhomocysteinemia and preeclamptic toxemia.

With respect to the methionine-homocysteine pathway, both folate deficiency and a MTHFR polymorphism play a role—alone or in combination.<sup>6</sup> In fact, 1 study reported that hyperhomocysteinemia alone did not display a significant association with early fetal loss, suggesting that folate deficiency and MTHFR polymorphisms may operate through additional, unidentified variables. However, in a meta-analysis, hyperhomocysteinemia alone had a pooled risk of habitual pregnancy loss of 2.7 (95% CI, 1.4–5.2) and 4.2 (95% CI, 2.0–8.8) for fasting and postmethionine levels, respectively.<sup>11</sup> The relatively weak association with hyperhomocysteinemia, the diversity in definition of habitual pregnancy loss, and the variability inherent in homocysteine testing likely contribute to these conflicting findings.

**Preeclampsia.** Several studies have confirmed an association between APC resistance and preeclamptic toxemia. In fact, the

American College of Obstetricians and Gynecologists reports a 2.4-fold increase in factor V Leiden, a major determinant of APC resistance, among women with severe preeclamptic toxemia.<sup>12</sup> However, analyses in non-Caucasian populations typically do not identify factor V Leiden as a risk factor.<sup>13,14</sup>

An association between the prothrombin 20210A polymorphism and severe preeclampsia is variable and not confirmed in all populations. The lower background frequency of this polymorphism may limit a sufficient sample size for detection of a significant association.

With regard to homocysteine, a variety of studies support a link between mild to moderate elevations and preeclamptic toxemia. Homocysteine is mildly elevated during pregnancies complicated by preeclamptic toxemia, and this elevation persists postpartum.<sup>15,16</sup> Homocysteine elevations are also present in the second trimester before blood pressure increases occur.<sup>16</sup> Even at 15 weeks' gestation, an elevated homocysteine level indicates an almost 3-fold increased risk of severe preeclampsia.<sup>17</sup> Based on a meta-analysis of 5 studies, mild to moderate increases in homocysteine alone are significantly associated with a risk for severe preeclampsia, while folate deficiency alone is not (TABLE 1). The hyperhomocysteinemia is explained only in part by the increased rate of MTHFR polymorphism, with other contributors unidentified.<sup>18</sup>

**Placental abruption and infarct.** Hyperhomocysteinemia, folate deficiency, and MTHFR polymorphisms may also contribute to abruption and placental infarct. Hyperhomocysteinemia is 3 times more common in study patients versus controls (31% versus 9%), with fasting levels more significantly increased than postmethionine levels.<sup>19</sup> Likewise, MTHFR polymorphisms have been significantly associated with abruption and fetal growth restriction. The combination of heterozygosity for 2 MTHFR polymorphisms, C677T and A1298C, occurs in almost a quarter of placental abruptions.<sup>20</sup>

CONTINUED

**TABLE 1****Meta-analysis of the association between preeclampsia and selected metabolic defects<sup>6</sup>**

<b>METABOLIC DEFECT</b>	<b>ODDS RATIO</b>	<b>95% CI</b>
Folate deficiency	1.2	0.5–2.7
Hyperhomocysteinemia	20.9	3.6–121.6
Methylene tetrahydrofolate reductase polymorphisms	2.6	1.4–5.1

**Other contributors to placental thrombosis**

Several factors reflecting normal physiologic changes of pregnancy also contribute to the increased thrombotic risk when thrombophilia polymorphisms are present during pregnancy. The low vascular resistance of the placenta, key to the adequate perfusion of the fetus, conversely sets the stage for fibrin deposition and clot formation. The normal increase in the levels of fibrinogen and clotting proteins during pregnancy also contribute to clot formation. These physiologic changes of pregnancy may be modulated by diverse environmental changes. When the most common polymorphism of MTHFR is present, the rate of this enzyme's functioning is further impaired by low folate, a condition often present due to physiologic decreases of folate during pregnancy. Suppression of available folate associated with cigarette smoking may further compound the deleterious effects of the MTHFR polymorphism on homocysteine levels.

**Multiple thrombophilias increase risk**

In a similar additive fashion, an individual with multiple thrombophilia polymorphisms faces even greater increases in thrombotic risk. For example, factor V Leiden working concomitantly with protein C or protein S deficiency, or factor V Leiden or prothrombin 20210A paired with hyperhomocysteinemia carry relative risks of venous thrombosis

greater than any of these elements alone. For patients with a history of thrombotic disease from the European Prospective Cohort on Thrombophilia (EPCOT) study, the highest odds for a stillbirth (odds ratio [OR], 14.3; 95% CI, 2.4–86.0) occurred in women with combined thrombophilia abnormalities.<sup>8</sup>

Multiple inherited thrombophilias also may interact at the maternal-fetal interface. Consistent with Mendelian inheritance, the fetus will inherit 1 of the maternal alleles at each gene of the clotting-cascade proteins. Chronologically, the fetal arterial supply is established as maternal spiral arteries perfuse the intervillous spaces, with the maternal and fetal blood supply of the placenta present 3 to 4 weeks after conception. Histologically, evidence of placental ischemia can be found on either the maternal or fetal side. It is unknown whether the risk of placental compromise is greater in the presence of maternal or fetal thrombophilia, alone or in combination. Initial study of factor V Leiden from spontaneous miscarriages suggests a slight skewing toward increased fetal inheritance of the maternal polymorphism, suggesting a further contributory role of the fetus to overall risk.<sup>21</sup> Interestingly, MTHFR polymorphisms in combination (C677T and A1298C) occur significantly more often among spontaneous losses than in fetal bloods at delivery, suggesting a decreased potential for viability among such fetuses.<sup>22</sup> In populations with no association between preeclampsia and maternal factor V Leiden, the MTHFR polymorphism or

CONTINUED

**TABLE 2**

**Folate supplementation decreases homocysteine levels in women\*<sup>30</sup>**

TIME OF ASSESSMENT	MEAN FASTING HOMOCYSTEINE (UMOL/L)	95% CI	MEAN POST-LOAD HOMOCYSTEINE (UMOL/L)	95% CI
Presupplementation	16.6	13.0–20.1	68.5	60.8–76.2
Postsupplementation	6.1	5.6–6.7	29.3	25.6–33.0

CI=confidence interval  
 \*All subjects had a history of preeclampsia.

prothrombin polymorphism fetal studies are likewise unrevealing.<sup>23</sup>

**Promising results of enoxaparin, folate studies**

Inherited thrombophilias may be amenable to treatment with baby aspirin, heparin, folic acid, intravenous immunoglobulin G (IgG), or factor concentrates. Currently, no randomized, controlled intervention trials have been conducted. Further, since initial adverse pregnancy outcomes in inherited thrombophilia patients are relatively rare, it will be difficult to identify a sufficiently sizable population at risk.

For inherited thrombophilia carriers with recurrent adverse pregnancy outcomes, 2 areas for intervention deserve greater attention: enoxaparin treatment for women experiencing habitual late fetal loss<sup>24</sup> and folic acid treatment for severe preeclampsia.<sup>30</sup>

**Enoxaparin and recurrent fetal loss.** The Brenner enoxaparin trial examined 96 women who had experienced recurrent pregnancy loss (3 or more in the first trimester [7-12 weeks], 2 or more in the second trimester, or 1 in the third trimester). Only losses with a fetal pole previously documented by ultrasound were counted. Approximately 66.6% of the participants had a thrombophilia polymorphism identified.

Subsequent pregnancies among these women were treated with 40 mg enoxaparin daily (80 mg daily for women with combined abnormalities). A significant difference in pregnancy outcome was noted when these

patients were compared with their historical outcomes, although no differential in risk for solitary versus combined defects (83% versus 69%, *P*=.37) could be determined. Results appeared equally efficacious with either 40 mg or 80 mg enoxaparin a day.

Although these findings are encouraging, the use of historical controls for pregnancy outcome studies is notorious for overestimating the benefit of any single intervention. More complete analysis of the benefits of enoxaparin must await randomized controlled trials.

Several other studies have addressed the use of preconception aspirin (81 mg per day) and postconception unfractionated heparin among women with a history of habitual miscarriage. In 1 investigation involving 149 women whose recurrent miscarriages were due to a variety of procoagulant defects as well as factor V Leiden, fewer than 1% failed such therapy.<sup>25</sup> Among a smaller population (*n*=34), treatment with postconception heparin resulted in decreased fetal loss, although the drug had no effect on other obstetric complications.<sup>26</sup> In addition, among women with a combination of infertility, miscarriage, and thrombophilia, who experienced an 85% historical rate of early-pregnancy loss, that rate declined to 15% when either preconception enoxaparin (history of infertility) or post-conception enoxaparin (miscarriage alone) was administered.<sup>27</sup>

**Folic acid supplementation and preeclamptic toxemia.** Among all individuals—pregnant or not—folate supplementa-

CONTINUED

**TABLE 3****Folic acid supplementation in women with a history of severe preeclampsia<sup>30</sup>**

<b>TIME OF ASSESSMENT</b>	<b>GESTATIONAL AGE AT DELIVERY (WEEKS)</b>	<b>BIRTH WEIGHT (G)</b>
Presupplementation	29.5 ± 3.7	1,088 ± 570
Postsupplementation	36.7 ± 2.2	2,867 ± 648

tion significantly lowers homocysteine levels. Efficacy is greatest at the highest pretreatment homocysteine levels and least at the lowest pretreatment levels.<sup>28</sup>

Decreases in homocysteine levels are seen with 0.5 mg to 5 mg folic acid daily plus vitamin B<sub>6</sub>. No additional benefit is obtained with supplemental vitamin B<sub>12</sub>. Folate appears most beneficial in lowering homocysteine in the presence of MTHFR polymorphisms, as such individuals have higher homocysteine levels at given folate deficiencies than individuals without this polymorphism.<sup>29</sup>

Based on these observations, folic acid supplements have been examined for their role in preeclampsia and intrauterine growth retardation (IUGR).<sup>30</sup> In 1 study, women with a history of preeclampsia; hemolysis, elevated liver enzymes and low platelet count (HELLP) syndrome; or

eclampsia (n=181) and IUGR (n=26) were identified following delivery. Assessment in the nonpregnant state—by measurement of both fasting and post-methionine-load levels—indicated positive results (above the 97.5 percentile in healthy premenopausal females) in 17.7% of preeclamptic toxemia cases and 19.2% of IUGR cases.

Of this group, 27 women proceeded to daily supplementation with folate (5 mg) and vitamin B<sub>6</sub> (250 mg) for a minimum of 10 weeks and achieved significantly decreased homocysteine levels (TABLE 2). (Interestingly,

CONTINUED

analysis by fasting levels alone would have missed more than 50% of the hyperhomocysteinemic women; their latent metabolic insufficiencies were documented only following load to the enzymatic system with methionine.) The supplementation protocol was continued preconception in 14 eligible women (11 had been hospitalized for preeclamptic toxemia and 3 had a history of IUGR), with a baby aspirin added daily at 10 to 12 weeks. Although the recurrence rate (64%) was unaltered from the expected, disease onset occurred later in pregnancy, resulting in infants delivered at a significantly later gestational ages with better birth weights. These results anticipate reduced neonatal mortality and morbidity (TABLE 3).

### Summary

With rapid expansion of human genome sequencing and low-cost assays for the detection of DNA sequence changes, the panel of suspect polymorphisms will likely increase. Randomized, controlled studies are needed to optimize the treatment choices. ■

#### REFERENCES

1. Brenner B, Blumenfeld Z. Thrombophilia and fetal loss. *Blood Rev.* 1997;11(2):72-79.
2. Fauci AS, Braunwald E, Isselbacher KJ, Martin JB, eds. *Harrison's Principles of Internal Medicine.* 14th ed. 1998: 339-345.
3. den Heijer M, Koster T, Blom HJ, et al. Hyperhomocysteinemia as a risk factor for deep-vein thrombosis. *N Engl J Med.* 1996;334(12):759-762.
4. Frosst P, Blom HJ, Milos R, et al. A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. *Nat Genet.* 1995;10(1):111-113.
5. van der Put NM, Gabreels F, Stevens EM, et al. A second common mutation in the methylenetetrahydrofolate reductase gene: an additional risk factor for neural-tube defects? *Am J Hum Genet.* 1998;62(5):1044-1051.
6. Ray JG, Laskin CA. Folic acid and homocyst(e)ine metabolic defects and the risk of placental abruption, pre-eclampsia and spontaneous pregnancy loss: A systematic review. *Placenta.* 1999;20(7):519-529.
7. Bare SN, Poka R, Balogh I, et al. Factor V Leiden as a risk factor for miscarriage and reduced fertility. *Aust N Z J Obstet Gynaecol.* 2000;40(2):186-190.
8. Preston FE, Rosendaal FR, Walker ID, et al. Increased fetal loss in women with heritable thrombophilia. *Lancet.* 1996;348(9032):913-916.
9. Pihusch R, Buchholz T, Lohse P, et al. Thrombophilic gene mutations and recurrent spontaneous abortion: Prothrombin mutation increases the risk in the first trimester. *Am J Reprod Immunol.* 2001;46(2):124-131.
10. Ridker PM, Miletich JR, Buring JE, et al. Factor V Leiden mutation as a risk factor for recurrent pregnancy loss. *Ann Intern Med.* 1998;128(12 Pt 1):1000-1003.
11. Nelen WL, Blom HJ, Steegers EA, et al. Hyperhomocysteinemia and recurrent early pregnancy loss: a meta-analysis. *Fertil Steril.* 2000;74(6):1196-1199.
12. Dizon-Townson DS, Nelson LM, Easton K, et al. The factor V Leiden mutation may predispose women to severe preeclampsia. *Am J Obstet Gynecol.* 1996;175(4 Pt 1):902-905.

13. Zusterzeel PL, Visser W, Blom HJ, et al. Methylenetetrahydrofolate reductase polymorphisms in preeclampsia and the HELLP syndrome. *Hypertens Pregnancy.* 2000;19(3):299-307.
14. Kim YJ, Williamson RA, Murray JC, et al. Genetic susceptibility to preeclampsia: Roles of cytosine-thymine substitution at nucleotide 677 of the gene for methylenetetrahydrofolate reductase, 68-base pair insertion at nucleotide 844 of the gene for cystathionine beta-synthase, and factor V Leiden mutation. *Am J Obstet Gynecol.* 2001;184(6):1211-1217.
15. Dekker GA, DeVries JI, Doelitzsch PM, et al. Underlying disorders associated with severe early-onset preeclampsia. *Am J Obstet Gynecol.* 1995;173(4):1042-1048.
16. Rajkovic A, Mahomed K, Malinow MR, et al. Plasma homocyst(e)ine concentrations in eclamptic and preeclamptic African women postpartum. *Obstet Gynecol.* 1999;94(3):355-360.
17. Cotter AM, Molloy AM, Scott JM, et al. Elevated plasma homocysteine in early pregnancy: a risk factor for the development of severe preeclampsia. *Am J Obstet Gynecol.* 2001;185(4):781-785.
18. Lachmeijer AM, Arngrimsson R, Bastiaans EJ, et al. Mutations in the gene for methylenetetrahydrofolate reductase, homocysteine levels, and vitamin status in women with a history of preeclampsia. *Am J Obstet Gynecol.* 2001;184(3):394-402.
19. Goddijn-Wessel TA, Wouters MG, van de Molen EF, et al. Hyperhomocysteinemia: a risk factor for placental abruption or infarction. *Eur J Obstet Gynecol Reprod Biol.* 1996;66(1):23-29.
20. Gebhardt GS, Scholtz CL, Hillermann R, et al. Combined heterozygosity for methylenetetrahydrofolate reductase (MTHFR) mutations C677T and A1298C is associated with abruptio placentae but not with intrauterine growth restriction. *Eur J Obstet Gynecol Reprod Biol.* 2001;97(2):174-177.
21. Dizon-Townson DS, Meline L, Nelson LM, et al. Fetal carriers of the factor V Leiden mutation are prone to miscarriage and placental infarction. *Am J Obstet Gynecol.* 1997;177(2):402-405.
22. Isotalo PA, Wells GA, Donnelly JG. Neonatal and fetal methylenetetrahydrofolate reductase genetic polymorphisms: An examination of C677T and A1298C mutations. *Am J Hum Genet.* 2000;67(4):986-990.
23. Livingston JC, Barton JR, Park V, et al. Maternal and fetal inherited thrombophilias are not related to the development of severe preeclampsia. *Am J Obstet Gynecol.* 2001;185(1):153-157.
24. Brenner B, Hoffman R, Blumenfeld Z, et al. Gestational outcome in thrombophilic women with recurrent pregnancy loss treated by enoxaparin. *Thromb Haemost.* 2000;83(5):693-697.
25. Bick RL. Recurrent miscarriage syndrome due to blood coagulation protein/platelet defects: Prevalence, treatment and outcome results. DRW Metrex Recurrent Miscarriage Syndrome Cooperative Group. *Clin Appl Thromb Hemost.* 2000;6(3):115-125.
26. Ogueh O, Chen MF, Spurlin G, et al. Outcome of pregnancy in women with hereditary thrombophilia. *Int J Gynaecol Obstet.* 2001;74(3):247-253.
27. Sarto A, Rocha M, Geller M, et al. Treatment with enoxaparin adapted to the fertility programs in women with recurrent abortion and thrombophilia. *Medicina.* 2001;61(4):406-412.
28. Clarke R, Armitage J. Vitamin supplements and cardiovascular risk: Review of the randomized trials of homocysteine-lowering vitamin supplements. *Semin Thromb Hemost.* 2000;26(3):341-348.
29. Malinow MR, Nieto FJ, Kruger WD, et al. The effects of folic acid supplementation on plasma total homocysteine are modulated by multivitamin use and methylenetetrahydrofolate reductase genotypes. *Arterioscler Thromb Vasc Biol.* 1997;17(6):1157-1162.
30. Leeda M, Ryazi N, DeVries JI, et al. Effects of folic acid and vitamin B6 supplementation on women with hyperhomocysteinemia and a history of preeclampsia or fetal growth restriction. *Am J Obstet Gynecol.* 1998;179(1):135-139.
31. Grandone E, Margaglione M, Colaizzo D, et al. Factor V Leiden is associated with repeated and recurrent unexplained fetal losses. *Thromb Haemost.* 1997;77(5):822-824.
32. Rai R, Shlebak A, Cohen H, et al. Factor V Leiden and acquired activated protein C resistance among 1000 women with recurrent miscarriage. *Hum Reprod.* 2001;16(5):961-965.
33. Younis JS, Brenner B, Ohel G, Tal J, Lanir N, Ben-Ami M. Activated protein C resistance and factor V Leiden mutation can be associated with first—as well as second—trimester recurrent pregnancy loss. *Am J Reprod Immunol.* 2000;43(1):31-35.

*Dr. Wilkins-Haug reports no affiliations or financial arrangements with any of the manufacturers of products mentioned in this article.*