


# INSTANT POLL

## RESULTS

Here's how your peers voted

### Which tocolytic agent do you use most often to treat preterm labor?



#### COMMENTS

“Between 23 and 26 weeks, magnesium has been the most effective therapy in my practice”

“Many physicians see magnesium's efficacy each week. The Cochrane Database should not be read as Bible.”

From: January 2007 OBG MANAGEMENT


### “I want an IUD”— how would you respond?

Your 24-year-old patient reports 3 sexual partners over 6 months; they have used a condom “occasionally.” Physical exam is normal; a test for *C. trachomatis* is negative. She had a first dose of the HPV vaccine.

Because she has multiple partners and a significant risk of sexually transmitted infection, I would not insert an IUD

Because the IUD is so effective, I would insert one and counsel her to have her partners use a condom

Because she has not had a prior delivery, I would be reluctant to insert an IUD and prefer that she use an oral contraceptive and condoms for her partners



From: April 2007 OBG MANAGEMENT

## Can you prognosticate the **future** of the specialty?

Gazing into the future, which of the following “-ist” models do you think ObGyn practices are most likely to heavily rely on to boost the career satisfaction of practicing obstetrician-gynecologists?

- ☐ The *laborist* model
- ☐ The *nocturnalist* model
- ☐ The *weekendist* model
- ☐ I don't foresee major changes to the current model

**What do you say the future holds? Tell us at**  
[www.obgmanagement.com](http://www.obgmanagement.com)

Read what your peers predict when INSTANT POLL results are published in an upcoming issue