



BRIEF ANSWERS
TO SPECIFIC
CLINICAL
QUESTIONS

Breast-feeding is considered safe for renal transplant recipients taking prednisone, azathioprine, cyclosporine, and tacrolimus

1-MINUTE CONSULT

SILVI SHAH, MD

Division of Nephrology, University of Cincinnati,
Cincinnati, OH

Q: Is pregnancy safe after kidney transplant?

A: Since the first successful pregnancy in a kidney transplant recipient in 1958,¹ hundreds of kidney recipients have had successful pregnancies. Chronic kidney disease disrupts the hypothalamic-pituitary-gonadal axis, leading to anovulation and infertility. However, within 6 months of kidney transplant, the hypothalamic-pituitary-gonadal axis and sex hormone levels return to normal,² and the renal allograft is able to adapt to the various physiologic changes of pregnancy.³

Successful pregnancy after kidney transplant requires a team approach to care that includes the primary care physician, a transplant nephrologist, and an obstetrician with expertise in high-risk pregnancies. But equally important is educating and counseling the patient about the risks and challenges. This should begin at the first pretransplant visit.⁴

Below are answers to questions often asked by renal transplant recipients who wish to become pregnant.

■ WHAT IS THE IDEAL TIME TO BECOME PREGNANT AFTER KIDNEY TRANSPLANT?

According to American Society of Transplantation and European best-practice guidelines, as outlined in **Table 1**, the ideal time to conceive is 1 to 2 years after renal transplant if graft function is stable, proteinuria is minimal, there are no recent episodes of acute rejection, and the patient is not taking teratogenic medications. Because transplant recipients take teratogenic immunosuppressive drugs such as mycophenolate mofetil, women should be counseled to start contraception as soon as

possible after kidney transplant.^{5,6}

Mycophenolate mofetil and sirolimus are contraindicated in pregnancy and should be stopped at least 6 weeks before conception. Mycophenolate mofetil increases the risk of congenital malformations and spontaneous abortion. Data on sirolimus from clinical studies are limited, but in animal studies it is associated with delay in ossification of skeletal structure and with an increase in fetal mortality.⁷

■ WHAT INCREASES THE RISK OF A POOR PREGNANCY OUTCOME AFTER RENAL TRANSPLANT?

Risk factors for poor maternal and fetal outcomes include an elevated prepregnancy serum creatinine level (≥ 1.4 mg/dL), hypertension, and proteinuria (≥ 500 mg/24 hours). Younger age at transplant and at conception is associated with better pregnancy outcome.^{5,8}

■ WHAT ARE THE POSSIBLE MATERNAL COMPLICATIONS?

Kidney transplant recipients who become pregnant have a risk of developing preeclampsia 6 times higher than normal, and the incidence rate ranges between 24% and 38%.^{9,10} The risk of cesarean delivery is 5 times higher than in the general population, and the incidence rate is 43% to 64%.^{10,11}

Low-dose aspirin reduces the risk of preeclampsia and should be prescribed to all pregnant women who are kidney transplant recipients. Angiotensin-converting enzyme inhibitors are contraindicated due to the risk of teratogenic effects, ie, pulmonary hypoplasia and oligohydramnios.⁴

doi:10.3949/ccjm.84a.16047

■ WHAT ARE THE POSSIBLE FETAL COMPLICATIONS?

Women who become pregnant after kidney transplant are at greater risk of preterm delivery (40% to 60% higher risk), having a baby with low birth weight (42% to 46% higher risk), and intrauterine growth restriction (30% to 50% higher risk). But the risk of perinatal mortality is not increased in the absence of the above-mentioned risk factors.^{10,11}

■ DOES PREGNANCY INCREASE THE RISK OF GRAFT FAILURE?

Pregnancy does not increase the risk of allograft loss as long as the patient has a pre-pregnancy serum creatinine below 1.4 mg/dL, no hypertension, and urine protein excretion less than 500 mg/24 hours.¹²

■ WHAT CHANGES TO IMMUNE SUPPRESSION ARE REQUIRED BEFORE AND DURING PREGNANCY?

Careful management of immunosuppression is critical in renal transplant recipients before and during pregnancy because of the risks of teratogenicity and other adverse effects.

As stated above, mycophenolate mofetil and sirolimus are teratogenic and should be stopped 6 weeks before conception. The recommended maintenance immunosuppression during pregnancy includes calcineurin inhibitors (tacrolimus and cyclosporine), azathioprine, and low-dose prednisone.

A 20% to 25% increase in the dose of calcineurin inhibitor is required during pregnancy due to an increase in metabolic activity of cytochrome P450 and an increase in the volume of distribution.^{5,6,13} However, this dosing increase requires more frequent monitoring throughout the pregnancy to ensure the safest possible therapeutic levels.

■ REFERENCES

1. Murray JE, Reid DE, Harrison JH, Merrill JP. Successful pregnancies after human renal transplantation. *N Engl J Med* 1963; 269:341–343.
2. Saha MT, Saha HH, Niskanen LK, Salmela KT, Pasternack AI. Time course of serum prolactin and sex hormones following successful renal transplantation. *Nephron* 2002; 92:735–737.
3. Davison JM. The effect of pregnancy on kidney function in renal allograft recipients. *Kidney Int* 1985; 27:74–79.
4. Shah S, Verma P. Overview of pregnancy in renal transplant patients. *Int J Nephrol* 2016; 2016:4539342.
5. McKay DB, Josephson MA, Armenti VT, et al; Women's Health

TABLE 1

Criteria for pregnancy after renal transplant

- From 1 to 2 years after transplant
- Stable allograft function, creatinine < 1.4 mg/dL
- No recent episodes of acute rejection
- Blood pressure controlled to ≤ 140/90 mm Hg
- No proteinuria, or proteinuria ≤ 500 mg/24 hours
- Prednisone dosage ≤ 15 mg/day
- Azathioprine dosage ≤ 2 mg/kg/day
- Discontinue mycophenolate mofetil and sirolimus 6 weeks before conception

■ DOES PREGNANCY INCREASE THE RISK OF INFECTION?

Because of their immunosuppressed state, renal transplant recipients are prone to infection; the incidence rate of urinary tract infection is as high as 40% due to mild reflux and pregnancy-related dilation of ureters and collecting ducts.⁶ Women should be screened for urinary tract infection at every visit with urine dipstick testing and with urine culture every 4 weeks. Antibiotics such as nitrofurantoin, amoxicillin, and cephalexin are safe to treat urinary tract infection during pregnancy.⁶

■ IS BREAST-FEEDING SAFE IN RENAL TRANSPLANT RECIPIENTS?

Breast-feeding is considered safe for women with renal transplant who are on prednisone, azathioprine, cyclosporine, and tacrolimus. Women should avoid breast-feeding if they are taking mycophenolate mofetil, sirolimus, everolimus, or belatacept, as clinical data on safety are not adequate.¹⁴

Committee of the American Society of Transplantation. Reproduction and transplantation: report on the AST Consensus Conference on Reproductive Issues and Transplantation. *Am J Transplant* 2005; 5:1592–1599.

6. EBPG Expert Group on Renal Transplantation. European best practice guidelines for renal transplantation. Section IV: long-term management of the transplant recipient. IV.10. Pregnancy in renal transplant recipients. *Nephrol Dial Transplant* 2002; 17(suppl 4):50–55.
7. Armenti VT, Moitz MJ, Cardonick EH, Davison JM. Immunosuppression in pregnancy: choices for infant and maternal health. *Drugs* 2002; 62:2361–2375.

PREGNANCY AFTER TRANSPLANT

8. **Bramham K, Chusney G, Lee J, Lightstone L, Nelson-Piercy C.** Breastfeeding and tacrolimus: serial monitoring in breast-fed and bottle-fed infants. *Clin J Am Soc Nephrol* 2013; 8:563–567.
9. **Deshpande NA, James NT, Kucirka LM, et al.** Pregnancy outcomes in kidney transplant recipients: a systematic review and meta-analysis. *Am J Transplant* 2011; 11:2388–2404.
10. **Bramham K, Nelson-Piercy C, Gao H, et al.** Pregnancy in renal transplant recipients: a UK national cohort study. *Clin J Am Soc Nephrol* 2013; 8:290–298.
11. **Coscia LA, Constantinescu S, Moritz MJ, et al.** Report from the National Transplantation Pregnancy Registry (NTPR): outcomes of pregnancy after transplantation. *Clin Transpl* 2010: 65–85.
12. **Sibanda N, Briggs JD, Davison JM, Johnson RJ, Rudge CJ.** Pregnancy

after organ transplantation: a report from the UK transplant pregnancy registry. *Transplantation* 2007; 83:1301–1307.

13. **Kim H, Jeong JC, Yang J, et al.** The optimal therapy of calcineurin inhibitors for pregnancy in kidney transplantation. *Clin Transplant* 2015; 29:142–148.
14. **Constantinescu S, Pai A, Coscia LA, Davison JM, Moritz MJ, Armentani VT.** Breast-feeding after transplantation. *Best Pract Res Clin Obstet Gynaecol* 2014; 28:1163–1173.

ADDRESS: Silvi Shah, MD, Division of Nephrology, University of Cincinnati, Department of Internal Medicine, 231 Albert Sabin Way, Medical Sciences Building Room 6065, PO Box 670557, Cincinnati, OH 45267-0557; silvishah2108@gmail.com



R.J. Fasenmyer Center for Clinical Immunology

Basic and Clinical Immunology

Complimentary CME-certified
Webcast Series

Do you feel left behind by the rapid advances in laboratory immunology being applied to patient care?

IF SO, DON'T MISS THIS UNIQUE OPPORTUNITY TO GET UP TO SPEED IN BASIC AND CLINICAL IMMUNOLOGY.

As a shareholder in the evolving field of immune-based therapeutics, the **Basic and Clinical Immunology webcast series** is your source for education and training in this field.

Watch these webcasts

- Overview of Innate Immunity, the Microbiome, and the Integrated Immune Response
- T Cell Effector Function (TH1, TH2, TH17) and Cytokine Biology
- Antigen Presentation, T Cell Activation and Deactivation
- Multiple Sclerosis – Cell Trafficking and the Integrated Immune Response
- Tolerance and Autoimmunity – Lessons from Animal Models and Primary Immunodeficiency Syndromes
- Immunoglobulins and Complement in Health and Disease
- Bone Immunology – Ankylosing Spondylitis and Psoriatic Arthritis
- B Cell Biology and T Follicular Helper Cells – The Fundamentals

These activities have been approved for *AMA PRA Category 1 Credit*™.

View today! ccfcme.org/clinicimmun

