

## Jaspreet Jaura, MD; Gary Kelsberg, MD

Valley Family Medicine Residency, University of Washington at Valley Medical Center, Renton

## Sarah Safranek, MLIS

University of Washington Health Sciences Library, Seattle

#### DEPUTY EDITOR

## Jon Neher, MD Valley Family Medicine Residency, University of Washington at Valley Medical Center, Renton

# Q/Does vitamin D supplementation reduce asthma exacerbations?

## **EVIDENCE-BASED ANSWER**

YES, to some extent it does, and primarily in patients with low vitamin D levels. Supplementation reduces asthma exacerbations requiring systemic steroids by 30% overall in adults and children with mild-to-moderate asthma (number needed to treat [NNT] = 7.7). The outcome is driven by the effect in patients with vitamin D levels < 25 nmol/L (NNT = 4.3), however; supplementation doesn't decrease exacerbations in patients with higher levels. Supplementation also reduces, by a smaller amount (NNT = 26.3), the odds of exacerbations requiring emer-

gency department care or hospitalization (strength of recommendation [SOR]: **A**, meta-analysis of randomized controlled trials [RCTs]).

In children, vitamin D supplementation may also reduce exacerbations and improve symptom scores (SOR: **C**, low-quality RCTs).

Vitamin D doesn't improve forced expiratory volume in 1 second (FEV<sub>1</sub>) or standardized asthma control test scores. Also, it isn't associated with serious adverse effects (SOR: **A**, meta-analysis of RCTs).

## **Evidence summary**

A Cochrane systematic review of vitamin D for managing asthma performed meta-analyses on RCTs that evaluated several outcomes.¹ The review found improvement in the primary outcome of asthma exacerbations requiring systemic steroids, mainly in adult patients, and in the secondary outcomes of emergency department visits or hospitalization, in a mix of adults and children (TABLE¹-6).

Most participants had mild-to-moderate asthma; trials lasted 4 to 12 months. Vitamin D dosage regimens varied, with a median daily dose of 900 IU/d (range, 400-4000 IU/d). Six RCTs were rated high-quality, and 1 had unclear risk of bias.

# Supplementation reduced exacerbations in patients with low vitamin D levels

A subsequent (2017) systematic review and meta-analysis evaluating the primary outcome of exacerbations requiring steroids<sup>7</sup> included another study<sup>8</sup> (in addition to the 6 RCTs in the Cochrane review).

When researchers reanalyzed individual participant data from the trials in the Cochrane review, plus the additional RCT, to include baseline vitamin D levels, they found that vitamin D supplementation reduced exacerbations overall (NNT = 7.7) and in patients with low baseline vitamin D levels (25[OH] vitamin D < 25 nmol/L; 92 participants in 3 RCTs; NNT = 4.3) but not in patients with higher baseline levels (764 participants in 6 RCTs). Vitamin D supplementation reduced the asthma exacerbation rate in patients with low baseline vitamin D levels (0.19 vs 0.42 events per participant-year; P = .046).

## Smaller benefit found on ED visits and hospitalizations

The Cochrane review, with 2 RCTs with

Effect of Vitamin D supplementation on frequency and severity of asthma exacerbations in patients with mild-to-moderate persistent asthma

Outcome measured	Studies	Population (mean ages)	Intervention vitamin D dose vs placebo <sup>a</sup>	Result (significant results in bold)	Comment
Asthma exacerbations requiring systemic steroids	658 Adults (2 RCTs¹) 22 Children (1 RCT¹)	Adults (40-48 y) Children (1-15 y)	Adults: 2000-4000 IU/d  Children: 400-2000 IU/d	Overall: 0.28 vs 0.44 events/person-year	Low vitamin D = < 25 nmol/L
Asthma exacerbations requiring emergency department visit ± hospitalization	658 Adults (2 RCTs¹) 305 Children (5 RCTs²-6)	Adults (40-48 y) Children (1-18 y)	Adults: 2000-4000 IU/d  Children: 400-1200 IU/d	25/1000 vs 63/1000 events in adults; NNT = 26.3	

NNT, number needed to treat.

adults (n = 658)<sup>1</sup> and 5 RCTs with children (n = 305),<sup>2-6</sup> evaluated whether Vitamin D reduced the need for emergency department visits and hospitalization with asthma exacerbations; they found a smaller benefit (NNT = 26.3).

## Effects on FEV<sub>1</sub>, daily asthma symptoms, and serious adverse effects

Several RCTs included in the 2017 metaanalysis found no effect of vitamin D supplementation on  $\text{FEV}_1$ , daily asthma symptoms (evaluated with the standardized Asthma Control Test Score), or reported serious adverse events.  $^{2-6,9,10}$  No deaths occurred in any trial.

## Additional findings in children from lower-quality studies

A 2015 systematic review and meta-analysis of RCTs evaluating vitamin D supplementation for children with asthma found<sup>11</sup>:

- moderate-quality evidence for decreased emergency department visits
   (1 RCT from India, 100 children ages 3 to 14 years, decrease not specified; *P* = .015);
- low-quality evidence for reduced exacerbations (6 RCTs [3 RCTs also in Cochrane review], 507 children ages 3 to 17 years; risk ratio = 0.41; 95% confidence interval, 0.27-0.63); and
- low-quality evidence for reduced standardized asthma symptom scores

(6 RCTs [2 RCTs also in Cochrane review], 231 children ages 3 to 17 years; amount of reduction not listed; P = .01).

## **Recommendations**

No published guidelines discuss using vitamin D in managing asthma. An American Academy of Family Physicians (AAFP) summary of the Cochrane systematic review recommends that family physicians await further studies and updated guidelines before recommending vitamin D for patients with asthma. <sup>12</sup> The AAFP also points out that the Endocrine Society has recommended vitamin D supplementation for adults (1500-2000 IU/d) and children (at least 1000 IU/d) at risk for deficiency.

## **Editor's takeaway**

In the meta-analyses highlighted here, researchers evaluated asthma patients with a wide range of ages, baseline vitamin D levels, and vitamin D supplementation protocols. Although vitamin D reduced asthma exacerbations requiring steroids overall, the effect was driven by 3 studies of patients with low baseline vitamin D levels. As a result, disentangling who might benefit the most remains a challenge. The conservative course for now is to manage asthma according to cur-

<sup>&</sup>lt;sup>a</sup>Additional Vitamin D dosing details: 400 IU/d,<sup>2</sup> 1000 IU/wk,<sup>3</sup> 500 IU/d,<sup>4</sup> 800 IU/d,<sup>5</sup> 1200 IU/d.<sup>6</sup>

rent guidelines and supplement vitamin D in patients at risk for, or with known, deficiency.

#### References

- Martineau AR, Cates CJ, Urashima M, et al. Vitamin D for the management of asthma. Cochrane Database Syst Rev. 2016;9:CD011511.
- Jensen M, Mailhot G, Alos N, et al. Vitamin D intervention in preschoolers with viral-induced asthma (DIVA): a pilot randomised controlled trial. *Trials*. 2016;26:17:353.
- Majak P, Jerzyiska J, Smejda K, et al. Correlation of vitamin D with Foxp3 induction and steroid-sparing effect of immunotherapy in asthmatic children. Ann Allergy Asthma Immunol. 2012;109: 329-335.
- Majak P, Olszowiec-Chlebna M, Smejda K, et al. Vitamin D supplementation in children may prevent asthma exacerbation triggered by acute respiratory infection. J Allergy Clin Immunol. 2011;127:1294-1296.
- Tachimoto H, Mezawa H, Segawa T, et al. Improved control of childhood asthma with low-dose, short-term vitamin D supple-

- mentation: a randomized, double-blind, placebo-controlled trial. *Allergy*. 2016;71:1001-1009.
- Urashima M, Segawa T, Okazaki M, et al. Randomized trial of vitamin D supplementation to prevent seasonal influenza A in school children. Am J Clin Nutr. 2010;91:1255-1260.
- Joliffe DA, Greenberg L, Hooper RL, et al. Vitamin D supplementation to prevent asthma exacerbations: a systematic review and meta-analysis of individual participant data. *Lancet* 2017;5:881-890.
- Kerley CP, Hutchinson K, Cormical L, et al. Vitamin D3 for uncontrolled childhood asthma: a pilot study. *Pediatr Allergy Immunol*. 2016;27:404-412.
- Castro M, King TS, Kunselman SJ, et al. Effect of vitamin D3 on asthma treatment failures in adults with symptomatic asthma and lower vitamin D levels: the VIDA randomized clinical trial. JAMA. 2014;311:2083-2091.
- Martineau AR, MacLaughlin BD, Hooper RL, et al. Double-blind multi-centre randomised controlled trial of vitamin D3 supplementation in adults with inhaled corticosteroid-treated asthma (ViDiAs). *Thorax*. 2015:70:451-457.
- Riverin B, Maguire J, Li P. Vitamin D supplementation for childhood asthma: a systematic review and meta-analysis. PLOS One. 2015;10:e0136841.
- 12. Qiu J. Vitamin D for the management of asthma. Am Fam Phys. 2017;96:290-293.