

Appendix Table 3. Sensitivity Analysis of Episode Spending by Components

All Episodes				
	Episode Component, %	Low SNF Cost (n=1,333)	High SNF Cost (n=445)	p-value
	Hospitalization	82.2	63.6	<0.001
	SNF	9.7	28.4	
	Other	8.1	8.0	
Individual Episode Types				
Episode Type	Episode Component, \$	Other Spending (n=159)	High Spending (n=52)	p-value
Aortic Aneurysm	Hospitalization	34,094	45,762	<0.001
	SNF	611	2,033	
	Other	1,783	3,806	
		Other Spending (n=628)	High Spending (n=208)	
Cellulitis	Hospitalization	6,651	7,303	<0.001
	SNF	1,509	3,293	
	Other	1,007	1,699	
		Other Spending (n=356)	High Spending (n=118)	
Cholecystectomy	Hospitalization	12,975	14,591	<0.001
	SNF	619	1,891	
	Other	1,000	2,015	

		Other Spending (n=1,000)	High Spending (n=333)	
Gastrointestinal Hemorrhage	Hospitalization	8,293	9,355	<0.001
	SNF	1,078	2,442	
	Other	968	1,826	
		Other Spending (n=1,085)	High Spending (n=358)	
Kidney/UTI	Hospitalization	6,194	6,772	<0.001
	SNF	2,590	4,881	
	Other	601	1,021	
		Other Spending (n=582)	High Spending (n=194)	
Spinal Fusion	Hospitalization	30,913	39,674	<0.001
	SNF	872	2,861	
	Other	2,079	4,291	

