

Doripenem Eases Ventilator-Related Pneumonia

BY BRUCE JANCIN

SAN DIEGO — The use of doripenem as initial therapy in patients with ventilator-associated pneumonia shortened the duration of mechanical ventilation and hospital length of stay, according to a pooled analysis of the two randomized trials that compared the drug with imipenem or piperacillin/tazobactam.

The explanation for the more efficient

use of medical resources in the doripenem-treated group may lie in the broad-spectrum carbapenem's robust activity against *Pseudomonas aeruginosa*, the most common baseline pathogen identified in the 625 participants in the two studies, Sanjay Merchant, Ph.D., said at the annual meeting of the American College of Chest Physicians.

P. aeruginosa was eradicated by doripenem (Doribax) in 67% of affected

patients, compared with 42% of those on the comparator agents. The minimum inhibitory concentration (MIC) 90 for doripenem was just 4 mcg/mL, one-fourth that of imipenem. The median duration of mechanical ventilation in the subgroup with *P. aeruginosa* ventilator-associated pneumonia (VAP) was 7 days with doripenem and 13 days with the other agents, reported Dr. Merchant of Johnson & Johnson, Raritan, N.J.

In the pooled analysis, the median duration of mechanical ventilation in the overall population was 7 days in the 312 doripenem-treated patients, compared with 10 days with imipenem or piperacillin/tazobactam (Zosyn). The median hospital length of stay was also significantly shorter in the doripenem group, at 22 versus 26 days.

These advantages for doripenem remained significant in a Cox regression analysis with adjustment for baseline APACHE II scores and other potential confounders.

In-hospital mortality was about 16% regardless of the antibiotic used.

Intravenous doripenem was dosed in the two trials at 500 mg every 8 hours. Imipenem was dosed at 500 mg every 6 hours or 1,000 mg every 8 hours, while piperacillin/tazobactam was given on a schedule of 4.5 g every 6 hours.

Doripenem is approved for the treatment of complicated urinary tract or intra-abdominal infections and is under Food and Drug Administration review for an indication for VAP and other nosocomial pneumonias.

Disclosures: This analysis was funded by Johnson & Johnson.

In Respiratory Failure, RA May Boost Mortality

SAN DIEGO — Much has been made of late of rheumatoid arthritis patients' substantially reduced life expectancy because of cardiovascular disease. Far less widely known is that they also have increased in-hospital mortality following acute respiratory failure.

In a study of 22,121 adults in the U.S. Nationwide Inpatient Sample database who had an emergency hospitalization for acute respiratory failure in 2003-2006, the 1,621 with comorbid rheumatoid arthritis or collagen vascular disease had an adjusted 21% increased risk of in-hospital mortality, Dr. David S. Kountz said at the annual meeting of the American College of Chest Physicians.

Unadjusted in-hospital mortality rates were 24.9% in the rheumatologic group and 24.3% in the 20,500 patients without such comorbidity. But the rheumatoid arthritis/collagen vascular disease group was younger, wealthier, and far more likely to be female.

After adjustment for these and other potential confounders, such as hospital teaching status, in a multivariate logistic regression analysis, the rheumatologic group had significantly greater in-hospital mortality, according to Dr. Kountz of Jersey Shore University Medical Center, Neptune, N.J.

—Bruce Jancin

values than many full-term neonates and older infants. Therefore, these pre-term neonates should be initiated with a dosing regimen of 10 mg/kg q12h. Consideration may be given to the use of a 10 mg/kg q8h regimen in neonates with a sub-optimal clinical response. All neonatal patients should receive 10 mg/kg q8h by 7 days of life. In limited clinical experience, 5 out of 6 (83%) pediatric patients with infections due to Gram-positive pathogens with MICs of 4 µg/mL treated with ZYVOX had clinical cures. However, pediatric patients exhibit wider variability in linezolid clearance and systemic exposure (AUC) compared with adults. In pediatric patients with a sub-optimal clinical response, particularly those with pathogens with MIC of 4 µg/mL, lower systemic exposure, site and severity of infection, and the underlying medical condition should be considered when assessing clinical response. **Geriatric Use** Of the 2046 patients treated with ZYVOX in Phase 3 comparator-controlled clinical trials, 589 (29%) were 65 years or older and 253 (12%) were 75 years or older. No overall differences in safety or effectiveness were observed between these patients and younger patients.

ADVERSE REACTIONS Adult Patients The safety of ZYVOX formulations was evaluated in 2046 adult patients enrolled in seven Phase 3 comparator-controlled clinical trials, who were treated for up to 28 days. In these studies, 85% of the adverse events reported with ZYVOX were described as mild to moderate in intensity. The incidence (%) of adverse events reported in at least 2% of patients treated with either ZYVOX (n=2046) or all comparators* (n=2001) in these trials were as follows: diarrhea 8.5 and 6.3; headache 6.5 and 5.5; nausea 6.2 and 4.6; vomiting 3.7 and 2.0; insomnia 2.5 and 1.7; constipation 2.2 and 2.1; rash 2.0 and 2.2; dizziness 2.0 and 1.9; and fever 1.6 and 2.1 respectively. The most common adverse events in patients treated with ZYVOX were diarrhea (incidence across studies: 2.8% to 11.0%), headache (incidence across studies: 0.5% to 11.3%), and nausea (incidence across studies: 3.4% to 9.6%). The percent of drug-related adverse events in at least 1% of adult patients in a trial involving the treatment of uncomplicated skin and skin structure infection comparing ZYVOX 400 mg q12h (n=548) to clarithromycin 250 mg q12h (n=537) were 25.4 and 19.6 respectively. The percent of patients discontinuing drug due to drug-related adverse events† were 3.5 and 2.4 respectively. The incidence of drug-related adverse events occurring in >1% of adult patients were diarrhea 5.3 and 4.8; nausea 3.5 and 3.5; headache 2.7 and 2.2; taste alteration 1.8 and 2.0; vaginal moniliasis 1.6 and 1.3; fungal infection 1.5 and 0.2; abnormal liver function tests 0.4 and 0.0; vomiting 0.9 and 0.4; tongue discoloration 1.1 and 0.0; dizziness 1.1 and 1.5; and oral moniliasis 0.4 and 0.0 respectively. The percent of drug-related adverse events in at least 1% of adult patients in all other indications of ZYVOX 600 mg q12h (n=1498) versus all other comparators‡ (n=1464) with at least 1 drug-related adverse event were 20.4 and 14.3 respectively. The percent of adult patients discontinuing due to drug-related adverse events† was 2.1 and 1.7 respectively. The incidence of drug-related adverse events occurring in >1% of adult patients were diarrhea 4.0 and 2.7; nausea 3.3 and 1.8; headache 1.9 and 1.0; taste alteration 0.9 and 0.2; vaginal moniliasis 1.0 and 0.4; fungal infection 0.1 and <0.1; abnormal liver function tests 1.3 and 0.5; vomiting 1.2 and 0.4; tongue discoloration 0.2 and 0.0; dizziness 0.4 and 0.3; and oral moniliasis 1.1 and 0.4. Other adverse events reported in Phase 2 and Phase 3 studies included oral moniliasis, vaginal moniliasis, hypertension, dyspepsia, localized abdominal pain, pruritus, and tongue discoloration. **Pediatric Patients** The safety of ZYVOX formulations was evaluated in 215 pediatric patients ranging in age from birth through 11 years, and in 248 pediatric patients aged 5 through 17 years (146 of these 248 were age 5 through 11 and 102 were age 12 to 17). These patients were enrolled in two Phase 3 comparator-controlled clinical trials and were treated for up to 28 days. In these studies, 83% and 99%, respectively, of the adverse events reported with ZYVOX were described as mild to moderate in intensity. In the study of hospitalized pediatric patients (birth through 11 years) with Gram-positive infections, who were randomized 2 to 1 (linezolid:vancomycin), mortality was 6.0% (13/215) in the linezolid arm and 3.0% (3/101) in the vancomycin arm. However, given the severe underlying illness in the patient population, no causality could be established. The incidence of adverse events reported in ≥2% of pediatric patients treated for uncomplicated skin and skin structure infections§ with ZYVOX (n=248) or cefadroxil (n=251) were fever 2.9 and 3.6; diarrhea 7.8 and 8.0; vomiting 2.9 and 6.4; rash 1.6 and 1.2; headache 6.5 and 4.0; upper respiratory infection 3.7 and 5.2; nausea 3.7 and 3.2; trauma 3.3 and 4.8; pharyngitis 2.9 and 1.6; cough 2.4 and 4.0; generalized abdominal pain 2.4 and 2.8; localized abdominal pain 2.4 and 2.8; loose stools 1.6 and 0.8; localized pain 2.0 and 1.6; skin disorder 2.0 and 0.0 respectively. The incidence of adverse events reported in ≥2% of pediatric patients treated for all other indications¶ with either ZYVOX (n=215) or vancomycin (n=101) in comparator-controlled trials were fever 14.1 and 14.1; diarrhea 10.8 and 12.1; vomiting 9.4 and 9.1; sepsis 8.0 and 7.1; rash 7.0 and 15.2; headache 0.9 and 0.0; anemia 5.6 and 7.1; thrombocytopenia 4.7 and 2.0; upper respiratory infection 4.2 and 1.0; nausea 1.9 and 0.0; dyspnea 3.3 and 1.0; reaction at site of injection or of vascular catheter 3.3 and 5.1; trauma 2.8 and 2.0; pharyngitis 0.5 and 1.0; convulsion 2.8 and 2.0; hypokalemia 2.8 and 3.0; pneumonia 2.8 and 2.0; thrombocytopenia 2.8 and 2.0; cough 0.9 and 0.0; generalized abdominal pain 0.9 and 2.0; localized abdominal pain 0.5 and 1.0; apnea 2.3 and 2.0; gastrointestinal bleeding 2.3 and 1.0; generalized edema 2.3 and 1.0; loose stools 2.3 and 3.0; localized pain 0.9 and 0.0; and skin disorder 0.9 and 1.0. The percent of pediatric patients treated for uncomplicated skin and skin structure infections§ with either ZYVOX (n=248) or cefadroxil (n=251) and with ≥1 drug-related adverse event occurring in more than 1% of patients were 19.2 and 14.1 respectively. The percent of pediatric patients discontinuing due to a drug-related adverse event was 1.6 and 2.4 respectively. The incidence of drug-related adverse events reported in more than 1% of pediatric patients (and more than 1 patient) were diarrhea 5.7 and 5.2; nausea 3.3 and 2.0; headache 2.4 and 0.8; loose stools 1.2 and 0.8; vomiting 1.2 and 2.4; generalized abdominal pain 1.6 and 1.2; localized abdominal pain 1.6 and 1.2; eosinophilia 0.4 and 0.4; rash 0.4 and 1.2; vertigo 1.2 and 0.4 and pruritus at non-application site 0.4 and 0.0 respectively. The percent of pediatric patients treated for all other indications¶ with either ZYVOX (n=215) or vancomycin (n=101) and with ≥1 drug-related adverse event occurring in more than 1% of patients were 18.8 and 34.3 respectively. The percent of patients discontinuing due to a drug-related adverse event were 0.9 and 6.1 respectively. The incidence of drug-related adverse events reported in more than 1% of pediatric patients (and more than 1 patient) were diarrhea 3.8 and 6.1; nausea 1.4 and 0.0; loose stools 1.9 and 0.0; thrombocytopenia 1.9 and 0.0; vomiting 1.9 and 1.0; anemia 1.4 and 1.0; eosinophilia 1.4 and 0.0; rash 1.4 and 7.1; oral moniliasis 0.9 and 4.0; fever 0.5 and 3.0; pruritus at non-application site 0.0 and 2.0; and anaphylaxis 0.0 and 10.1* respectively. **Laboratory Changes** ZYVOX has been associated with thrombocytopenia when used in doses up to and including 600 mg every 12 hours for up to 28 days. In Phase 3 comparator-controlled trials, the percentage of adult patients who developed a substantially low platelet count (defined as less than 75% of lower limit of normal and/or baseline) was 2.4% (range among studies: 0.3 to 10.0%) with ZYVOX and 1.5% (range among studies: 0.4 to 7.0%) with a comparator. In a study of hospitalized pediatric patients ranging in age from birth through 11 years, the percentage of patients who developed a substantially low platelet count (defined as less than 75% of lower limit of normal and/or baseline) was 12.9% with ZYVOX and 13.4% with vancomycin. In an outpatient study of pediatric patients aged from 5 through 17 years, the percentage of patients who developed a substantially low platelet count was 0% with ZYVOX and 0.4% with cefadroxil. Thrombocytopenia associated with the use of ZYVOX appears to be dependent on duration of therapy, (generally greater than 2 weeks of treatment). The platelet counts for most patients returned to the normal range/baseline during the follow-up period. No related clinical adverse events were

identified in Phase 3 clinical trials in patients developing thrombocytopenia. Bleeding events were identified in thrombocytopenic patients in a compassionate use program for ZYVOX; the role of linezolid in these events cannot be determined (see **WARNINGS**). Changes seen in other laboratory parameters, without regard to drug relationship, revealed no substantial differences between ZYVOX and the comparators. These changes were generally not clinically significant, did not lead to discontinuation of therapy, and were reversible. The percent of adult patients with at least one substantially abnormal hematologic** value in patients treated with ZYVOX 400 mg q12h or clarithromycin 250 mg q12h for uncomplicated skin and skin structure infections were as follows: hemoglobin (g/dL) 0.9 and 0.0; platelet count (x 10³/mm³) 0.7 and 0.8; WBC (x 10³/mm³) 0.2 and 0.6; neutrophils (x 10³/mm³) 0.0 and 0.2 respectively. The percent of adult patients with at least one substantially abnormal hematologic** value in patients treated with ZYVOX 600 mg q12h or a comparator‡ were as follows: hemoglobin (g/dL) 7.1 and 6.6; platelet count (x 10³/mm³) 3.0 and 1.8; WBC (x 10³/mm³) 2.2 and 1.3 and neutrophils (x 10³/mm³) 1.1 and 1.2 respectively. The percent of adult patients with at least one substantially abnormal serum chemistry†† value in patients treated with ZYVOX 400 mg q12h or clarithromycin 250 mg q12h for uncomplicated skin and skin structure infections were as follows: AST (U/L) 1.7 and 1.3; ALT (U/L) 1.7 and 1.7; LDH (U/L) 0.2 and 0.2; alkaline phosphatase (U/L) 0.2 and 0.2; lipase (U/L) 2.8 and 2.6; amylase (U/L) 0.2 and 0.2; total bilirubin (mg/dL) 0.2 and 0.0; BUN (mg/dL) 0.2 and 0.0; and creatinine (mg/dL) 0.2 and 0.0 respectively. The percent of adult patients with at least one substantially abnormal serum chemistry†† value in patients treated with ZYVOX 600 mg q12h or a comparator‡ were as follows: AST (U/L) 5.0 and 6.8; ALT (U/L) 9.6 and 9.3; LDH (U/L) 1.8 and 1.5; alkaline phosphatase (U/L) 3.5 and 3.1; lipase (U/L) 4.3 and 4.2; amylase (U/L) 2.4 and 2.0; total bilirubin (mg/dL) 0.9 and 1.1; BUN (mg/dL) 2.1 and 1.5; and creatinine (mg/dL) 0.2 and 0.6 respectively. The percent of pediatric patients with at least one substantially abnormal hematologic** value in patients treated with ZYVOX or cefadroxil for uncomplicated skin and skin structure infections§ were as follows: hemoglobin (g/dL) 0.0 and 0.0; platelet count (x 10³/mm³) 0.0 and 0.4; WBC (x 10³/mm³) 0.8 and 0.8; neutrophils (x 10³/mm³) 1.2 and 0.8 respectively. The percent of pediatric patients with at least one substantially abnormal hematologic** value in patients treated with ZYVOX or vancomycin for any other indication¶ were as follows: hemoglobin (g/dL) 15.7 and 12.4; platelet count (x 10³/mm³) 12.9 and 13.4; WBC (x 10³/mm³) 12.4 and 10.3 and neutrophils (x 10³/mm³) 5.9 and 4.3 respectively. The percent of pediatric patients with at least one substantially abnormal serum chemistry‡ value in patients treated with ZYVOX or cefadroxil for uncomplicated skin and skin structure infections§ were as follows: ALT (U/L) 0.0 and 0.0; lipase (U/L) 0.4 and 1.2; and creatinine (mg/dL) 0.4 and 0.0 respectively. The percent of pediatric patients with at least one substantially abnormal serum chemistry‡ value in patients treated with ZYVOX or vancomycin for any other indication¶ were as follows: ALT (U/L) 10.1 and 12.5; amylase (U/L) 0.6 and 1.3; total bilirubin (mg/dL) 6.3 and 5.2; and creatinine (mg/dL) 2.4 and 1.0 respectively. **Postmarketing Experience** Myelosuppression (including anemia, leukopenia, pancytopenia, and thrombocytopenia) has been reported during postmarketing use of ZYVOX (see **WARNINGS**). Peripheral neuropathy, and optic neuropathy sometimes progressing to loss of vision, have been reported in patients treated with ZYVOX. Lactic acidosis has been reported with the use of ZYVOX (see **PRECAUTIONS**). Although these reports have primarily been in patients treated for longer than the maximum recommended duration of 28 days, these events have also been reported in patients receiving shorter courses of therapy. Serotonin syndrome has been reported in patients receiving concomitant serotonergic agents, including antidepressants such as selective serotonin reuptake inhibitors (SSRIs) and ZYVOX (see **PRECAUTIONS**). Convulsions have been reported with the use of ZYVOX (see **PRECAUTIONS**). Anaphylaxis, angioedema, and bullous skin disorders such as those described as Stevens Johnson syndrome have been reported. These events have been chosen for inclusion due to either their seriousness, frequency of reporting, possible causal connection to ZYVOX, or a combination of these factors. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made and causal relationship cannot be precisely established. **OVERDOSAGE** In the event of overdosage, supportive care is advised, with maintenance of glomerular filtration. Hemodialysis may facilitate more rapid elimination of linezolid. In a Phase 1 clinical trial, approximately 30% of a dose of linezolid was removed during a 3-hour hemodialysis session beginning 3 hours after the dose of linezolid was administered. Data are not available for removal of linezolid with peritoneal dialysis or hemoperfusion. Clinical signs of acute toxicity in animals were decreased activity and ataxia in rats and vomiting and tremors in dogs treated with 3000 mg/kg/day and 2000 mg/kg/day, respectively.

* MDRSP refers to isolates resistant to 2 or more of the following antibiotics: penicillin, second-generation cephalosporins, macrolides, tetracycline, and trimethoprim/sulfamethoxazole.

† Comparators included cefepodoxime proxetil 200 mg PO q12h; ceftriaxone 1 g IV q12h; clarithromycin 250 mg PO q12h; dicloxacillin 500 mg PO q6h; oxacillin 2 g IV q6h; vancomycin 1 g IV q12h.

‡ The most commonly reported drug-related adverse events leading to discontinuation in patients treated with ZYVOX were nausea, headache, diarrhea, and vomiting.

§ Comparators included cefepodoxime proxetil 200 mg PO q12h; ceftriaxone 1 g IV q12h; dicloxacillin 500 mg PO q6h; oxacillin 2 g IV q6h; vancomycin 1 g IV q12h.

¶ Patients 5 through 11 years of age received ZYVOX 10 mg/kg PO q12h or cefadroxil 15 mg/kg PO q12h. Patients 12 years or older received ZYVOX 600 mg PO q12h or cefadroxil 500 mg PO q12h.

** Patients from birth through 11 years of age received ZYVOX 10 mg/kg IV/PO q8h or vancomycin 10 to 15 mg/kg IV q6-24h, depending on age and renal clearance.

†† These reports were of 'red-man syndrome,' which were coded as anaphylaxis.

** <75% (<50% for neutrophils) of Lower Limit of Normal (LLN) for values normal at baseline; <75% (<50% for neutrophils) of LLN and of baseline for values abnormal at baseline.

†† >2 x Upper Limit of Normal (ULN) for values normal at baseline; >2 x ULN and >2 x baseline for values abnormal at baseline.

‡† <75% (<50% for neutrophils) of Lower Limit of Normal (LLN) for values normal at baseline; <75% (<50% for neutrophils) of LLN and <75% (<50% for neutrophils, <90% for hemoglobin) if baseline <LLN) of baseline for values abnormal at baseline.

§ >2 x Upper Limit of Normal (ULN) for values normal at baseline; >2 x ULN and >2 (<1.5 for total bilirubin) x baseline for values abnormal at baseline.

Rx only

Rev. May 2008