

REMICADE-maintenance experienced elevations in ALT at >1 to <3 times the ULN compared to 34% of patients treated with placebo-maintenance. ALT elevations  $\geq 3$  times the ULN were observed in 5% of patients who received REMICADE-maintenance compared with 4% of patients who received placebo-maintenance. ALT elevations  $\geq 5$  times ULN were observed in 2% of patients who received REMICADE-maintenance compared to none in patients treated with placebo-maintenance. In UC clinical trials (median follow up 30 weeks. Specifically, the median duration of follow-up was 30 weeks for placebo and 31 weeks for REMICADE.), 17% of patients receiving REMICADE experienced elevations in ALT at >1 to <3 times the ULN compared to 12% of patients treated with placebo. ALT elevations  $\geq 3$  times the ULN were observed in 2% of patients who received REMICADE compared with 1% of patients who received placebo. ALT elevations  $\geq 5$  times ULN were observed in <1% of patients in both REMICADE and placebo groups. In an AS clinical trial (median follow up 24 weeks for placebo group and 102 weeks for REMICADE group) 51% of patients receiving REMICADE experienced elevations in ALT at >1 to <3 times the ULN compared to 15% of patients treated with placebo. ALT elevations  $\geq 3$  times the ULN were observed in 10% of patients who received REMICADE compared to none in patients who received placebo. ALT elevations  $\geq 5$  times ULN were observed in 4% of patients who received REMICADE compared to none in patients treated with placebo. In a PsA clinical trial (median follow up 39 weeks for REMICADE group and 18 weeks in placebo group) 50% of patients receiving REMICADE experienced elevations in ALT at >1 to <3 times the ULN compared to 16% of patients treated with placebo. ALT elevations  $\geq 3$  times the ULN were observed in 7% of patients who received REMICADE compared to none in patients who received placebo. ALT elevations  $\geq 5$  times ULN were observed in 2% of patients who received REMICADE compared to none in patients treated with placebo. In PsO clinical trials, (ALT values are obtained in 2 phase 3 psoriasis studies with median follow-up of 50 weeks for REMICADE and 16 weeks for placebo). 49% of patients receiving REMICADE experienced elevations in ALT at >1 to <3 times the ULN compared to 24% of patients treated with placebo. ALT  $\geq 3$  x ULN were observed in 8% of patients who received REMICADE compared to <1 % who received placebo. ALT elevations  $\geq 5$  x ULN were observed in 3% of patients who received REMICADE compared to none in patients treated with placebo. **Adverse Reactions in Pediatric Crohn's Disease** There were some differences observed in the adverse reactions observed in the pediatric patients receiving REMICADE compared to those observed in adults with CD. The following adverse events were reported more commonly in 103 randomized pediatric CD patients administered 5 mg/kg REMICADE through 54 weeks than in 385 adult CD patients receiving a similar treatment regimen: anemia (11%), blood in stool (10%), leukopenia (9%), flushing (9%), viral infection (8%), neutropenia (7%), bone fracture (7%), bacterial infection (6%), and respiratory tract allergic reaction (6%). Infections were reported in 56% of randomized pediatric patients in Study Peds Crohn's and in 50% of adult patients in Study Crohn's I. In Study Peds Crohn's, infections were reported more frequently for patients who received every 8 week as opposed to every 12 week infusions (74% and 38%, respectively), while serious infections were reported for 3 patients in the every 8 week and 4 patients in the every 12 week maintenance treatment group. The most commonly reported infections were upper respiratory tract infection and pharyngitis, and the most commonly reported serious infection was abscess. Pneumonia was reported for 3 patients, (2 in the every 8 week and 1 in the every 12 week maintenance treatment groups). Herpes zoster was reported for 2 patients in the every 8 week maintenance treatment group. In Study Peds Crohn's, 18% of randomized patients experienced one or more infusion reactions, with no notable difference between treatment groups. Of the 112 patients in Study Peds Crohn's, there were no serious infusion reactions, and 2 patients had non-serious anaphylactoid reactions. Antibodies to REMICADE developed in 3% of pediatric patients in Study Peds Crohn's. Elevations of ALT up to 3 times the upper limit of normal (ULN) were seen in 18% of pediatric patients in CD clinical trials; 4% had ALT elevations  $\geq 3$  x ULN, and 1% had elevations  $\geq 5$  x ULN. (Median follow-up was 53 weeks.) The most common serious adverse events reported in the post-marketing experience in children were infections (some fatal) including opportunistic infections and tuberculosis, infusion reactions, and hypersensitivity reactions. Serious adverse events in the post-marketing experience with REMICADE in the pediatric population have also included malignancies, including hepatosplenic T-cell lymphomas (see **Boxed WARNINGS** and **WARNINGS**), transient hepatic enzyme abnormalities, lupus-like syndromes, and the development of autoantibodies. **Adverse Reactions in Psoriasis Studies** During the placebo-controlled portion across the three clinical trials up to Week 16, the proportion of patients who experienced at least 1 SAE (defined as resulting in death, life threatening, requires hospitalization, or persistent or significant disability/incapacity) was 1.7% in the 3 mg/kg REMICADE group, 3.2% in the placebo group, and 3.9% in the 5 mg/kg REMICADE group. Among patients in the 2 Phase 3 studies, 12.4% of patients receiving REMICADE 5 mg/kg every 8 weeks through one year of maintenance treatment experienced at least 1 SAE in Study I. In Study II, 4.1% and 4.7% of patients receiving REMICADE 3 mg/kg and 5 mg/kg every 8 weeks, respectively, through one year of maintenance treatment experienced at least 1 SAE. One death due to bacterial sepsis occurred 25 days after the second infusion of 5 mg/kg REMICADE. Serious infections included sepsis, and abscesses. In Study I, 2.7% of patients receiving REMICADE 5 mg/kg every 8 weeks through 1 year of maintenance treatment experienced at least 1 serious infection. In Study II, 1.0% and 1.3% of patients receiving REMICADE 3 mg/kg and 5 mg/kg, respectively, through 1 year of treatment experienced at least 1 serious infection. The most common serious infections (requiring hospitalization) were abscesses (skin, throat, and peri-rectal) reported by 5 (0.7%) patients in the 5 mg/kg REMICADE group. Two active cases of tuberculosis were reported: 6 weeks and 34 weeks after starting REMICADE. In placebo-controlled portion of the psoriasis studies, 7 of 1123 patients who received REMICADE at any dose were diagnosed with at least one NMSC compared to 0 of 334 patients who received placebo. In the psoriasis studies, 1% (15/1373) of patients experienced serum sickness or a combination of arthralgia and/or myalgia with fever, and/or rash, usually early in the treatment course. Of these patients, 6 required hospitalization due to fever, severe myalgia, arthralgia, swollen joints, and immobility. **Other Adverse Reactions** Safety data are available from 4779 REMICADE-treated adult patients, including 1304 with RA, 1106 with CD, 484 with UC, 202 with AS, 293 with PsA, 1373 with plaque PsO and 17 with other conditions. (For information on other adverse reactions in pediatric patients, see **ADVERSE REACTIONS, Adverse Reactions in Pediatric Crohn's Disease**.) Adverse events reported in  $\geq 5\%$  of all patients with RA receiving 4 or more infusions are listed below. The types and frequencies of adverse reactions observed were similar in REMICADE-treated RA, AS, PsA, plaque PsO and CD patients except for abdominal pain, which occurred in 26% of REMICADE-treated patients with CD. In the CD studies, there were insufficient numbers and duration of follow-up for patients who never received REMICADE to provide meaningful comparisons. The percentages of adverse events for placebo-treated patients (n=350; average weeks of follow-up 59) and REMICADE-treated patients (n=1129; average weeks of follow-up 66), respectively, are: **Gastrointestinal:** Nausea: 20, 21; Abdominal pain: 8, 12; Diarrhea: 12, 12; Dyspepsia: 7, 10; **Respiratory:** Upper respiratory tract infection: 25, 32; Sinusitis: 8, 14; Pharyngitis: 8, 12; Coughing: 8, 12; Bronchitis: 9, 10; Rhinitis: 5, 8; **Skin and appendages disorders:** Rash: 5, 10; Pruritus: 2, 7; **Body as a whole—general disorders:** Fatigue: 7, 9; Pain: 7, 8; **Resistance mechanism disorders:** Fever: 4, 7; Moniliasis: 3, 5; **Central and peripheral nervous system disorders:** Headache: 14, 18; **Musculoskeletal system disorders:** Back pain: 5, 8; Arthralgia: 7, 8; **Urinary system disorders:** Urinary tract infection: 6, 8; **Cardiovascular disorders, general:** Hypertension: 5, 7. Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in clinical trials of a drug cannot be directly compared to rates in clinical trials of another drug and may not predict the rates observed in broader patient populations in clinical practice. The most common serious adverse events observed in clinical trials were infections (see **ADVERSE REACTIONS, Infections**). Other serious, medically relevant adverse events  $\geq 0.2\%$  or clinically significant adverse events by body system were as follows: **Body as a whole:** allergic reaction, diaphragmatic hernia, edema, surgical/procedural sequela; **Blood:** pancytopenia; **Cardiovascular:** circulatory failure, hypotension, syncope; **Gastrointestinal:** constipation, gastrointestinal hemorrhage, ileus, intestinal obstruction, intestinal perforation, intestinal stenosis, pancreatitis, peritonitis, proctalgia; **Central & Peripheral Nervous:** meningitis, neuritis, peripheral neuropathy, dizziness; **Heart Rate and Rhythm:** arrhythmia, bradycardia, cardiac arrest, tachycardia; **Liver and Biliary:** biliary pain, cholecystitis, cholelithiasis, hepatitis; **Metabolic and Nutritional:** dehydration; **Musculoskeletal:** intervertebral disk herniation, tendon disorder; **Myo-, Endo-, Pericardial, and Coronary Valve:** myocardial infarction; **Platelet, Bleeding, and Clotting:** thrombocytopenia; **Neoplasms:** basal cell, breast, lymphoma; **Psychiatric:** confusion, suicide attempt; **Red Blood Cell:** anemia, hemolytic anemia; **Reproductive:** menstrual irregularity; **Resistance Mechanism:** cellulitis, sepsis, serum sickness; **Respiratory:** adult respiratory distress syndrome, lower respiratory tract infection (including pneumonia), pleural effusion, pleurisy, pulmonary edema, respiratory insufficiency; **Skin and Appendages:** increased sweating, ulceration; **Urinary:** renal calculus, renal failure; **Vascular (Extracardiac):** brain infarction, pulmonary embolism, thrombophlebitis; **White Cell and Reticuloendothelial:** leukopenia, lymphadenopathy. **Post-marketing Adverse Events** The following adverse events, some with fatal outcome, have been reported during post-approval use of REMICADE: neutropenia (see **WARNINGS, Hematologic Events**), interstitial lung disease (including pulmonary fibrosis/ interstitial pneumonitis and very rare rapidly progressive disease), idiopathic thrombocytopenic purpura, thrombotic thrombocytopenic purpura, pericardial effusion, systemic and cutaneous vasculitis, erythema multiforme, Stevens-Johnson Syndrome, toxic epidermal necrolysis, Guillain-Barré syndrome, psoriasis (including new onset and pustular, primarily palmar/plantar), transverse myelitis, and neuropathies (additional neurologic events have also been observed, see **WARNINGS, Neurologic Events**) and acute liver failure, jaundice, hepatitis, and cholestasis (see **WARNINGS, Hepatotoxicity**). Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to REMICADE exposure. The following serious adverse events have been reported in the post-marketing experience in children: infections (some fatal) including opportunistic infections and tuberculosis, infusion reactions, and hypersensitivity reactions. Serious adverse events in the post-marketing experience with REMICADE in the pediatric population have also included malignancies, including hepatosplenic T-cell lymphomas (see **Boxed WARNINGS** and **WARNINGS**), transient hepatic enzyme abnormalities, lupus-like syndromes, and the development of autoantibodies. **OVERDOSAGE:** Single doses up to 20 mg/kg have been administered without any direct toxic effect. In case of overdosage, it is recommended that the patient be monitored for any signs or symptoms of adverse reactions or effects and appropriate symptomatic treatment instituted immediately. **Administration Instructions Regarding Infusion Reactions** Adverse effects during administration of REMICADE have included flu-like symptoms, headache, dyspnea, hypotension, transient fever, chills, gastrointestinal symptoms, and skin rashes. Anaphylaxis might occur at any time during REMICADE infusion. Approximately 20% of REMICADE-treated patients in all clinical trials experienced an infusion reaction compared with 10% of placebo-treated patients (see **ADVERSE REACTIONS, Infusion-related Reactions**). Prior to infusion with REMICADE, premedication may be administered at the physician's discretion. Premedication could include antihistamines (anti-H1 +/- anti-H2), acetaminophen and/or corticosteroids. During infusion, mild to moderate infusion reactions may improve following slowing or suspension of the infusion, and upon resolution of the infusion, reinitiation at a lower infusion rate and/or therapeutic administration of antihistamines, acetaminophen, and/or corticosteroids. For patients that do not tolerate the infusion following these interventions, REMICADE should be discontinued. During or following infusion, patients that have severe infusion-related hypersensitivity reactions should be discontinued from further REMICADE treatment. The management of severe infusion reactions should be dictated by the signs and symptoms of the reaction. Appropriate personnel and medication should be available to treat anaphylaxis if it occurs.

**REFERENCES:** 1. *Am J Respir Crit Care Med.* 2000;161:S221-S247. 2. See latest Centers for Disease Control guidelines and recommendations for tuberculosis testing in immunocompromised patients. 3. Gardam MA, Keystone EC, Menzies R, et al. Anti-tumor necrosis factor agents and tuberculosis risk: mechanisms of action and clinical management. *Lancet Infect Dis.* 2003;3:148-155. 4. Belhadi K, Reyes F, Farost JP, et al. Hepatosplenic  $\gamma$ -T-cell lymphoma is a rare clinicopathologic entity with poor outcome: report on a series of 21 patients. *Blood.* 2003;102(13):4261-4269.

© 2007 Centocor, Inc.  
Malvern, PA 19355, USA  
1-800-457-6399

License #1242  
Revised April 2007  
IN07515

# Drug Utilization Boosts Nation's Health Tab

BY ALICIA AULT

Associate Editor, Practice Trends

WASHINGTON — The nation spent \$2 trillion, or \$7,000 per person, on health care in 2006. While that was only a small increase from the previous year, America's prescription drug tab increased by 8.5%, fueled largely by the new Medicare Part D drug benefit.

Health spending as a share of the nation's gross domestic product continues to rise, hitting 16% in 2006.

Total spending on physician and clinical services grew 5.9% to \$448 billion, the slowest rate of growth since 1999. Physician pay crawled almost to a halt, largely because of the freeze in Medicare's reimbursement rates in 2006. Private insurers seemed to have followed suit, said Cathy Cowan, an economist at the Centers for Medicare and Medicaid Services. Cowan, a coauthor of an annual analysis of the nation's health spending, spoke at a briefing on the report, which was published in the January/February issue of Health Affairs.

Medicare had the fastest rate of growth since 1981, according to the report. Spending increased 19% in 2006 to \$401 billion, driven largely by the prescription drug benefit and the cost of administration for that benefit and for Medicare Advantage, a managed care program.

Medicaid spending dropped for the first time since the program began in 1965. The 0.9% decrease was largely due to Medicaid enrollees who were shifted into Medicare for their prescription drugs.

Overall drug spending grew 8.5% in 2006—a far cry from the double-digit increases seen in the late 1990s, but still an increase from the 5.8% rise in spending in 2005. Half of the 2006 increase was due to greater utilization, not surprising given that about 23 million Medicare beneficia-

ries took advantage of the new benefit. Prescription prices increased by only a little over 3%, according to an annual analysis by actuaries at the Centers for Medicare and Medicaid Services.

The change in the drug rebate picture also contributed to rising drug costs. Under Medicaid, states received an average 30% rebate from drugmakers. Medicare, however, got only about 5% from manufacturers for the millions of beneficiaries who shifted out of Medicaid.

Medicare spent \$41 billion on Part D in 2006, with \$35 billion for drug purchases and \$6 billion for administration and “net cost of insurance”—that is, the cost of subsidizing premiums for low-income beneficiaries and costs for transferring beneficiaries into private plans. Medicare paid for 18% of all retail drugs, compared with only 2% in 2005.

The largest increase in drug utilization came from beneficiaries using the Part D benefit. But there was also increased drug use due to new indications for existing drugs, growth in several therapeutic classes, and rising use of specialty drugs such as injectable biologics for rheumatoid arthritis and multiple sclerosis, and anemia drugs for oncology. Hypnotics saw the largest rise in use of any drug class.

The authors said the data they had at hand and their analysis did not allow them to determine whether the prescription drug benefit had increased or lowered overall health care spending. “Sooner or later, somebody's going to do a dynamite study and figure this out,” said Richard Foster, the chief actuary at CMS.

Mr. Foster told reporters that the study showed that the “overall cost of prescription drugs has changed very little as a result of Part D.”

A study by Consumers Union, however, seemed to refute that claim. (See box.) ■

## Drug Prices Up Too: Consumers Union

Government economists have concluded that the Medicare Part D prescription drug benefit did not affect the price of pharmaceuticals in 2006, the program's first full year, but Consumers Union has issued another in a series of studies charging that drug prices are indeed rising.

Each month since December 2005, the consumer advocacy group has tracked the prices of five drugs commonly used by Medicare beneficiaries in a single ZIP code in each of five states—California, New York, Illinois, Florida, and Texas. The data are taken directly from Medicare.gov. According to Consumers Union, the data show that the majority of private insurers have consistently raised prices, sometimes at 2-3 times the rate of inflation.

Medicare beneficiaries might be bearing the brunt of price increases, especially because they usually are liable for a percentage of the drug's

price as a copayment. “We're seeing a lot of inflation,” said Consumers Union Senior Policy Analyst Bill Vaughan in an interview.

The group also found that prices generally rise the most from December to January—after a beneficiary has locked into a plan for the upcoming year. The average increase for the five drugs as a package (Lipitor, Celebrex, Zolof, nifedipine ER, and Altace) was \$369 from December 2007 to January 2008, according to Consumers Union.

“Most of these Medicare drug plans are increasing costs [at] double or triple the rate of inflation, which really torpedoes the insurance industry's claim that they are getting the best deal for seniors,” said Mr. Vaughan. “These continual price hikes are Exhibit A for Congress to give renewed attention to negotiating drug prices on behalf of America's taxpayers and seniors.”