

Neonatal Outcomes Vary With Diabetes Type

BY PATRICE WENDLING
Chicago Bureau

DALLAS — Women with type 2 diabetes had a similar incidence of adverse obstetric outcomes as those with type 1 diabetes but fewer adverse neonatal outcomes in a retrospective cohort analysis of 384 pregnancies.

As expected, patients with both type 1 and type 2 diabetes had worse obstetric and neonatal outcomes, compared with

nondiabetic controls, Dr. Kristin M. Knight and colleagues at the University of Rochester (N.Y.) reported in a poster at the annual meeting of the Society for Maternal-Fetal Medicine.

Using a preexisting database of pregestational diabetes patients, the researchers analyzed maternal and fetal outcomes of singleton pregnancies between July 2000 and August 2006 in 64 women with type 1 diabetes, 64 women with type 2 diabetes, and 256 matched controls with normal

glucose screening during pregnancy.

Patients with type 2 diabetes were significantly older (mean 30.1 years) than patients with type 1 diabetes (26.8 years) or controls (27.4 years), and had a significantly higher prepregnancy body mass index (37 vs. 27 vs. 24 kg m²). Mean hemoglobin A_{1c} values did not differ significantly between women with type 1 and type 2 diabetes (7.1% vs. 6.9%). The majority of women with type 2 diabetes (91%) were on insulin during pregnancy.

Both groups with diabetes had higher incidences of cesarean delivery, preeclampsia, preterm delivery, polyhydramnios, large-for-gestational-age infants, and neonatal ICU admission than did the 256 nondiabetic controls. However, the incidences were not different for women with type 2 vs. type 1 diabetes, the researchers noted.

Women with type 1 diabetes had a higher incidence of composite poor neonatal outcome (perinatal death, respiratory distress syndrome, sepsis, meconium aspiration, hypoglycemia, seizures, necrotizing enterocolitis, or intubation) than did women with type 2 diabetes and controls.

Those with type 1 diabetes had significantly more fetal congenital anomalies than did controls (6.3% vs. 1.2%). While such anomalies were elevated in women with type 2 diabetes, they were not significantly different (3.2%).

This finding differs from some of the other literature that's available, Dr. Knight said in an interview. This might be because the study population included only women seeking prenatal care, which is not the norm in clinical practice, and thus might have underrepresented poorer outcomes in patients with type 2 diabetes.

Overall, few data are available regarding pregnancy outcomes in type 2 diabetes, even though this type of diabetes is becoming increasingly common in reproductive-age women, she said. "Type 2 diabetic patients should receive the same degree of preconceptional counseling, diligent glucose control, and antenatal surveillance as type 1 diabetic patients, in order to minimize the occurrence of poor perinatal outcome," the authors wrote.

In a second poster at the meeting, diabetic macrovascular and microvascular disease during pregnancy was associated with reduced intrauterine fetal growth among 358 women with type 1 diabetes enrolled in a "Diabetes in Pregnancy" program at the University of Cincinnati.

The women were enrolled before 14 weeks' gestation, prospectively followed through the postpartum period, and treated with intensive insulin therapy.

They were classified at entry based on vascular status, with no vasculopathy present in 192, hypertension or background retinopathy in 79, proliferative retinopathy in only 18, nephropathy in only 42, and proliferative retinopathy and nephropathy in 26. Their mean ages were 24, 27, 27, 26, and 29 years, respectively; and they had been diagnosed with diabetes mellitus for 10, 15, 18, 14, and 18 years, respectively.

After controlling for gestation at delivery and maternal age and race, the odds ratio for delivery of a low-birth-weight infant (less than 2,500 g), compared with women without vasculopathy, was highest in women with proliferative retinopathy and nephropathy, Dr. Sina Haeri of the department of obstetrics and gynecology, Washington Hospital Center, and associates reported. Likewise, after controlling for maternal age and race, the odds ratio for delivery of a small-for-gestational-age infant, compared with women with no vasculopathy, was highest in those with proliferative retinopathy and nephropathy. ■

References: 1. Brange J, Volund A. Insulin analogs with improved pharmacokinetic profiles. *Adv Drug Deliv Rev.* 1999;35(2-3):307-335. 2. Raskin P, Guthrie RA, Leiter L, Riis A, Jovanovic L. Use of insulin aspart, a fast-acting insulin analog, as the mealtime insulin in the management of patients with type 1 diabetes. *Diabetes Care.* 2000;23(5):583-588. 3. Korytkowski M, Bell D, Jacobsen C, Suwannasari R, for the FlexPen® Study Team. A multicenter, randomized, open-label, comparative, two-period crossover trial of preference, efficacy, and safety profiles of a prefilled, disposable pen and conventional vial/syringe for insulin injection in patients with type 1 or 2 diabetes mellitus. *Clin Ther.* 2003;25(11):2836-2848. 4. Niskanen L, Jensen LE, Råstam J, Nygaard-Pedersen L, Erichsen K, Vora JP. Randomized, multinational, open-label, 2-period, crossover comparison of biphasic insulin aspart 30 and biphasic insulin lispro 25 and pen devices in adult patients with type 2 diabetes mellitus. *Clin Ther.* 2004;26(4):531-540.

NovoLog® Insulin aspart (rDNA origin) injection

BRIEF SUMMARY. PLEASE CONSULT PACKAGE INSERT FOR FULL PRESCRIBING INFORMATION.

INDICATIONS AND USAGE

NovoLog is indicated for the treatment of patients with diabetes mellitus, for the control of hyperglycemia. Because NovoLog has a more rapid onset and shorter duration of activity than human regular insulin, NovoLog given by injection should normally be used in regimens with an intermediate or long-acting insulin. NovoLog may also be infused subcutaneously by external insulin pumps. NovoLog may be administered intravenously under proper medical supervision in a clinical setting for glycemic control. (See WARNINGS; PRECAUTIONS [especially Usage in Pumps], Mixing of Insulins.)

CONTRAINDICATIONS

NovoLog is contraindicated during episodes of hypoglycemia and in patients hypersensitive to NovoLog or one of its excipients.

WARNINGS

NovoLog differs from regular human insulin by a more rapid onset and a shorter duration of activity. Because of the fast onset of action, the injection of NovoLog should immediately be followed by a meal. Because of the short duration of action of NovoLog, patients with diabetes also require a longer-acting insulin to maintain adequate glucose control. Glucose monitoring is recommended for all patients with diabetes and is particularly important for patients using external pump infusion therapy.

Hypoglycemia is the most common adverse effect of insulin therapy, including NovoLog. As with all insulins, the timing of hypoglycemia may differ among various insulin formulations. Any change of insulin dose should be made cautiously and only under medical supervision. Changes in insulin strength, manufacturer, type (e.g., regular, NPH, analog), species (animal, human), or method of manufacture (rDNA versus animal-source insulin) may result in the need for a change in dosage.

Insulin Pumps: When used in an external insulin pump for subcutaneous infusion, NovoLog should not be diluted or mixed with any other insulin. Physicians and patients should carefully evaluate information on pump use in the NovoLog physician and patient package inserts and in the pump manufacturer's manual (e.g., NovoLog-specific information should be followed for in-use time, frequency of changing infusion sets, or other details specific to NovoLog usage, because NovoLog-specific information may differ from general pump manual instructions).

Pump or infusion set malfunctions or insulin degradation can lead to hyperglycemia and ketosis in a short time because of the small subcutaneous depot of insulin. This is especially pertinent for rapid-acting insulin analogs that are more rapidly absorbed through skin and have shorter duration of action. These differences may be particularly relevant when patients are switched from multiple injection therapy or infusion with buffered regular insulin. Prompt identification and correction of the cause of hyperglycemia or ketosis is necessary. Interim therapy with subcutaneous injection may be required. (See PRECAUTIONS, Mixing of Insulins.)

PRECAUTIONS

General

Hypoglycemia and hypokalemia are among the potential clinical adverse effects associated with the use of all insulins. Because of differences in the action of NovoLog and other insulins, care should be taken in patients in whom such potential side effects might be clinically relevant (e.g., patients who are fasting, have autonomic neuropathy, or are using potassium-lowering drugs or patients taking drugs sensitive to serum potassium level). Insulin stimulates potassium movement into the cells, possibly leading to hypokalemia that left untreated may cause respiratory paralysis, ventricular arrhythmia, and death. Since intravenously administered insulin has a rapid onset of action, increased attention to hypoglycemia and hypokalemia is necessary. Therefore, glucose and potassium levels must be monitored closely when NovoLog or any other insulin is administered intravenously. Lipodystrophy and hypersensitivity are among other potential clinical adverse effects associated with the use of all insulins. As with all insulin preparations, the time course of NovoLog action may vary in different individuals or at different times in the same individual and is dependent on site of injection, blood supply, temperature, and physical activity. Adjustment of dosage of any insulin may be necessary if patients change their physical activity or their usual meal plan. Insulin requirements may be altered during illness, emotional disturbances, or other stresses.

Hypoglycemia - As with all insulin preparations, hypoglycemic reactions may be associated with the administration of NovoLog. Rapid changes in serum glucose levels may induce symptoms of hypoglycemia in persons with diabetes, regardless of the glucose value. Early warning symptoms of hypoglycemia may be different or less pronounced under certain conditions, such as long duration of diabetes, diabetic nerve disease, use of medications such as beta-blockers, or intensified diabetes control (see PRECAUTIONS, Drug Interactions). Such situations may result in severe hypoglycemia (and, possibly, loss of consciousness) prior to patients' awareness of hypoglycemia.

Renal Impairment - As with other insulins, the dose requirements for NovoLog may be reduced in patients with renal impairment.

Hepatic Impairment - As with other insulins, the dose requirements for NovoLog may be reduced in patients with hepatic impairment.

Allergy - Local Allergy - As with other insulin therapy, patients may experience redness, swelling, or itching at the site of injection. These minor reactions usually resolve in a few days to a few weeks, but in some occasions, may require discontinuation of NovoLog. In some instances, these reactions may be related to factors other than insulin, such as irritants in a skin cleansing agent or poor injection technique.

Systemic Allergy - Less common, but potentially more serious, is generalized allergy to insulin, which may cause rash (including pruritus) over the whole body, shortness of breath, wheezing, reduction in blood pressure, rapid pulse, or sweating. Severe cases of generalized allergy, including anaphylactic reaction, may be life threatening.

Localized reactions and generalized myalgias have been reported with the use of cresol as an injectable excipient.

In controlled clinical trials using injection therapy, allergic reactions were reported in 3 of 735 patients (0.4%) who received regular human insulin and 10 of 1394 patients (0.7%) who received NovoLog. During these and other trials, 3 of 2341 patients treated with NovoLog were discontinued due to allergic reactions.

Antibody Production - Increases in levels of anti-insulin antibodies that react with both human insulin and insulin aspart have been observed in patients treated with NovoLog. The number of patients treated with insulin aspart experiencing these increases is greater than the number among those treated with human regular insulin. Data from a 12-month controlled trial in patients with Type 1 diabetes suggest that the increase in these antibodies is transient. The differences in antibody levels between the human regular insulin and insulin aspart treatment groups observed at 3 and 6 months were no longer evident at 12 months. The clinical significance of these antibodies is not known. They do not appear to cause deterioration in HbA_{1c} or to necessitate increases in insulin dose.

Pregnancy and Lactation - Female patients should be advised to tell their physician if they intend to become, or if they become pregnant. Information is not available on the use of NovoLog during lactation (see PREGNANCY-TERATOGENIC EFFECTS-PREGNANCY CATEGORY).

Usage in Pumps - NovoLog is recommended for use in pump systems suitable for insulin infusion as listed below.

Pumps: Disetronic H-TRON® series, MiniMed 500 series and other equivalent pumps.

Reservoirs and Infusion Sets: NovoLog is recommended for use in any reservoir and infusion sets that are compatible with insulin and the specific pump. In-vitro studies have shown that pump malfunction, loss of cresol, and insulin degradation may occur when NovoLog is maintained in a pump system for more than 48 hours. Reservoirs and infusion sets should be changed at least every 48 hours.

NovoLog in clinical use should not be exposed to temperatures greater than 37°C (98.6°F). **NovoLog should not be mixed with other insulins or with a diluent when it is used in the pump.** (See WARNINGS; PRECAUTIONS, Mixing of Insulins.)

Laboratory Tests

As with all insulin therapy, the therapeutic response to NovoLog should be monitored by periodic blood glucose tests. Periodic measurement of glycosylated hemoglobin is recommended for the monitoring of long-term glycemic control. When NovoLog is administered intravenously, glucose and potassium levels must be closely monitored to avoid potentially fatal hypoglycemia and hypokalemia.

Drug Interactions

A number of substances affect glucose metabolism and may require insulin dose adjustment and particularly close monitoring.

- The following are examples of substances that may increase the blood-glucose-lowering effect and susceptibility to hypoglycemia: oral antidiabetic products, ACE inhibitors, disopyramide, fibrates, fluoxetine, monoamine oxidase (MAO) inhibitors, propoxyphene, salicylates, somatostatin analog (e.g., octreotide), sulfonamide antibiotics.

- The following are examples of substances that may reduce the blood-glucose-lowering effect: corticosteroids, niacin, danazol, diuretics, sympathomimetic agents (e.g., epinephrine, salbutamol, terbutaline), isoniazid, phenothiazine derivatives, somatropin, thyroid hormones, estrogens, progestogens (e.g., in oral contraceptives).

- Beta-blockers, clonidine, lithium salts, and alcohol may either potentiate or weaken the blood-glucose-lowering effect of insulin. Pentamidine may cause hypoglycemia, which may sometimes be followed by hyperglycemia.
- In addition, under the influence of sympatholytic medicinal products such as beta-blockers, clonidine, guanethidine, and reserpine, the signs of hypoglycemia may be reduced or absent.

Mixing of Insulins

- A clinical study in healthy male volunteers (n=24) demonstrated that mixing NovoLog with NPH human insulin immediately before injection produced some attenuation in the peak concentration of NovoLog, but that the time to peak and the total bioavailability of NovoLog were not significantly affected. If NovoLog is mixed with NPH human insulin, NovoLog should be drawn into the syringe first. The injection should be made immediately after mixing. Because there are no data on the compatibility of NovoLog and crystalline zinc insulin preparations, NovoLog should not be mixed with these preparations.

- The effects of mixing NovoLog with insulins of animal source or insulin preparations produced by other manufacturers have not been studied (see WARNINGS).

- Mixtures should not be administered intravenously.

- When used in external subcutaneous infusion pumps for insulin, NovoLog should not be mixed with any other insulins or diluent.**

Carcinogenicity, Mutagenicity, Impairment of Fertility

Standard 2-year carcinogenicity studies in animals have not been performed to evaluate the carcinogenic potential of NovoLog. In 52-week studies, Sprague-Dawley rats were dosed subcutaneously with NovoLog at 10, 50, and 200 U/kg/day (approximately 2, 8, and 32 times the human subcutaneous dose of 1.0 U/kg/day, based on U/body surface area, respectively). At a dose of 200 U/kg/day, NovoLog increased the incidence of mammary gland tumors in females when compared to untreated controls. The incidence of mammary tumors for NovoLog was not significantly different than for regular human insulin. The relevance of these findings to humans is not known. NovoLog was not genotoxic in the following tests: Ames test, mouse lymphoma cell forward gene mutation test, human peripheral blood lymphocyte chromosome aberration test, in vivo micronucleus test in mice, and in ex vivo UDS test in rat liver hepatocytes. In fertility studies in male and female rats, at subcutaneous doses up to 200 U/kg/day (approximately 32 times the human subcutaneous dose, based on U/body surface area), no direct adverse effects on male and female fertility, or general reproductive performance of animals was observed.

Pregnancy - Teratogenic Effects - Pregnancy Category B

All pregnancies have a background risk of birth defects, loss, or other adverse outcome regardless of drug exposure. This background risk is increased in pregnancies complicated by hyperglycemia and may be decreased with good metabolic control. It is essential for patients with diabetes or history of

gestational diabetes to maintain good metabolic control before conception and throughout pregnancy. Insulin requirements may decrease during the first trimester, generally increase during the second and third trimesters, and rapidly decline after delivery. Careful monitoring of glucose control is essential in such patients.

An open-label, randomized study compared the safety and efficacy of NovoLog versus human insulin in the treatment of pregnant women with Type 1 diabetes (322 exposed pregnancies (NovoLog: 157, human insulin: 165)). Two-thirds of the enrolled patients were already pregnant when they entered the study. Since only one-third of the patients enrolled before conception, the study was not large enough to evaluate the risk of congenital malformations. Mean HbA_{1c} of ~6% was observed in both groups during pregnancy, and there was no significant difference in the incidence of maternal hypoglycemia.

Subcutaneous reproduction and teratology studies have been performed with NovoLog and regular human insulin in rats and rabbits. In these studies, NovoLog was given to female rats before mating, during mating, and throughout pregnancy, and to rabbits during organogenesis. The effects of NovoLog did not differ from those observed with subcutaneous regular human insulin. NovoLog, like human insulin, caused pre- and post-implantation losses and visceral/skeletal abnormalities in rats at a dose of 200 U/kg/day (approximately 32 times the human subcutaneous dose of 1.0 U/kg/day, based on U/body surface area) and in rabbits at a dose of 10 U/kg/day (approximately three times the human subcutaneous dose of 1.0 U/kg/day, based on U/body surface area). The effects are probably secondary to maternal hypoglycemia at high doses. No significant effects were observed in rats at a dose of 50 U/kg/day and rabbits at a dose of 3 U/kg/day. These doses are approximately 8 times the human subcutaneous dose of 1.0 U/kg/day for rats and equal to the human subcutaneous dose of 1.0 U/kg/day for rabbits, based on U/body surface area.

Nursing Mothers

It is unknown whether insulin aspart is excreted in human milk. Many drugs, including human insulin, are excreted in human milk. For this reason, caution should be exercised when NovoLog is administered to a nursing mother.

Pediatric Use

A 24-week, parallel-group study of children and adolescents with Type 1 diabetes (n=283) age 6 to 18 years compared the following treatment regimens: NovoLog (n=187) or Novolin R (n=96). NPH insulin was administered as the basal insulin. NovoLog achieved glycemic control comparable to Novolin R, as measured by change in HbA_{1c}. The incidence of hypoglycemia was similar for both treatment groups. NovoLog and regular human insulin have also been compared in children with Type 1 diabetes (n=26) age 2 to 6 years. As measured by end-of-treatment HbA_{1c} and fructosamine, glycemic control with NovoLog was comparable to that obtained with regular human insulin. As observed in the 6 to 18 year old pediatric population, the rates of hypoglycemia were similar in both treatment groups.

Geriatric Use

Of the total number of patients (n=1,375) treated with NovoLog in 3 human insulin-controlled clinical studies, 2.6% (n=36) were 65 years of age or over. Half of these patients had Type 1 diabetes (18/1285) and half had Type 2 (18/90) diabetes. The HbA_{1c} response to NovoLog, as compared to human insulin, did not differ by age, particularly in patients with Type 2 diabetes. Additional studies in larger populations of patients 65 years of age or over are needed to permit conclusions regarding the safety of NovoLog in elderly compared to younger patients. Pharmacokinetic/pharmacodynamic studies to assess the effect of age on the onset of NovoLog action have not been performed.

ADVERSE REACTIONS

Clinical trials comparing NovoLog with regular human insulin did not demonstrate a difference in frequency of adverse events between the two treatments.

Adverse events commonly associated with human insulin therapy include the following:

Body as Whole - Allergic reactions (see PRECAUTIONS, Allergy).

Skin and Appendages - Injection site reaction, lipodystrophy, pruritus, rash (see PRECAUTIONS, Allergy, Usage in Pumps).

Other - Hypoglycemia, hyperglycemia and ketosis (see WARNINGS and PRECAUTIONS). In controlled clinical trials, small, but persistent elevations in alkaline phosphatase result were observed in some patients treated with NovoLog. The clinical significance of this finding is unknown.

OVERDOSAGE

Excess insulin may cause hypoglycemia and hypokalemia, particularly during IV administration. Hypoglycemia may occur as a result of an excess of insulin relative to food intake, energy expenditure, or both. Mild episodes of hypoglycemia usually can be treated with oral glucose. Adjustments in drug dosage, meal portions, or exercise may be needed. More severe episodes with coma, seizure, or neurologic impairment may be treated with intramuscular/subcutaneous glucagon or concentrated intravenous glucose. Sustained carbohydrate intake and observation may be necessary because hypoglycemia may recur after apparent clinical recovery. Hypokalemia must be corrected appropriately.

More detailed information is available on request.

Rx only

Date of Issue: January 26, 2007

Manufactured by Novo Nordisk A/S, DK-2880 Bagsvaerd, Denmark
Manufactured for Novo Nordisk Inc., Princeton, New Jersey 08540
www.novonordisk-us.com

NovoLog®, NovoPen® 3, NovoIn®, FlexPen®, and NovoFine® are registered trademarks of Novo Nordisk A/S.

H-TRON® is a registered trademark of Disetronic Medical Systems, Inc.

NovoLog® is covered by US Patent Nos 5,618,913; 5,866,538; and other patents pending.

© 2007 Novo Nordisk Inc. 130142R 5/07

