

Warfarin Less Effective in Real Life Than in Trials

BY MITCHEL L. ZOLER
Philadelphia Bureau

NEW ORLEANS — Real world experience with warfarin suggests that it is not as good at preventing strokes in patients with atrial fibrillation as clinical trial results have suggested, especially among African Americans.

A review of more than 23,000 Medicare patients with atrial fibrillation showed that overall, warfarin prophylaxis cut the

stroke rate by 34%, compared with a 65% cut in strokes that's been consistently seen in clinical trials, Brian F. Gage, M.D., reported at the 30th International Stroke Conference.

This lower efficacy in a real-world setting is "disappointing," said Dr. Gage, an internist at Washington University in St. Louis.

But warfarin performed even worse in the African Americans in the study. In this group, warfarin use was associated with a

trend toward more strokes, although this increase was not statistically significant, compared with African Americans not on warfarin.

In this analysis, the 95% confidence interval showed that, at best, warfarin prophylaxis in African Americans produced an 18% reduction in strokes, compared with untreated patients. This benefit is close to the 22% stroke reduction from aspirin prophylaxis in all patients, a suggestion that aspirin prophylaxis may be just as

good as or better than warfarin prophylaxis in African Americans. This hypothesis should be tested by analyzing results already collected in prior randomized, controlled studies, Dr. Gage told this newspaper.

The study used a national sample of 23,657 Medicare patients with atrial fibrillation who were treated during April 1998 through March 1999. Warfarin prophylaxis was used by 43% of African Americans in the study and by 50% of white patients.

Information culled from the medical records of the patients on warfarin therapy showed that this prophylaxis was often used in a less-than-ideal manner. Patients who regularly receive warfarin should have their dosage adjusted based on their international normalized ratio (INR), a

BRIEF SUMMARY: For full Prescribing Information, see package insert.

INDICATIONS AND USAGE CRESTOR is indicated: 1. as an adjunct to diet to reduce elevated total-C, LDL-C, ApoB, nonHDL-C, and TG levels and to increase HDL-C in patients with primary hypercholesterolemia (heterozygous familial and nonfamilial) and mixed dyslipidemia (Fredrickson Type IIa and IIb); 2. as an adjunct to diet for the treatment of patients with elevated serum TG levels (Fredrickson Type IV); 3. to reduce LDL-C, total-C, and ApoB in patients with homozygous familial hypercholesterolemia as an adjunct to other lipid-lowering treatments (e.g., LDL apheresis) or if such treatments are unavailable.

CONTRAINDICATIONS CRESTOR is contraindicated in patients with a known hypersensitivity to any component of this product. Rosuvastatin is contraindicated in patients with active liver disease or with unexplained persistent elevations of serum transaminases (see WARNINGS, Liver Enzymes). **Pregnancy and Lactation** Atherosclerosis is a chronic process and discontinuation of lipid-lowering drugs during pregnancy should have little impact on the outcome of long-term therapy of primary hypercholesterolemia. Cholesterol and other products of cholesterol biosynthesis are essential components for fetal development (including synthesis of steroids and cell membranes). Since HMG-CoA reductase inhibitors decrease cholesterol synthesis and possibly the synthesis of other biologically active substances derived from cholesterol, they may cause fetal harm when administered to pregnant women. Therefore, HMG-CoA reductase inhibitors are contraindicated during pregnancy and in nursing mothers. ROSUVASTATIN SHOULD BE ADMINISTERED TO WOMEN OF CHILDBEARING AGE ONLY WHEN SUCH PATIENTS ARE HIGHLY UNLIKELY TO CONCEIVE AND HAVE BEEN INFORMED OF THE POTENTIAL HAZARDS. If the patient becomes pregnant while taking this drug, therapy should be discontinued immediately and the patient apprised of the potential hazard to the fetus.

WARNINGS **Liver Enzymes** HMG-CoA reductase inhibitors, like some other lipid-lowering therapies, have been associated with biochemical abnormalities of liver function. The incidence of persistent elevations (>3 times the upper limit of normal [ULN]) occurring on 2 or more consecutive occasions in serum transaminases in fixed dose studies was 0.4, 0, 0, and 0.1% in patients who received rosuvastatin 5, 10, 20, and 40 mg, respectively. In most cases, the elevations were transient and resolved or improved on continued therapy or after a brief interruption in therapy. There were two cases of jaundice, for which a relationship to rosuvastatin therapy could not be determined, which resolved after discontinuation of therapy. There were no cases of liver failure or irreversible liver disease in these trials. It is recommended that liver function tests be performed before and at 12 weeks following both the initiation of therapy and any elevation of dose, and periodically (e.g., semiannually) thereafter. Liver enzyme changes generally occur in the first 3 months of treatment with rosuvastatin. Patients who develop increased transaminase levels should be monitored until the abnormalities have resolved. Should an increase in ALT or AST of >3 times ULN persist, reduction of dose or withdrawal of rosuvastatin is recommended. Rosuvastatin should be used with caution in patients who consume substantial quantities of alcohol and/or have a history of liver disease (see CLINICAL PHARMACOLOGY, Special Populations, Hepatic Insufficiency). Active liver disease or unexplained persistent transaminase elevations are contraindications to the use of rosuvastatin (see CONTRAINDICATIONS).

Myopathy/Rhabdomyolysis Rare cases of rhabdomyolysis with acute renal failure secondary to myoglobinuria have been reported with rosuvastatin and with other drugs in this class. Uncomplicated myalgia has been reported in rosuvastatin-treated patients (see ADVERSE REACTIONS). Creatine kinase (CK) elevations (>10 times upper limit of normal) occurred in 0.2% to 0.4% of patients taking rosuvastatin at doses of up to 40 mg in clinical studies. Treatment-related myopathy, defined as muscle aches or muscle weakness in conjunction with increases in CK values >10 times upper limit of normal, was reported in up to 0.1% of patients taking rosuvastatin doses of up to 40 mg in clinical studies. Rare cases of rhabdomyolysis were seen with higher than recommended doses (80 mg) of rosuvastatin in clinical trials. Factors that may predispose patients to myopathy with HMG-CoA reductase inhibitors include advanced age (>65 years), hypothyroidism, and renal insufficiency. The incidence of myopathy increased at doses of rosuvastatin above the recommended dosage range. Consequently: 1. Rosuvastatin should be prescribed with caution in patients with predisposing factors for myopathy, such as, renal impairment (see DOSAGE AND ADMINISTRATION), advanced age, and hypothyroidism. 2. Patients should be advised to promptly report unexplained muscle pain, tenderness, or weakness, particularly if accompanied by malaise or fever. Rosuvastatin therapy should be discontinued if markedly elevated CK levels occur or myopathy is diagnosed or suspected. 3. The risk of myopathy during treatment with rosuvastatin may be increased with concurrent administration of other lipid-lowering therapies or cyclosporine, (see CLINICAL PHARMACOLOGY, Drug Interactions, PRECAUTIONS, Drug Interactions, and DOSAGE AND ADMINISTRATION). The benefit of further alterations in lipid levels by the combined use of rosuvastatin with fibrates or niacin should be carefully weighed against the potential risks of this combination. Combination therapy with rosuvastatin and gemfibrozil should generally be avoided. (See DOSAGE AND ADMINISTRATION and PRECAUTIONS, Drug Interactions). 4. The risk of myopathy during treatment with rosuvastatin may be increased in circumstances which increase rosuvastatin drug levels (see CLINICAL PHARMACOLOGY, Special Populations, Race and Renal Insufficiency, and PRECAUTIONS, General). 5. Rosuvastatin therapy should also be temporarily withheld in any patient with an acute, serious condition suggestive of myopathy or predisposing to the development of renal failure secondary to rhabdomyolysis (e.g., sepsis, hypotension, major surgery, trauma, severe metabolic, endocrine, and electrolyte disorders, or uncontrolled seizures).

PRECAUTIONS **General** Before instituting therapy with rosuvastatin, an attempt should be made to control hypercholesterolemia with appropriate diet and exercise, weight reduction in obese patients, and treatment of underlying medical problems (see INDICATIONS AND USAGE). Administration of rosuvastatin 20 mg to patients with severe renal impairment ($Cl_{CR} < 30$ mL/min/1.73 m²) resulted in a 3-fold increase in plasma concentrations of rosuvastatin compared with healthy volunteers (see WARNINGS, Myopathy/Rhabdomyolysis and DOSAGE AND ADMINISTRATION). Pharmacokinetic studies show an approximate 2-fold elevation in median exposure in Japanese subjects residing in Japan and in Chinese subjects residing in Singapore compared with Caucasians residing in North America and Europe. The contribution of environmental and genetic factors to the difference observed has not been determined. However, these increases should be considered when making rosuvastatin dosing decisions for patients of Japanese and Chinese ancestry. (See WARNINGS, Myopathy/Rhabdomyolysis; CLINICAL PHARMACOLOGY, Special Populations, Race.) **Information for Patients** Patients should be advised to report promptly unexplained muscle pain, tenderness, or weakness, particularly if accompanied by malaise or fever. When taking rosuvastatin with an aluminum and magnesium hydroxide combination antacid, the antacid should be taken at least 2 hours after rosuvastatin administration (see CLINICAL PHARMACOLOGY, Drug Interactions). **Laboratory Tests** In the rosuvastatin clinical trial program, dipstick-positive proteinuria and microscopic hematuria were observed among rosuvastatin-treated patients, predominantly in patients dosed above the recommended dose range (i.e., 80 mg). However, this finding was more frequent in patients taking rosuvastatin 40 mg, when compared to lower doses of rosuvastatin or comparator statins, though it was generally transient and was not associated with worsening renal function. Although the clinical significance of this finding is unknown, a dose reduction should be considered for patients on rosuvastatin 40 mg therapy with unexplained persistent proteinuria during routine urinalysis testing. **Drug Interactions** **Cyclosporine:** When rosuvastatin 10 mg was coadministered with cyclosporine in cardiac transplant patients, rosuvastatin mean C_{max} and mean AUC were increased 11-fold and 7-fold, respectively, compared with healthy volunteers. These

increases are considered to be clinically significant and require special consideration in the dosing of rosuvastatin to patients taking concomitant cyclosporine (see WARNINGS, Myopathy/Rhabdomyolysis, and DOSAGE AND ADMINISTRATION). **Warfarin:** Coadministration of rosuvastatin to patients on stable warfarin therapy resulted in clinically significant rises in INR (>4, baseline 2-3). In patients taking coumarin anticoagulants and rosuvastatin concomitantly, INR should be determined before starting rosuvastatin and frequently enough during early therapy to ensure that no significant alteration of INR occurs. Once a stable INR time has been documented, INR can be monitored at the intervals usually recommended for patients on coumarin anticoagulants. If the dose of rosuvastatin is changed, the same procedure should be repeated. Rosuvastatin therapy has not been associated with bleeding or with changes in INR in patients not taking anticoagulants. **Gemfibrozil:** Coadministration of a single rosuvastatin dose to healthy volunteers on gemfibrozil (600 mg twice daily) resulted in a 2.2- and 1.9-fold, respectively, increase in mean C_{max} and mean AUC of rosuvastatin (see DOSAGE AND ADMINISTRATION). **Endocrine Function** Although clinical studies have shown that rosuvastatin alone does not reduce basal plasma cortisol concentration or impair adrenal reserve, caution should be exercised if any HMG-CoA reductase inhibitor or other agent used to lower cholesterol levels is administered concomitantly with drugs that may decrease the levels or activity of endogenous steroid hormones such as ketoconazole, spironolactone, and cimetidine. **CNS Toxicity** CNS vascular lesions, characterized by perivascular hemorrhages, edema, and mononuclear cell infiltration of perivascular spaces, have been observed in dogs treated with several other members of this drug class. A clinically similar drug in this class produced dose-dependent optic nerve degeneration (Wallerian degeneration of retinogeniculate fibers) in dogs, at a dose that produced plasma drug levels about 30 times higher than the mean drug level in humans taking the highest recommended dose. Edema, hemorrhage, and partial necrosis in the interstitium of the choroid plexus was observed in a female dog sacrificed moribund at day 24 at 90 mg/kg/day by oral gavage (systemic exposures 100 times the human exposure at 40 mg/day based on AUC comparisons). Corneal opacity was seen in dogs treated for 52 weeks at 6 mg/kg/day by oral gavage (systemic exposures 20 times the

ADVERSE REACTIONS Rosuvastatin is generally well tolerated. Adverse reactions have usually been mild and transient. In clinical studies of 10,275 patients, 3.7% were discontinued due to adverse experiences attributable to rosuvastatin. The most frequent adverse events thought to be related to rosuvastatin were myalgia, constipation, asthenia, abdominal pain, and nausea. **Clinical Adverse Experiences** Adverse experiences, regardless of causality assessment, reported in ≥2% of patients in placebo-controlled clinical studies of rosuvastatin are shown in Table 1; discontinuations due to adverse events in these studies of up to 12 weeks duration occurred in 3% of patients on rosuvastatin and 5% on placebo.

Table 1. Adverse Events in Placebo-Controlled Studies

Adverse event	Rosuvastatin N=744	Placebo N=382
Pharyngitis	9.0	7.6
Headache	5.5	5.0
Diarrhea	3.4	2.9
Dyspepsia	3.4	3.1
Nausea	3.4	3.1
Myalgia	2.8	1.3
Asthenia	2.7	2.6
Back pain	2.6	2.4
Flu syndrome	2.3	1.8
Urinary tract infection	2.3	1.6
Rhinitis	2.2	2.1
Sinusitis	2.0	1.8

In addition, the following adverse events were reported, regardless of causality assessment, in ≥1% of 10,275 patients treated with rosuvastatin in clinical studies. The events in *italics* occurred in ≥2% of these patients. **Body as a Whole:** Abdominal pain, accidental injury, chest pain, infection, pain, pelvic pain, and neck pain. **Cardiovascular System:** Hypertension, angina pectoris, vasodilatation, and palpitation. **Digestive System:** Constipation, gastroenteritis, vomiting, flatulence, periodontal abscess, and gastritis. **Endocrine:** Diabetes mellitus. **Hemic and Lymphatic System:** Anemia and ecchymosis. **Metabolic and Nutritional Disorders:** Peripheral edema. **Musculoskeletal System:** Arthritis, arthralgia, and pathological fracture. **Nervous System:** Dizziness, insomnia, hypertonnia, paresthesia, depression, anxiety, vertigo, and neuralgia. **Respiratory System:** Bronchitis, cough increased, dyspnea, pneumonia, and asthma. **Skin and Appendages:** Rash and pruritus. **Laboratory Abnormalities:** In the rosuvastatin clinical trial program, dipstick-positive proteinuria and microscopic hematuria were observed among rosuvastatin-treated patients, predominantly in patients dosed above the recommended dose range (i.e., 80 mg). However, this finding was more frequent in patients taking rosuvastatin 40 mg, when compared to lower doses of rosuvastatin or comparator statins, though it was generally transient and was not associated with worsening renal function. (See PRECAUTIONS, Laboratory Tests.) Other abnormal laboratory values reported were elevated creatine phosphokinase, transaminases, hyperglycemia, glutamyl transpeptidase, alkaline phosphatase, bilirubin, and thyroid function abnormalities. Other adverse events reported less frequently than 1% in the rosuvastatin clinical study program, regardless of causality assessment, included arrhythmia, hepatitis, hypersensitivity reactions (i.e., face edema, thrombocytopenia, leukopenia, vesiculobullous rash, urticaria, and angioedema), kidney failure, syncope, myasthenia, myositis, pancreatitis, photosensitivity reaction, myopathy, and rhabdomyolysis.

OVERDOSAGE There is no specific treatment in the event of overdose. In the event of overdose, the patient should be treated symptomatically and supportive measures instituted as required. Hemodialysis does not significantly enhance clearance of rosuvastatin. **DOSAGE AND ADMINISTRATION** The patient should be placed on a standard cholesterol-lowering diet before receiving CRESTOR and should continue on this diet during treatment. CRESTOR can be administered as a single dose at any time of day, with or without food. **Hypercholesterolemia (Heterozygous Familial and Nonfamilial) and Mixed Dyslipidemia (Fredrickson Type IIa and IIb)** The dose range for CRESTOR is 5 to 40 mg once daily. Therapy with CRESTOR should be individualized according to goal of therapy and response. The usual recommended starting dose of CRESTOR is 10 mg once daily. Initiation of therapy with 5 mg once daily may be considered for patients requiring less aggressive LDL-C reductions or who have predisposing factors for myopathy (see WARNINGS, Myopathy/Rhabdomyolysis). For patients with marked hypercholesterolemia (LDL-C > 190 mg/dL) and aggressive lipid targets, a 20-mg starting dose may be considered. The 40-mg dose of CRESTOR should be reserved for those patients who have not achieved goal LDL-C at 20 mg (see WARNINGS, Myopathy/Rhabdomyolysis). After initiation and/or upon titration of CRESTOR, lipid levels should be analyzed within 2 to 4 weeks and dosage adjusted accordingly. **Homozygous Familial Hypercholesterolemia** The recommended starting dose of CRESTOR is 20 mg once daily in patients with homozygous FH. The maximum recommended daily dose is 40 mg. CRESTOR should be used in these patients as an adjunct to other lipid-lowering treatments (e.g., LDL apheresis) or if such treatments are unavailable. Response to therapy should be estimated from pre-apheresis LDL-C levels. **Dosage in Patients Taking Cyclosporine** In patients taking cyclosporine, therapy should be limited to CRESTOR 5 mg once daily (see WARNINGS, Myopathy/Rhabdomyolysis, and PRECAUTIONS, Drug Interactions). **Concomitant Lipid-Lowering Therapy** The effect of CRESTOR on LDL-C and total-C may be enhanced when used in combination with a bile acid binding resin. If CRESTOR is used in combination with gemfibrozil, the dose of CRESTOR should be limited to 10 mg once daily (see WARNINGS, Myopathy/Rhabdomyolysis, and PRECAUTIONS, Drug Interactions). **Dosage in Patients With Renal Insufficiency** No modification of dosage is necessary for patients with mild to moderate renal insufficiency. For patients with severe renal impairment ($Cl_{CR} < 30$ mL/min/1.73 m²) not on hemodialysis, dosing of CRESTOR should be started at 5 mg once daily and not to exceed 10 mg once daily (see PRECAUTIONS, General, and CLINICAL PHARMACOLOGY, Special Populations, Renal Insufficiency).

Rx only

References: 1. Data on file, DA-CRS-13. 2. Grundy SM, Cleeman JJ, Merz CN, et al. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. *Circulation*. 2004;110:227-239. 3. IMS National Prescription Audit: November 2003-October 2004. 4. Shepherd J, Hunnigake DB, Stein EA, et al. The safety of rosuvastatin. *Am J Cardiol*. 2004;94:882-888. 5. Prescribing Information for CRESTOR. AstraZeneca, Wilmington, DE. 6. Rosuvastatin Information Web site. Rosuvastatin Clinical Information-Postmarketing Experience, Safety Information. Available at: <http://www.rosuvastatininformation.com>. Accessed November 30, 2004. 7. Data on file, DA-CRS-01.

CRESTOR is a registered trademark of the AstraZeneca group of companies.

Please visit our Web site at www.crestor.com

© AstraZeneca 2004

Licensed from SHIONOGI & CO., LTD., Osaka, Japan

Manufactured for: AstraZeneca Pharmaceuticals LP

Wilmington, DE 19850

By: IPR Pharmaceuticals, Inc.

Carolina, PR 00984

PCC 630100

10/04 224758

AstraZeneca



Prophylaxis with aspirin may be just as good as or better than prophylaxis with warfarin in African Americans.

DR. GAGE

measure of clotting time. Ideally, INRs should be measured about every 28 days in patients who regularly take warfarin.

Among all white patients, the average time between INR measurements was 26 days, and among African Americans the average interval was 30 days. But 25% of the white patients on warfarin had an interval of 39 days or longer between INR measurements. Among African Americans on warfarin, 25% had an interval of 57 days or longer between measurements, Dr. Gage said at the conference, sponsored by the American Stroke Association. For these subgroups, the interval between INR measurements was "way too long," he said.

But substandard INR monitoring was not the only reason patients got less benefit from warfarin prophylaxis, compared with the benchmark of clinical trials. The rate of protection from stroke remained unexpectedly low among African Americans even in an analysis that controlled for the frequency of INR monitoring as well as clinical variables that predispose patients to strokes.

Other factors that were not controlled for in this analysis, and that may help explain warfarin's underachievement, include poor compliance with the warfarin regimen, inadequacies in the health care setting, and inadequate access to anticoagulant services, said Dr. Gage, who is also medical director of the Blood Thinner Clinic at Barnes-Jewish Hospital in St. Louis.

The clinical trials that assessed warfarin's efficacy for preventing strokes in patients with atrial fibrillation were highly selective; more than 90% of patients who were initially assessed for these trials were eventually excluded. The clinical trial results therefore came from patients that were mostly white, less than age 75 years, and followed very closely, and these results may not be generalizable to other health care settings, Dr. Gage said.