

Strategies to Limit Bleeding in PCI Underused

BY MARY ANN MOON

FROM JAMA

Strategies to limit periprocedural bleeding in percutaneous coronary interventions are effective, particularly in the patients at highest risk for bleeding, according to a study of a nationwide database.

Unfortunately, these strategies are underused, and the very patients most like-

ly to benefit are the least likely to receive them, reported Dr. Steven P. Marso of Saint Luke's Mid America Heart Institute, Kansas City, Mo., and his associates.

Targeting bleeding complications "holds great potential for improving the safety and cost-effectiveness of PCI," since more than 1 million of the procedures are performed in the United States every year and bleeding complications occur in 2%-6% of them. Major bleeding events raise

the risks for early and late mortality, MI, and stroke, and also "result in an average 4- to 6-day increase in length of stay and, on average, increase hospital costs by \$6,000-\$8,000," the investigators noted.

They studied bleeding complications using the National Cardiovascular Data Registry, a nationwide database of catheterization and PCI procedures at more than 1,100 medical centers. They assessed data on more than 1.5 million

patients who underwent PCI via the femoral artery approach during the period from Jan. 1, 2004, to Sept. 30, 2008.

To mitigate bleeding, vascular closure devices such as Angio-Seal (St. Jude Medical) and Perclose A-T (Abbott Vascular), the drug bivalirudin (Angiomax), or both were used in 24%, 23%, and 18% of patients, respectively. Manual compression, used in 35% of patients, served as the control strategy.

ZYVOX® linezolid injection, tablets and for oral suspension
Brief summary of prescribing information.

INDICATIONS AND USAGE ZYVOX formulations are indicated in the treatment of the following infections caused by susceptible strains of the designated microorganisms (see **PRECAUTIONS, Pediatric Use**). **Vancomycin-Resistant *Enterococcus faecium* infections**, including cases with concurrent bacteremia. **Nosocomial pneumonia** caused by *Staphylococcus aureus* (methicillin-susceptible and -resistant strains), or *Streptococcus pneumoniae* (including multidrug-resistant strains [MDRSP¹]). **Complicated skin and skin structure infections, including diabetic foot infections, without concomitant osteomyelitis**, caused by *Staphylococcus aureus* (methicillin-susceptible and -resistant strains), *Streptococcus pyogenes*, or *Streptococcus agalactiae*. ZYVOX has not been studied in the treatment of decubitus ulcers. **Uncomplicated skin and skin structure infections** caused by *Staphylococcus aureus* (methicillin-susceptible only) or *Streptococcus pyogenes*. **Community-acquired pneumonia** caused by *Streptococcus pneumoniae* (including multidrug-resistant strains [MDRSP¹]), including cases with concurrent bacteremia, or *Staphylococcus aureus* (methicillin-susceptible strains only). To reduce the development of drug-resistant bacteria and maintain the effectiveness of ZYVOX and other antibacterial drugs, ZYVOX should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

CONTRAINDICATIONS ZYVOX formulations are contraindicated for use in patients who have known hypersensitivity to linezolid or any of the other product components. ZYVOX should not be used in patients taking any medicinal product which inhibits monoamine oxidases A or B (e.g. phenelzine, isocarboxazid) or within 2 weeks of taking any such medicinal product. Unless patients are monitored for potential increases in blood pressure, ZYVOX should not be administered to patients with uncontrolled hypertension, pheochromocytoma, thyrotoxicosis and/or patients taking any of the following types of medications: directly and indirectly acting sympathomimetic agents (e.g. pseudoephedrine), vasopressor agents (e.g. epinephrine, norepinephrine), and dopaminergic agents (e.g. dopamine, dobutamine). Unless patients are carefully observed for signs and/or symptoms of serotonin syndrome, ZYVOX should not be administered to patients with carcinoid syndrome and/or patients taking any of the following medications: serotonin re-uptake inhibitors, tricyclic antidepressants, serotonin 5-HT₁ receptor agonists (triptans), meperidine, or buspirone. **WARNINGS** Myelosuppression (including anemia, leukopenia, pancytopenia, and thrombocytopenia) has been reported in patients receiving ZYVOX. In cases where the outcome is known, when ZYVOX was discontinued, the affected hematologic parameters have risen toward pretreatment levels. Complete blood counts should be monitored weekly in patients who receive ZYVOX, particularly in those who receive ZYVOX for longer than two weeks, those with pre-existing myelosuppression, those receiving concomitant drugs that produce bone marrow suppression, or those with a chronic infection who have received previous or concomitant antibiotic therapy. Discontinuation of therapy with ZYVOX should be considered in patients who develop or have worsening myelosuppression. In adult and juvenile dogs and rats, myelosuppression, reduced extramedullary hematopoiesis in spleen and liver, and lymphoid depletion of thymus, lymph nodes, and spleen were observed. **Mortality imbalance in an Investigational Study in Patients With Catheter-related Bloodstream Infections, Including Those With Catheter-site Infections.** ZYVOX is not approved and should not be used for the treatment of patients with catheter-related bloodstream infections or catheter-site infections. In an open-label investigational study in seriously ill patients with intravascular catheter-related infections, an imbalance in mortality was seen in patients treated with ZYVOX compared with vancomycin/dicloxacillin/oxacillin. While causality has not been established, mortality was higher in patients treated with ZYVOX who were infected with Gram-negative organisms alone, with both Gram-positive and Gram-negative organisms, or who had no infection when they entered the study. Patients with Gram-positive infections had no difference in mortality. ZYVOX has no clinical activity against Gram-negative pathogens and is not indicated for the treatment of Gram-negative infections. It is critical that specific Gram-negative therapy be initiated immediately if a concomitant Gram-negative pathogen is documented or suspected. *Clostridium difficile*-associated diarrhea (CDAD) has been reported with the use of nearly all antibacterial agents, including ZYVOX, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*. *C. difficile* produces toxins A and B, which contribute to the development of CDAD. Hypertoxin-producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur more than 2 months after the administration of antibacterial agents. If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

PRECAUTIONS General Lactic acidosis has been reported with the use of ZYVOX. In reported cases, patients experienced repeated episodes of nausea and vomiting. Patients who develop recurrent nausea or vomiting, unexplained acidosis, or a low bicarbonate level while receiving ZYVOX should receive immediate medical evaluation. Spontaneous reports of serotonin syndrome associated with the co-administration of ZYVOX and serotonergic agents, including antidepressants such as selective serotonin reuptake inhibitors (SSRIs), have been reported (see **PRECAUTIONS, Drug Interactions**). Where administration of ZYVOX and concomitant serotonergic agents is clinically appropriate, patients should be closely observed for signs and symptoms of serotonin syndrome such as cognitive dysfunction, hyperpyrexia, hyperreflexia, and incoordination. If signs or symptoms occur physicians should consider discontinuation of either one or both agents. If the concomitant serotonergic agent is withdrawn, discontinuation symptoms can be observed (see package insert of the specified agent(s) for a description of the associated discontinuation symptoms). Peripheral and optic neuropathy have been reported in patients treated with ZYVOX, primarily those patients treated for longer than the maximum recommended duration of 28 days. In cases of optic neuropathy that progressed to loss of vision, patients were treated for extended periods beyond the maximum recommended duration. Visual blurring has been reported in some patients treated with ZYVOX for less than 28 days. If patients experience symptoms of visual impairment, such as changes in visual acuity, changes in color vision, blurred vision, or visual field defect, prompt ophthalmic evaluation is recommended. **Visual function should be monitored in all patients taking ZYVOX for extended periods (≥3 months) and in all patients reporting new visual symptoms regardless of length of therapy with ZYVOX.** If peripheral or optic neuropathy occurs, the continued use of ZYVOX in these patients should be weighed against the potential risks. Convulsions have been reported in patients treated with ZYVOX. In some of these cases, a history of seizures or risk factors for seizures was reported. The use of antibiotics may promote the overgrowth of nonsusceptible organisms. Should superinfection occur during therapy, appropriate measures should be taken. ZYVOX has not been studied in patients with uncontrolled hypertension, pheochromocytoma, carcinoid syndrome, or untreated hyperthyroidism. The safety and efficacy of ZYVOX formulations given for longer than 28 days have not been evaluated in controlled clinical trials. Prescribing ZYVOX in the absence of a proven or

strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria. **Information for Patients** Patients should be advised that: ZYVOX may be taken with or without food. They should inform their physician if they have a history of hypertension. Large quantities of foods or beverages with high tyramine content should be avoided while taking ZYVOX. Quantities of tyramine consumed should be less than 100 mg per meal. Foods high in tyramine content include those that may have undergone protein changes by aging, fermentation, pickling, or smoking to improve flavor, such as aged cheeses (0 to 15 mg tyramine per ounce); fermented or air-dried meats (0.1 to 8 mg tyramine per ounce); sauerkraut (8 mg tyramine per 8 ounces); soy sauce (5 mg tyramine per 1 teaspoon); tap beers (4 mg tyramine per 12 ounces); red wines (0 to 6 mg tyramine per 8 ounces). The tyramine content of any protein-rich food may be increased if stored for long periods or improperly refrigerated. They should inform their physician if taking medications containing pseudoephedrine HCl or phenylpropanolamine HCl, such as cold remedies and decongestants. They should inform their physician if taking serotonin re-uptake inhibitors or other antidepressants.

Phenylketonurics: Each 5 mL of the 100 mg/5 mL ZYVOX for Oral Suspension contains 20 mg phenylalanine. The other ZYVOX formulations do not contain phenylalanine. Contact your physician or pharmacist. They should inform their physician if they experience changes in vision. They should inform their physician if they have a history of seizures. Diarrhea is a common problem caused by antibiotics, which usually ends when the antibiotic is discontinued. Sometimes after starting treatment with antibiotics, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibiotic. If this occurs, patients should contact their physician as soon as possible. Patients should be counseled that antibacterial drugs including ZYVOX should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When ZYVOX is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by ZYVOX or other antibacterial drugs in the future. **Drug Interactions**

Monoamine Oxidase Inhibition: Linezolid is a reversible, nonselective inhibitor of monoamine oxidase. Therefore, linezolid has the potential for interaction with adrenergic and serotonergic agents. **Adrenergic Agents:** Some individuals receiving ZYVOX may experience a reversible enhancement of the pressor response to indirect-acting sympathomimetic agents, vasopressor or dopaminergic agents. Commonly used drugs such as phenylpropanolamine and pseudoephedrine have been specifically studied. Initial doses of adrenergic agents, such as dopamine or epinephrine, should be reduced and titrated to achieve the desired response. **Serotonergic Agents:** Co-administration of linezolid and serotonergic agents was not associated with serotonin syndrome in Phase 1, 2 or 3 studies. Spontaneous reports of serotonin syndrome associated with co-administration of ZYVOX and serotonergic agents, including antidepressants such as selective serotonin reuptake inhibitors (SSRIs), have been reported. Patients who are treated with ZYVOX and concomitant serotonergic agents should be closely observed as described in the **PRECAUTIONS, General Section**. **Drug-Laboratory Test Interactions** There are no reported drug-laboratory test interactions. **Pregnancy Teratogenic Effects. Pregnancy Category C:** Linezolid was not teratogenic in mice, rats, or rabbits at exposure levels 6.5-fold (in mice), equivalent to (in rats), or 0.5-fold (in rabbits) the expected human exposure level, based on AUCs. However, embryo and fetal toxicities were seen (see **Non-teratogenic Effects**). There are no adequate and well-controlled studies in pregnant women. ZYVOX should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Non-teratogenic Effects In mice, embryo and fetal toxicities were seen only at doses that caused maternal toxicity (clinical signs and reduced body weight gain). A dose of 450 mg/kg/day (6.5-fold the estimated human exposure level based on AUCs) correlated with increased postimplantation embryo death, including total litter loss, decreased fetal body weights, and an increased incidence of costal cartilage fusion. In rats, mild fetal toxicity was observed at 15 and 50 mg/kg/day (exposure levels 0.22-fold to approximately equivalent to the estimated human exposure, respectively based on AUCs). The effects consisted of decreased fetal body weights and reduced ossification of sternbrae, a finding often seen in association with decreased fetal body weights. Slight maternal toxicity, in the form of reduced body weight gain, was seen at 50 mg/kg/day. In rabbits, reduced fetal body weight occurred only in the presence of maternal toxicity (clinical signs, reduced body weight gain and food consumption) when administered at a dose of 15 mg/kg/day (0.5-fold the estimated human exposure based on AUCs). When female rats were treated with 50 mg/kg/day (approximately equivalent to the estimated human exposure based on AUCs) of linezolid during pregnancy and lactation, survival of pups was decreased on postnatal days 1 to 4. Male and female pups permitted to mature to reproductive age, when mated, showed an increase in preimplantation loss. **Nursing Mothers** Linezolid and its metabolites are excreted in the milk of lactating rats. Concentrations in milk were similar to those in maternal plasma. It is not known whether linezolid is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when ZYVOX is administered to a nursing woman. **Pediatric Use** The safety and effectiveness of ZYVOX for the treatment of pediatric patients with the following infections are supported by evidence from adequate and well-controlled studies in adults, pharmacokinetic data in pediatric patients, and additional data from a comparator-controlled study of Gram-positive infections in pediatric patients ranging in age from birth through 11 years (see **INDICATIONS AND USAGE**): nosocomial pneumonia, complicated skin and skin structure infections, community-acquired pneumonia (also supported by evidence from an uncontrolled study in patients ranging in age from 8 months through 12 years), vancomycin-resistant *Enterococcus faecium* infections. The safety and effectiveness of ZYVOX for the treatment of pediatric patients with the following infection have been established in a comparator-controlled study in pediatric patients ranging in age from 5 through 17 years: uncomplicated skin and skin structure infections caused by *Staphylococcus aureus* (methicillin-susceptible strains only) or *Streptococcus pyogenes*. Pharmacokinetic information generated in pediatric patients with ventriculoperitoneal shunts showed variable cerebrospinal fluid (CSF) linezolid concentrations following single and multiple dosing of linezolid; therapeutic concentrations were not consistently achieved or maintained in the CSF. Therefore, the use of linezolid for the empiric treatment of pediatric patients with central nervous system infections is not recommended. The C_{max} and the volume of distribution (V_{ss}) of linezolid are similar regardless of age in pediatric patients. However, linezolid clearance is a function of age. Excluding neonates less than a week of age, clearance is most rapid in the youngest age groups ranging from >1 week old to 11 years, resulting in lower single-dose systemic exposure (AUC) and shorter half-life as compared with adults. As age of pediatric patients increases, the clearance of linezolid gradually decreases, and by adolescence, mean clearance values approach those observed for the adult population. There is wider inter-subject variability in linezolid clearance and in systemic drug exposure (AUC) across all pediatric age groups as compared with adults. Similar mean daily AUC values were observed in pediatric patients from birth to 11 years of age dosed q8h relative to adolescents or adults dosed q12h. Therefore, the dosage for pediatric patients up to 11 years of age should be 10 mg/kg q8h. Pediatric patients 12 years and older should receive 600 mg q12h. Recommendations for the dosage regimen for pre-term neonates less than 7 days of age (gestational age less than 34 weeks) are based on pharmacokinetic data from 9 pre-term neonates. Most of these pre-term neonates have lower systemic linezolid clearance values and larger AUC

Periprocedural bleeding, the primary outcome for this study, occurred in more than 30,000 patients (2%). This was defined as bleeding that required a blood transfusion or a prolonged hospital stay, or bleeding that was associated with a greater than 3-g/dL decline in hemoglobin level.

Bleeding events occurred in 2.1% of patients in whom vascular closure devices were used, 1.6% who received bivalirudin, and 0.9% who received both preventive strategies. In comparison, bleeding events occurred in 2.8% of patients who received manual compression.

Independently of preprocedural risk of bleeding, vascular closure devices were associated with 6.7 fewer bleeding events per 1,000 patients, bivalirudin was associated with 8.5 fewer events per 1,000, and the combination of both devices and bivalirudin was associated with 14.2 fewer events, compared with manual compression.

In a further analysis of the data, patients

were stratified according to their estimated risk for bleeding before the procedure began. As bleeding risk increased, differences in actual bleeding rates between the four strategies became more pronounced. Among the highest-risk patients, “the use of vascular closure devices plus bivalirudin was associated with an absolute 3.8% lower bleed-

Targeting bleeding complications ‘holds great potential for improving the safety and cost-effectiveness’ of percutaneous coronary interventions.

ing rate, which translates into an estimated number needed to treat of 33 to prevent 1 bleeding event, as compared with manual compression,” Dr. Marso and his colleagues wrote (JAMA 2010; 303:2156-64).

A total of 40.3% of patients at highest risk received manual compression, compared with 30.8% of those with the lowest risk. However, 14.4% of the highest-risk patients received bivalirudin plus vascular closure, compared with 21.0% of the lowest-risk patients.

Translating these study findings into change in clinical practice will be “challenging” for several reasons, the researchers noted.

“First, assessing the risk for bleeding in clinical practice is neither inherently intuitive nor commonly used. Second, physicians have more experience using bivalirudin in lower-risk patients, since it was first studied in patients undergoing elective PCI and only recently in higher-risk patients,” they said.

In addition, some patients are not suited to one or the other of these strategies. Bivalirudin is not recommended in those taking anticoagulants or those who have a chronic total occlusion, and closure devices are not recommended in patients with certain anatomical limitations such as severe calcification or peripheral artery disease.

Disclosures: Dr. Marso and his associates reported receiving support from many sources, including The Medicines Company (maker of bivalirudin), Amylin Pharmaceuticals, Boston Scientific, Volcano Corp., Terumo Corp., Abbott Vascular, and NovoNordisk.

values than many full-term neonates and older infants. Therefore, these pre-term neonates should be initiated with a dosing regimen of 10 mg/kg q12h. Consideration may be given to the use of a 10 mg/kg q8h regimen in neonates with a sub-optimal clinical response. All neonatal patients should receive 10 mg/kg q8h by 7 days of life. In limited clinical experience, 5 out of 6 (83%) pediatric patients with infections due to Gram-positive pathogens with MICs of 4 µg/mL treated with ZYVOX had clinical cures. However, pediatric patients exhibit wider variability in linezolid clearance and systemic exposure (AUC) compared with adults. In pediatric patients with a sub-optimal clinical response, particularly those with pathogens with MIC of 4 µg/mL, lower systemic exposure, site and severity of infection, and the underlying medical condition should be considered when assessing clinical response. Geriatric Use of the 2046 patients treated with ZYVOX in Phase 3 comparator-controlled clinical trials, 589 (29%) were 65 years or older and 253 (12%) were 75 years or older. No overall differences in safety or effectiveness were observed between these patients and younger patients.

ADVERSE REACTIONS Adult Patients The safety of ZYVOX formulations was evaluated in 2046 adult patients enrolled in seven Phase 3 comparator-controlled clinical trials, who were treated for up to 28 days. In these studies, 85% of the adverse events reported with ZYVOX were described as mild to moderate in intensity. The incidence (%) of adverse events reported in at least 2% of patients treated with either ZYVOX (n=2046) or all comparators* (n=2001) in these trials were as follows: diarrhea 8.5 and 6.3; headache 6.5 and 5.5; nausea 6.2 and 4.6; vomiting 3.7 and 2.0; insomnia 2.5 and 1.7; constipation 2.2 and 2.1; rash 2.0 and 2.2; dizziness 2.0 and 1.9; and fever 1.6 and 2.1 respectively. The most common adverse events in patients treated with ZYVOX were diarrhea (incidence across studies: 2.8% to 11.0%), headache (incidence across studies: 0.5% to 11.3%), and nausea (incidence across studies: 3.4% to 9.6%). The percent of drug-related adverse events in at least 1% of adult patients in a trial involving the treatment of uncomplicated skin and skin structure infection comparing ZYVOX 400 mg q12h (n=548) to clarithromycin 250 mg q12h (n=537) were 25.4 and 19.6 respectively. The percent of patients discontinuing drug due to drug-related adverse events† were 3.5 and 2.4 respectively. The incidence of drug-related adverse events occurring in >1% of adult patients were diarrhea 5.3 and 4.8; nausea 3.5 and 3.5; headache 2.7 and 2.2; taste alteration 1.8 and 2.0; vaginal moniliasis 1.6 and 1.3; fungal infection 1.5 and 0.2; abnormal liver function tests 0.4 and 0.0; vomiting 0.9 and 0.4; tongue discoloration 1.1 and 0.0; dizziness 1.1 and 1.5; and oral moniliasis 0.4 and 0.0 respectively. The percent of drug-related adverse events in at least 1% of adult patients in all other indications of ZYVOX 600 mg q12h (n=1498) versus all other comparators‡ (n=1464) with at least 1 drug-related adverse event was 20.4 and 14.3 respectively. The percent of adult patients discontinuing due to drug-related adverse events‡ was 2.1 and 1.7 respectively. The incidence of drug-related adverse events occurring in >1% of adult patients were diarrhea 4.0 and 2.7; nausea 3.3 and 1.8; headache 1.9 and 1.0; taste alteration 0.9 and 0.2; vaginal moniliasis 1.0 and 0.4; fungal infection 0.1 and <0.1; abnormal liver function tests 1.3 and 0.5; vomiting 1.2 and 0.4; tongue discoloration 0.2 and 0.0; dizziness 0.4 and 0.3; and oral moniliasis 1.1 and 0.4. Other adverse events reported in Phase 2 and Phase 3 studies included oral moniliasis, vaginal moniliasis, hypertension, dyspepsia, localized abdominal pain, pruritus, and tongue discoloration. **Pediatric Patients** The safety of ZYVOX formulations was evaluated in 215 pediatric patients ranging in age from birth through 11 years, and in 248 pediatric patients aged 5 through 17 years (146 of these 248 were age 5 through 11 and 102 were age 12 to 17). These patients were enrolled in two Phase 3 comparator-controlled clinical trials and were treated for up to 28 days. In these studies, 83% and 99%, respectively, of the adverse events reported with ZYVOX were described as mild to moderate in intensity. In the study of hospitalized pediatric patients (birth through 11 years) with Gram-positive infections, who were randomized 2 to 1 (linezolid:vancomycin), mortality was 6.0% (13/215) in the linezolid arm and 3.0% (3/101) in the vancomycin arm. However, given the severe underlying illness in the patient population, no causality could be established. The incidence of adverse events reported in ≥2% of pediatric patients treated for uncomplicated skin and skin structure infections† with ZYVOX (n=248) or cefadroxil (n=251) were fever 2.9 and 3.6; diarrhea 7.8 and 8.0; vomiting 2.9 and 6.4; rash 1.6 and 1.2; headache 6.5 and 4.0; upper respiratory infection 3.7 and 5.2; nausea 3.7 and 3.2; trauma 3.3 and 4.8; pharyngitis 2.9 and 1.6; cough 2.4 and 4.0; generalized abdominal pain 2.4 and 2.8; localized abdominal pain 2.4 and 2.8; loose stools 1.6 and 0.8; localized pain 2.0 and 1.6; skin disorder 2.0 and 0.0 respectively. The incidence of adverse events reported in ≥2% of pediatric patients treated for all other indications† with either ZYVOX (n=215) or vancomycin (n=101) in comparator-controlled trials were fever 14.1 and 14.1; diarrhea 10.8 and 12.1; vomiting 9.4 and 9.1; sepsis 8.0 and 7.1; rash 7.0 and 15.2; headache 0.9 and 0.0; anemia 5.6 and 7.1; thrombocytopenia 4.7 and 2.0; upper respiratory infection 4.2 and 1.0; nausea 1.9 and 0.0; dyspnea 3.3 and 1.0; reaction at site of injection or of vascular catheter 3.3 and 5.1; trauma 2.8 and 2.0; pharyngitis 0.5 and 1.0; convulsion 2.8 and 2.0; hypokalemia 2.8 and 3.0; pneumonia 2.8 and 2.0; thrombocytopenia 2.8 and 2.0; cough 0.9 and 0.0; generalized abdominal pain 0.9 and 2.0; localized abdominal pain 0.5 and 1.0; apnea 2.3 and 2.0; gastrointestinal bleeding 2.3 and 1.0; generalized edema 2.3 and 1.0; loose stools 2.3 and 3.0; localized pain 0.9 and 0.0; and skin disorder 0.9 and 1.0. The percent of pediatric patients treated for uncomplicated skin and skin structure infections† with either ZYVOX (n=248) or cefadroxil (n=251) and with ≥1 drug-related adverse event occurring in more than 1% of patients were 19.2 and 14.1 respectively. The percent of pediatric patients discontinuing due to a drug-related adverse event was 1.6 and 2.4 respectively. The incidence of drug-related adverse events reported in more than 1% of pediatric patients (and more than 1 patient) were diarrhea 5.7 and 5.2; nausea 3.3 and 2.0; headache 2.4 and 0.8; loose stools 1.2 and 0.8; vomiting 1.2 and 2.4; generalized abdominal pain 1.6 and 1.2; localized abdominal pain 1.6 and 1.2; eosinophilia 0.4 and 0.4; rash 0.4 and 1.2; vertigo 1.2 and 0.4 and pruritus at non-application site 0.4 and 0.0 respectively. The percent of pediatric patients treated for all other indications† with either ZYVOX (n=215) or vancomycin (n=101) and with ≥1 drug-related adverse event occurring in more than 1% of patients were 18.8 and 34.3 respectively. The percent of patients discontinuing due to a drug-related adverse event were 0.9 and 6.1 respectively. The incidence of drug-related adverse events reported in more than 1% of pediatric patients (and more than 1 patient) were diarrhea 3.8 and 6.1; nausea 1.4 and 0.0; loose stools 1.9 and 0.0; thrombocytopenia 1.9 and 0.0; vomiting 1.9 and 1.0; anemia 1.4 and 1.0; eosinophilia 1.4 and 0.0; rash 1.4 and 7.1; oral moniliasis 0.9 and 4.0; fever 0.5 and 3.0; pruritus at non-application site 0.0 and 2.0; and anaphylaxis 0.0 and 10.1† respectively. **Laboratory Changes ZYVOX** has been associated with thrombocytopenia when used in doses up to and including 600 mg every 12 hours for up to 28 days. In Phase 3 comparator-controlled trials, the percentage of adult patients who developed a substantially low platelet count (defined as less than 75% of lower limit of normal and/or baseline) was 2.4% (range among studies: 0.3 to 10.0%) with ZYVOX and 1.5% (range among studies: 0.4 to 7.0%) with a comparator. In a study of hospitalized pediatric patients ranging in age from birth through 11 years, the percentage of patients who developed a substantially low platelet count (defined as less than 75% of lower limit of normal and/or baseline) was 12.9% with ZYVOX and 13.4% with vancomycin. In an outpatient study of pediatric patients aged from 5 through 17 years, the percentage of patients who developed a substantially low platelet count was 0% with ZYVOX and 0.4% with cefadroxil. Thrombocytopenia associated with the use of ZYVOX appears to be dependent on duration of therapy, generally greater than 2 weeks of treatment. The platelet counts for most patients returned to the normal range/baseline during the follow-up period. No related clinical adverse events were

identified in Phase 3 clinical trials in patients developing thrombocytopenia. Bleeding events were identified in thrombocytopenic patients in a compassionate use program for ZYVOX; the role of linezolid in these events cannot be determined (see **WARNINGS**). Changes seen in other laboratory parameters, without regard to drug relationship, revealed no substantial differences between ZYVOX and the comparators. These changes were generally not clinically significant, did not lead to discontinuation of therapy, and were reversible. The percent of adult patients with at least one substantially abnormal hematologic‡ value in patients treated with ZYVOX 400 mg q12h or clarithromycin 250 mg q12h for uncomplicated skin and skin structure infections were as follows: hemoglobin (g/dL) 0.9 and 0.0; platelet count (x 10³/mm³) 0.7 and 0.8; WBC (x 10³/mm³) 0.2 and 0.6; neutrophils (x 10³/mm³) 0.0 and 0.2 respectively. The percent of adult patients with at least one substantially abnormal hematologic‡ value in patients treated with ZYVOX 600 mg q12h or a comparator§ were as follows: hemoglobin (g/dL) 7.1 and 6.6; platelet count (x 10³/mm³) 3.0 and 1.8; WBC (x 10³/mm³) 2.2 and 1.3 and neutrophils (x 10³/mm³) 1.1 and 1.2 respectively. The percent of adult patients with at least one substantially abnormal serum chemistry†† value in patients treated with ZYVOX 400 mg q12h or clarithromycin 250 mg q12h for uncomplicated skin and skin structure infections were as follows: AST (U/L) 1.7 and 1.3; ALT (U/L) 1.7 and 1.7; LDH (U/L) 0.2 and 0.2; alkaline phosphatase (U/L) 0.2 and 0.2; lipase (U/L) 2.8 and 2.6; amylase (U/L) 0.2 and 0.2; total bilirubin (mg/dL) 0.2 and 0.0; BUN (mg/dL) 0.2 and 0.0; and creatinine (mg/dL) 0.2 and 0.0 respectively. The percent of adult patients with at least one substantially abnormal serum chemistry†† value in patients treated with ZYVOX 600 mg q12h or a comparator§ were as follows: AST (U/L) 5.0 and 6.8; ALT (U/L) 9.6 and 9.3; LDH (U/L) 1.8 and 1.5; alkaline phosphatase (U/L) 3.5 and 3.1; lipase (U/L) 4.3 and 4.2; amylase (U/L) 2.4 and 2.0; total bilirubin (mg/dL) 0.9 and 1.1; BUN (mg/dL) 2.1 and 1.5; and creatinine (mg/dL) 0.2 and 0.6 respectively. The percent of pediatric patients with at least one substantially abnormal hematologic‡ value in patients treated with ZYVOX or cefadroxil for uncomplicated skin and skin structure infections§ were as follows: hemoglobin (g/dL) 0.0 and 0.0; platelet count (x 10³/mm³) 0.0 and 0.4; WBC (x 10³/mm³) 0.8 and 0.8; neutrophils (x 10³/mm³) 1.2 and 0.8 respectively. The percent of pediatric patients with at least one substantially abnormal hematologic‡ value in patients treated with ZYVOX or vancomycin for any other indication¶ were as follows: hemoglobin (g/dL) 15.7 and 12.4; platelet count (x 10³/mm³) 12.9 and 13.4; WBC (x 10³/mm³) 12.4 and 10.3 and neutrophils (x 10³/mm³) 5.9 and 4.3 respectively. The percent of pediatric patients with at least one substantially abnormal serum chemistry¶ value in patients treated with ZYVOX or cefadroxil for uncomplicated skin and skin structure infections§ were as follows: ALT (U/L) 0.0 and 0.0; lipase (U/L) 0.4 and 1.2; and creatinine (mg/dL) 0.4 and 0.0 respectively. The percent of pediatric patients with at least one substantially abnormal serum chemistry¶ value in patients treated with ZYVOX or vancomycin for any other indication¶ were as follows: ALT (U/L) 10.1 and 12.5; amylase (U/L) 0.6 and 1.3; total bilirubin (mg/dL) 6.3 and 5.2; and creatinine (mg/dL) 2.4 and 1.0 respectively. **Postmarketing Experience** Myelosuppression (including anemia, leukopenia, pancytopenia, and thrombocytopenia) has been reported during postmarketing use of ZYVOX (see **WARNINGS**). Peripheral neuropathy, and optic neuropathy sometimes progressing to loss of vision, have been reported in patients treated with ZYVOX. Lactic acidosis has been reported with the use of ZYVOX (see **PRECAUTIONS**). Although these reports have primarily been in patients treated for longer than the maximum recommended duration of 28 days, these events have also been reported in patients receiving shorter courses of therapy. Serotonin syndrome has been reported in patients receiving concomitant serotonergic agents, including antidepressants such as selective serotonin reuptake inhibitors (SSRIs) and ZYVOX (see **PRECAUTIONS**). Convulsions have been reported with the use of ZYVOX (see **PRECAUTIONS**). Anaphylaxis, angioedema, and bullous skin disorders such as those described as Stevens Johnson syndrome have been reported. These events have been chosen for inclusion due to either their seriousness, frequency of reporting, possible causal connection to ZYVOX, or a combination of these factors. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made and causal relationship cannot be precisely established. **OVERDOSAGE** In the event of overdosage, supportive care is advised, with maintenance of glomerular filtration. Hemodialysis may facilitate more rapid elimination of linezolid. In a Phase 1 clinical trial, approximately 30% of a dose of linezolid was removed during a 3-hour hemodialysis session beginning 3 hours after the dose of linezolid was administered. Data are not available for removal of linezolid with peritoneal dialysis or hemoperfusion. Clinical signs of acute toxicity in animals were decreased activity and ataxia in rats and vomiting and tremors in dogs treated with 3000 mg/kg/day and 2000 mg/kg/day, respectively.

* MDRSP refers to isolates resistant to 2 or more of the following antibiotics: penicillin, second-generation cephalosporins, macrolides, tetracycline, and trimethoprim/sulfamethoxazole.
† Comparators included cefpodoxime proxetil 200 mg PO q12h; ceftriaxone 1 g IV q12h; clarithromycin 250 mg PO q12h; dicloxacillin 500 mg PO q6h; oxacillin 2 g IV q6h; vancomycin 1 g IV q12h.
‡ The most commonly reported drug-related adverse events leading to discontinuation in patients treated with ZYVOX were nausea, headache, diarrhea, and vomiting.
§ Comparators included cefpodoxime proxetil 200 mg PO q12h; ceftriaxone 1 g IV q12h; dicloxacillin 500 mg PO q6h; oxacillin 2 g IV q6h; vancomycin 1 g IV q12h.
¶ Patients 5 through 11 years of age received ZYVOX 10 mg/kg PO q12h or cefadroxil 15 mg/kg PO q12h. Patients 12 years or older received ZYVOX 600 mg PO q12h or cefadroxil 500 mg PO q12h.
†† Patients from birth through 11 years of age received ZYVOX 10 mg/kg IV/PO q8h or vancomycin 10 to 15 mg/kg IV q6-24h, depending on age and renal clearance.
‡‡ These reports were of ‘red-man syndrome,’ which were coded as anaphylaxis.
§§ <75% (<50% for neutrophils) of Lower Limit of Normal (LLN) for values normal at baseline; <75% (<50% for neutrophils) of LLN and of baseline for values abnormal at baseline.
¶¶ >2 x Upper Limit of Normal (ULN) for values normal at baseline; >2 x ULN and >2 x baseline for values abnormal at baseline.
*** <75% (<50% for neutrophils) of Lower Limit of Normal (LLN) for values normal at baseline; <75% (<50% for neutrophils) of LLN and <75% (<50% for neutrophils, <90% for hemoglobin) if baseline <LLN) of baseline for values abnormal at baseline.
**** >2 x Upper Limit of Normal (ULN) for values normal at baseline; >2 x ULN and >2 (x1.5 for total bilirubin) x baseline for values abnormal at baseline.
Rx only Rev. May 2008

Hospital Quality Data Application

The Agency for Healthcare Research and Quality is offering a free, Microsoft Windows-based software application to help hospitals compile, analyze, and post hospital quality data.

MONAHRQ (My Own Network Powered by AHRQ) can be used to create custom Web sites that include information about patient safety, mortality rates, discharges, costs, length of stay, and more. For more information, call AHRQ Public Affairs at 301-427-1859 or 301-427-1855. ■

Can't Find Your Last Issue?



You have FREE access to articles from this issue and past issues of HOSPITALIST NEWS at www.ehospitalistnews.com