

Many Black MI Patients Skip Regular Checks

BY MARY ELLEN SCHNEIDER
New York Bureau

More than two-thirds of African American patients who have suffered a myocardial infarction say the event was a “wake-up call,” but a quarter of patients also report that they did not see their physician regularly after the attack, according to a survey released by the National Medical Association.

“Obviously, there’s a disconnect here,” said Dr. Clyde W. Yancy, medical director of the Heart and Vascular Institute at Baylor University Medical Center in Dallas.

Physicians and researchers need to better understand this contradiction because it’s an opportunity to improve outcomes among African American patients, Dr. Yancy said during a teleconference sponsored by the National Medical Association (NMA) and supported by GlaxoSmithKline.

The survey, which was commissioned by the NMA and supported by GlaxoSmithKline, was conducted online among 502 African American adults aged 18 and older who had experienced a myocardial infarction.

African Americans have a significantly higher risk for virtually every cardiovascular disease than their white counterparts, Dr. Yancy said. And when it comes to myocardial infarction, African American men have the highest incidence of first heart attacks, followed by white men, and closely followed by African American women.

But despite the increased risk, there is a lack of awareness. “The perception is that the African American community is not at risk, particularly for heart attacks. Awareness needs to be elevated in a major way.”

The NMA survey showed that most respondents saw their myocardial infarction as a significant event, with 64% saying they felt that they had been given a second chance at life, and 46% saying that they were significantly worried about having another heart attack.

However, the survey also found that they were not taking steps to avoid another cardiac event. For example, 22% of respondents reported not taking medication exactly as prescribed and 21% said that they do not monitor their eating habits.

The survey results also revealed that African American patients are in need of increased support in the period following a myocardial infarction. Fewer than half of respondents (47%) said they had family and friends who remind them to take their medications and 27% said they did not feel knowledgeable about how to manage their health after an attack.

Part of the problem may come down to socioeconomic factors, Dr. Yancy said. Patients may be neglecting their medications and physician visits because they lack the resources and support. Other factors include a possible lack of trust of physicians by African American patients, a belief that they have not received the best medicine, and a lack of education about potential side effects. ■

Post-AMI Anxiety More Common in Women

BY MITCHEL L. ZOLER
Philadelphia Bureau

CHICAGO — The high anxiety that many women have after an acute myocardial infarction may explain their high complication rate, compared with men, and it may offer a new way to improve patient outcomes.

“Early recognition and effective treatment of anxiety immediately after a myocardial infarction may decrease morbidity and mortality,” Debra K. Moser,

D.N.Sc., said during a poster presentation at the annual scientific sessions of the American Heart Association. “Anxiety is a target for intervention.” In AMI patients, “one of the things most lacking in clinical cardiology today is the assessment of anxiety and depression.”

The Brief Symptom Inventory (BSI) is the way that Dr. Moser and her associates have measured anxiety in patients after an acute myocardial infarction (AMI), and treatment can include an anxiolytic drug as well as psychosocial interventions such

as cognitive-behavioral therapy, said Dr. Moser, professor of nursing at the University of Kentucky, Lexington.

She and her associates examined the role of anxiety after AMI in 635 men and 244 women enrolled at several centers in the United States and four other countries. The patients’ anxiety levels were measured, using the BSI, within 72 hours of their hospital admission for an AMI.

The BSI is a reliable and validated measure that uses six questions to measure anxiety, said Dr. Moser. Each question is

FIRST
IN A NOVEL
CLASS OF
SLEEP
AGENTS

Rozerem is indicated for the treatment of insomnia characterized by difficulty with sleep onset. Rozerem can be prescribed for long-term use. Rozerem should not be used in patients with hypersensitivity to any components of the formulation, severe hepatic impairment, or in combination with fluvoxamine. Failure of insomnia to remit after a reasonable period of time should be medically evaluated, as this may be the result of an unrecognized underlying medical disorder. Hypnotics should be administered with caution to patients exhibiting signs and symptoms of depression. Rozerem has not been studied in patients with severe sleep apnea, severe COPD, or in children or adolescents. The effects in these populations are unknown. Avoid taking Rozerem with alcohol. Rozerem has been associated with decreased testosterone levels and increased prolactin levels. Health professionals should be mindful of any unexplained symptoms possibly associated with such changes in these hormone levels. Rozerem should not be taken with or immediately after a high-fat meal. Rozerem should be taken within 30 minutes before going to bed and activities confined to preparing for bed. The most common adverse events seen with Rozerem that had at least a 2% incidence difference from placebo were somnolence, dizziness, and fatigue.

Please see adjacent Brief Summary of Prescribing Information.