

CLINICAL CAPSULES

Cannabis Use May Predict Later Anxiety

Young people who use cannabis at 15 years of age are significantly more likely to show symptoms of anxiety and depression in young adulthood, compared with non-cannabis users, data show.

From a clinical standpoint, reducing cannabis use in adolescents could reduce anxiety and depression in young adulthood, wrote Dr. Mohammad R. Hayatbakhsh and his colleagues at the University of Queensland in Brisbane, Australia.

To assess the relationships among cannabis use and anxiety and depression,

the researchers reviewed data on 3,239 Australian young people from birth to age 21 and measured confounding factors at birth, age 14, and age 21 (J. Am. Acad. Child Adolesc. Psychiatry 2007;46:408-17).

A total of 1,586 youth (49%) reported ever using cannabis; 36.8% were occasional users (an average of one use per month) and 12.2% were frequent users (an average of one use every few days). Symptoms of anxiety and depression were assessed using the Youth Self-Report questionnaire.

After the researchers controlled for multiple factors, including gender, mother's

mental health, family income, and smoking and alcohol consumption, young people who were frequent cannabis users before 15 years of age were more than twice as likely to show signs of anxiety and depression as young adults than were non-cannabis users. The association persisted, whether or not the young people used other illicit drugs in addition to cannabis.

Conversely, the presence of anxiety and depression before age 15 was not significantly associated with cannabis use in young adulthood. However, the findings suggest that cannabis use in adolescence may predict mental health problems later in life, the researchers noted.

Risky Sex Linked to Sensation-Seeking
Black adolescent girls who reported risky sexual activities scored significantly higher on tests of sexual sensation-seeking traits than did those who didn't report risky sex, based on data from 1,245 sexually active females aged 15-21 years.

To examine the relationship between sexual sensation seeking and sexual risk taking in adolescents, Joshua Spitalnick, Ph.D., of Emory University, Atlanta, and his colleagues surveyed black girls who were seeking sexual health services at community health clinics (J. Adolescence 2007;30:165-73).

Most (84%) of these adolescents reported that they were involved in a sexual relationship. The average length of that relationship was 15 months, and the respondents reported an average of nine lifetime partners. The average age of the participants was 18 years.

They were evaluated using the Sexual Sensation Seeking for Adolescents (SSSA) scale, which included statements such as, "I enjoy the thrill of having sex in public places," with answer choices ranging from 1 (strongly disagree) to 4 (strongly agree). Scores ranged from 9 to 36, with an average score of 17.53.

Overall, respondents with high SSSA scores were significantly more likely to report risky sexual behavior including more frequent vaginal intercourse, more sexual partners, and inconsistent use of condoms or other protection than were girls who scored lower on the SSSA.

Delinquent Teens at Risk for Suicide

Teenage delinquency was significantly associated with an increased risk for suicidal behavior in girls, according to data from a nationally representative sample of American teens.

To prospectively examine the association between delinquency and suicide in teens, Martie P. Thompson, Ph.D., and her colleagues at Clemson (S.C.) University reviewed data on 15,034 teens aged 12-17 years from the National Longitudinal Study of Adolescent Health, a survey of factors that affect teens' health and behavior (J. Adolesc. Health 2007;40:232-7).

Delinquency was assessed using a 15-item survey of behaviors in the previous 12 months, with questions such as, "How often did you deliberately damage property that didn't belong to you?"

After the researchers controlled for age, race, gender, and urban dwelling status, delinquent teens were significantly more likely than their nondelinquent peers to report suicidal ideation, suicide attempts, and treatment for suicide attempts at both 1 and 7 years' follow-up. When the researchers controlled for behavioral risk factors such as depression, impulsivity, religiosity, and problem drinking, delinquency remained significantly associated with suicidal ideation 1 year later and with suicide attempts 1 and 7 years later among girls. The association for boys remained but was not statistically significant.

Minority teens were less likely than were white teens to report suicidal ideation after 1 year, but there were no significant racial differences in those who attempted suicide or who required medical treatment after attempting suicide, Dr. Thompson and her associates noted.

—Heidi Splette



Brief Summary of Prescribing Information
05-1114

ROZEREM™

(ramelteon) Tablets

INDICATIONS AND USAGE

ROZEREM is indicated for the treatment of insomnia characterized by difficulty with sleep onset.

CONTRAINDICATIONS

ROZEREM is contraindicated in patients with a hypersensitivity to ramelteon or any components of the ROZEREM formulation.

WARNINGS

Since sleep disturbances may be the presenting manifestation of a physical and/or psychiatric disorder, symptomatic treatment of insomnia should be initiated only after a careful evaluation of the patient. The failure of insomnia to remit after a reasonable period of treatment may indicate the presence of a primary psychiatric and/or medical illness that should be evaluated. Worsening of insomnia, or the emergence of new cognitive or behavioral abnormalities, may be the result of an unrecognized underlying psychiatric or physical disorder and requires further evaluation of the patient. As with other hypnotics, exacerbation of insomnia and emergence of cognitive and behavioral abnormalities were seen with ROZEREM during the clinical development program.

ROZEREM should not be used by patients with severe hepatic impairment.

ROZEREM should not be used in combination with fluvoxamine (see PRECAUTIONS: Drug Interactions).

A variety of cognitive and behavior changes have been reported to occur in association with the use of hypnotics. In primarily depressed patients, worsening of depression, including suicidal ideation, has been reported in association with the use of hypnotics.

Patients should avoid engaging in hazardous activities that require concentration (such as operating a motor vehicle or heavy machinery) after taking ROZEREM.

After taking ROZEREM, patients should confine their activities to those necessary to prepare for bed.

PRECAUTIONS

General

ROZEREM has not been studied in subjects with severe sleep apnea or severe COPD and is not recommended for use in those populations.

Patients should be advised to exercise caution if they consume alcohol in combination with ROZEREM.

Use in Adolescents and Children

ROZEREM has been associated with an effect on reproductive hormones in adults, e.g. decreased testosterone levels and increased prolactin levels. It is not known what effect chronic or even chronic intermittent use of ROZEREM may have on the reproductive axis in developing humans (see Pediatric Use).

Information for Patients

Patients should be advised to take ROZEREM within 30 minutes prior to going to bed and should confine their activities to those necessary to prepare for bed.

Patients should be advised to avoid engaging in hazardous activities (such as operating a motor vehicle or heavy machinery) after taking ROZEREM.

Patients should be advised that they should not take ROZEREM with or immediately after a high fat meal.

Patients should be advised to consult their health care provider if they experience worsening of insomnia or any new behavioral signs or symptoms of concern.

Patients should consult their health care provider if they experience one of the following: cessation of menses or galactorrhea in females, decreased libido, or problems with fertility.

Laboratory Tests

No standard monitoring is required.

For patients presenting with unexplained amenorrhea, galactorrhea, decreased libido, or problems with fertility, assessment of prolactin levels and testosterone levels should be considered as appropriate.

Drug Interactions

ROZEREM has a highly variable inter-subject pharmacokinetic profile (approximately 100% coefficient of variation in C_{max} and AUC). As noted above, CYP1A2 is the major isozyme involved in the metabolism of ROZEREM; the CYP2C subfamily and CYP3A4 isozymes are also involved to a minor degree.

Effects of Other Drugs on ROZEREM Metabolism

Fluvoxamine (strong CYP1A2 inhibitor): When fluvoxamine 100 mg twice daily was administered for 3 days prior to single-dose co-administration of ROZEREM 16 mg and fluvoxamine, the AUC_{0-24} for ramelteon increased approximately 190-fold, and the C_{max} increased approximately 70-fold, compared to ROZEREM administered alone. ROZEREM should not be used in combination with fluvoxamine (See WARNINGS). Other less potent CYP1A2 inhibitors have not been adequately studied. ROZEREM should be administered with caution to patients taking less strong CYP1A2 inhibitors.

Rifampin (strong CYP enzyme inducer): Administration of rifampin 600 mg once daily for 11 days resulted in a mean decrease of approximately 80% (40% to 90%) in total exposure to ramelteon and metabolite M-II, (both AUC_{0-24} and C_{max}) after a single 32 mg dose of ROZEREM. Efficacy may be reduced when ROZEREM is used in combination with strong CYP enzyme inducers such as rifampin.

Ketoconazole (strong CYP3A4 inhibitor): The AUC_{0-24} and C_{max} of ramelteon increased by approximately 84% and 36%, respectively, when a single 16 mg dose of ROZEREM was administered on the fourth day of ketoconazole 200 mg twice daily administration, compared to administration of ROZEREM alone. Similar increases were seen in M-II pharmacokinetic variables. ROZEREM should be administered with caution in subjects taking strong CYP3A4 inhibitors such as ketoconazole.

Flucanazole (strong CYP2C9 inhibitor): The total and peak systemic exposure (AUC_{0-24} and C_{max}) of ramelteon after a single 16 mg dose of ROZEREM was increased by approximately 150% when administered with flucanazole. Similar increases were also seen in M-II exposure. ROZEREM should be administered with caution in subjects taking strong CYP2C9 inhibitors such as flucanazole.

Interaction studies of concomitant administration of ROZEREM with fluoxetine (CYP2D6 inhibitor), omeprazole (CYP1A2 inducer/CYP2C19 inhibitor), theophylline (CYP1A2 substrate), and dextromethorphan (CYP2D6 substrate) did not produce clinically meaningful changes in either peak or total exposures to ramelteon or the M-II metabolite.

Effects of ROZEREM on Metabolism of Other Drugs

Concomitant administration of ROZEREM with omeprazole (CYP2C19 substrate), dextromethorphan (CYP2D6 substrate), midazolam (CYP3A4 substrate), theophylline (CYP1A2 substrate), digoxin (p-glycoprotein substrate), and warfarin (CYP2C9 [S]/CYP1A2 [R] substrate) did not produce clinically meaningful changes in peak and total exposures to these drugs.

Effect of Alcohol on Rozerem
Alcohol: With single-dose, daytime co-administration of ROZEREM 32 mg and alcohol (0.6 g/kg), there were no clinically meaningful or statistically sig-

nificant effects on peak or total exposure to ROZEREM. However, an additive effect was seen on some measures of psychomotor performance (i.e., the Digit Symbol Substitution Test, the Psychomotor Vigilance Task Test, and a Visual Analog Scale of sedation) at some post-dose time points. No additive effect was seen on the Delayed Word Recognition Test. Because alcohol by itself impairs performance, and the intended effect of ROZEREM is to promote sleep, patients should be cautioned not to consume alcohol when using ROZEREM.

Drug/Laboratory Test Interactions

ROZEREM is not known to interfere with commonly used clinical laboratory tests. In addition, *in vitro* data indicate that ramelteon does not cause false-positive results for benzodiazepines, opiates, barbiturates, cocaine, cannabinoids, or amphetamines in two standard urine drug screening methods *in vitro*.

Carcinogenesis, Mutagenesis, and Impairment of Fertility

Carcinogenesis

In a two-year carcinogenicity study, B6C3F₁ mice were administered ramelteon at doses of 0, 30, 100, 300, or 1000 mg/kg/day by oral gavage. Male mice exhibited a dose-related increase in the incidence of hepatic tumors at dose levels ≥ 100 mg/kg/day including hepatic adenoma, hepatic carcinoma, and hepatoblastoma. Females developed a dose-related increase in the incidence of hepatic adenomas at dose levels ≥ 300 mg/kg/day and hepatic carcinoma at the 1000 mg/kg/day dose level. The no-effect level for hepatic tumors in male mice was 30 mg/kg/day (103-times and 3-times the therapeutic exposure to ramelteon and the active metabolite M-II, respectively, at the maximum recommended human dose [MRHD] based on an area-under-the-curve [AUC] comparison). The no-effect level for hepatic tumors in female mice was 100 mg/kg/day (827-times and 12-times the therapeutic exposure to ramelteon and M-II, respectively, at the MRHD based on AUC).

In a two-year carcinogenicity study conducted in the Sprague-Dawley rat, male and female rats were administered ramelteon at doses of 0, 15, 60, 250 or 1000 mg/kg/day by oral gavage. Male rats exhibited a dose-related increase in the incidence of hepatic adenoma and benign Leydig cell tumors of the testis at dose levels ≥ 250 mg/kg/day and hepatic carcinoma at the 1000 mg/kg/day dose level. Female rats exhibited a dose-related increase in the incidence of hepatic adenoma at dose levels ≥ 60 mg/kg/day and hepatic carcinoma at the 1000 mg/kg/day dose level. The no-effect level for hepatic tumors and benign Leydig cell tumors in male rats was 60 mg/kg/day (1.429-times and 12-times the therapeutic exposure to ramelteon and M-II, respectively, at the MRHD based on AUC). The no-effect level for hepatic tumors in female rats was 15 mg/kg/day (472-times and 16-times the therapeutic exposure to ramelteon and M-II, respectively, at the MRHD based on AUC).

The development of hepatic tumors in rodents following chronic treatment with non-genotoxic compounds may be secondary to microsomal enzyme induction, a mechanism for tumor generation not thought to occur in humans. Leydig cell tumor development following treatment with non-genotoxic compounds in rodents has been linked to reductions in circulating testosterone levels with compensatory increases in luteinizing hormone release, which is a known proliferative stimulus to Leydig cells in the rat testis. Rat Leydig cells are more sensitive to the stimulatory effects of luteinizing hormone than human Leydig cells. In mechanistic studies conducted in the rat, daily ramelteon administration at 250 and 1000 mg/kg/day for 4 weeks was associated with a reduction in plasma testosterone levels.

In the same study, luteinizing hormone levels were elevated over a 24 hour period after the last ramelteon dose; however, the ability of this luteinizing hormone finding and its support for the proposed mechanistic explanation was not clearly established.

Although the rodent tumors observed following ramelteon treatment occurred at plasma levels of ramelteon and M-II in excess of mean clinical plasma concentrations at the MRHD, the relevance of both rodent hepatic tumors and benign rat Leydig cell tumors to humans is not known.

Mutagenesis

Ramelteon was not genotoxic in the following: *in vitro* bacterial reverse mutation (Ames) assay, *in vitro* mammalian cell gene mutation assay using the mouse lymphoma TK⁺ cell line; *in vivo/in vitro* unscheduled DNA synthesis assay in rat hepatocytes; and *in vivo* micronucleus assays conducted in mouse and rat. Ramelteon was positive in the chromosomal aberration assay in Chinese hamster lung cells in the presence of S9 metabolic activation.

Separate studies indicated that the concentration of the M-II metabolite formed by the rat liver S9 fraction used in the *in vitro* genetic toxicology studies described above, exceeded the concentration of ramelteon; therefore, the genotoxic potential of the M-II metabolite was also assessed in these studies.

Impairment of Fertility

Ramelteon was administered to male and female Sprague-Dawley rats in an initial fertility and early embryonic development study at dose levels of 6, 60, or 600 mg/kg/day. No effects on male or female mating or fertility were observed with a ramelteon dose up to 600 mg/kg/day (786-times higher than the MRHD on a mg/m² basis). Irregular estrus cycles, reduction in the number of implants, and reduction in the number of live embryos were noted with dosing females at ≥ 60 mg/kg/day (79-times higher than the MRHD on a mg/m² basis). A reduction in the number of corpora lutea occurred at the 600 mg/kg/day dose level. Administration of ramelteon up to 600 mg/kg/day to male rats for 7 weeks had no effect on sperm quality and when the treated male rats were mated with untreated female rats there was no effect on implants or embryos. In a repeat of this study using oral administration of ramelteon at 20, 60 or 200 mg/kg/day for the same study duration, females demonstrated irregular estrus cycles with doses ≥ 60 mg/kg/day, but no effects were seen on implantation or embryo viability. The no-effect dose for fertility endpoints was 20 mg/kg/day in females (26-times the MRHD on a mg/m² basis) and 600 mg/kg/day in males (786-times higher than the MRHD on a mg/m² basis) when considering all studies.

Pregnancy: Pregnancy Category C
Ramelteon has been shown to be a developmental teratogen in the rat when given in doses 197 times higher than the maximum recommended human dose [MRHD] on a mg/m² basis. There are no adequate and well-controlled studies in pregnant women. Ramelteon should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

The effects of ramelteon on embryo-fetal development were assessed in both the rat and rabbit. Pregnant rats were administered ramelteon by oral gavage at doses of 0, 10, 40, 150, or 600 mg/kg/day during gestation days 6-17, which is the period of organogenesis in this species. Evidence of maternal toxicity and fetal teratogenicity was observed at doses greater than or equal to 150 mg/kg/day. Maternal toxicity was chiefly characterized by decreased body weight and, at 600 mg/kg/day, ataxia and decreased spontaneous movement. At maternally toxic doses (150 mg/kg/day or greater), the fetuses demonstrated visceral malformations consisting of diaphragmatic hernia and minor anatomical variations of the skeleton (irregularly shaped scapula). At 600 mg/kg/day, reductions in fetal body weights and malformations including cysts on the external genitalia were additionally observed. The no-effect level for teratogenicity in this study was 40 mg/kg/day (1,892-times and 45-times higher than the therapeutic exposure to ramelteon and the active metabolite M-II, respectively, at the MRHD based on an area-under-the-curve [AUC] comparison). Pregnant rabbits were administered ramelteon by oral gavage at doses of 0, 12, 60, or 300 mg/kg/day during gestation days 6-18, which is the period of organogenesis in this species. Although maternal toxicity was apparent with a ramelteon dose of 300 mg/kg/day, no evidence of fetal effects or teratogenicity was associated with any dose level. The no-effect level for teratogenicity was, therefore, 300 mg/kg/day (11,862-times and 99-times

higher than the therapeutic exposure to ramelteon and M-II, respectively, at the MRHD based on AUC).

The effects of ramelteon on pre- and post-natal development in the rat were studied by administration of ramelteon to the pregnant rat by oral gavage at doses of 0, 30, 100, or 300 mg/kg/day from day 6 of gestation through parturition to postnatal (lactation) day 21, at which time offspring were weaned. Maternal toxicity was noted at doses of 100 mg/kg/day or greater and consisted of reduced body weight gain and increased adrenal gland weight.

Reduced body weight during the post-weaning period was also noticed in the offspring of the groups given 100 mg/kg/day and higher. Offspring in the 300 mg/kg/day group demonstrated physical and developmental delays including delayed eruption of the lower incisors, a delayed acquisition of the righting reflex, and an alteration of emotional response. These delays are often observed in the presence of reduced offspring body weight but may still be indicative of developmental delay. An apparent decrease in the viability of offspring in the 300 mg/kg/day group was likely due to altered maternal behavior and function observed at this dose level. Offspring of the 300 mg/kg/day group also showed evidence of diaphragmatic hernia, a finding observed in the embryo-fetal development study previously described.

There were no effects on the reproductive capacity of offspring and the resulting progeny were not different from those of vehicle-treated offspring. The no-effect level for pre- and postnatal development in this study was 30 mg/kg/day (39-times higher than the MRHD on a mg/m² basis).

Labor and Delivery

The potential effects of ROZEREM on the duration of labor and/or delivery, for either the mother or the fetus, have not been studied. ROZEREM has no established use in labor and delivery.

Nursing Mothers

Ramelteon is secreted into the milk of lactating rats. It is not known whether this drug is excreted in human milk. No clinical studies in nursing mothers have been performed. The use of ROZEREM in nursing mothers is not recommended.

Pediatric Use

Safety and effectiveness of ROZEREM in pediatric patients have not been established. Further study is needed prior to determining that this product may be used safely in pre-pubescent and pubescent patients.

Geriatric Use

A total of 654 subjects in double-blind, placebo-controlled, efficacy trials who received ROZEREM were at least 65 years of age; of these, 199 were 75 years of age or older. No overall differences in safety or efficacy were observed between elderly and younger adult subjects.

ADVERSE REACTIONS

Overview

The data described in this section reflect exposure to ROZEREM in 4251 subjects, including 346 exposed for 6 months or longer, and 473 subjects for one year.

Adverse Reactions Resulting in Discontinuation of Treatment
Five percent of the 3594 individual subjects exposed to ROZEREM in clinical studies discontinued treatment owing to an adverse event, compared with 2% of the 1370 subjects receiving placebo. The most frequent adverse events leading to discontinuation in subjects receiving ROZEREM were somnolence (0.8%), dizziness (0.5%), nausea (0.3%), fatigue (0.3%), headache (0.3%), and insomnia (0.3%).

ROZEREM Most Commonly Observed Adverse Events in Phase 3 trials
The incidence of adverse events during the Phase 1 through 3 trials (% placebo, n=1370; % ramelteon [8 mg], n=1250) were: headache NOS (7%, 7%), somnolence (3%, 5%), fatigue (2%, 4%), dizziness (3%, 5%), nausea (2%, 3%), insomnia exacerbated (2%, 3%), upper respiratory tract infection NOS (2%, 3%), diarrhea NOS (2%, 2%), myalgia (1%, 2%), depression (1%, 2%), dysgeusia (1%, 2%), arthralgia (1%, 2%), influenza (0, 1%), blood cortisol decreased (0, 1%).

Because clinical trials are conducted under widely varying conditions, adverse reaction rates in the clinical trials of a drug cannot be directly compared to rates in clinical trials of other drugs, and may not reflect the rates observed in practice. The adverse reaction information from clinical trials does, however, provide a basis for identifying the adverse events that appear to be related to drug use and for approximating rates.

DRUG ABUSE AND DEPENDENCE

ROZEREM is not a controlled substance.

Human Data: See the CLINICAL TRIALS section, Studies Pertinent to Safety Concerns for Sleep-Promoting Agents in the Complete Prescribing Information.

Animal Data: Ramelteon did not produce any signals from animal behavioral studies indicating that the drug produces rewarding effects. Monkeys did not prefer rates in the clinical trials of a drug cannot be directly compared to rates in clinical trials of other drugs, and may not reflect the rates observed in practice. The adverse reaction information from clinical trials does, however, provide a basis for identifying the adverse events that appear to be related to drug use and for approximating rates.

Discontinuation of ramelteon in animals or in humans after chronic administration did not produce withdrawal signs. Ramelteon does not appear to produce physical dependence.

OVERDOSAGE

Signs and Symptoms

No cases of ROZEREM overdose have been reported during clinical development. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.

ROZEREM was administered in single doses up to 160 mg in an abuse liability trial. No safety or tolerability concerns were seen.