

# Lobular Carcinoma More Alarming Than Thought

BY BRUCE JANCIN  
Denver Bureau

SAN ANTONIO — Lobular carcinoma in situ is not always the indolent disease it has long been considered to be, Dr. Bruno Cutuli said at a breast cancer symposium sponsored by the Cancer Therapy and Research Center.

Nor is lobular carcinoma in situ (LCIS) simply a risk factor for invasive breast cancer, as has been the traditional view. In

fact, some LCIS lesions are precursors of invasive breast cancer—and as such, may warrant more aggressive local therapy, such as whole-breast radiotherapy following surgical excision, said Dr. Cutuli, a radiation oncologist at the Courlancy Polyclinic, Reims, France.

LCIS accounts for 1%-2% of all breast cancers and 15%-20% of all in situ carcinomas. The age-specific incidence reportedly jumped 2.6-fold between 1980 and 2001, for two likely reasons.

One is increased mammographic screening. Although LCIS isn't usually visible on mammography, it is often discovered on biopsy of a cyst, fibroadenoma, or other mammographically detected abnormality.

The other factor involved in the steep rise in LCIS since 1980 was probably the widespread use of hormone therapy, according to Dr. Cutuli.

Dr. Cutuli presented a retrospective study of 330 cases of pure LCIS in 325 patients treated during 1985-2000 at 12 French can-

cer centers. The patients' median age was 49 years, and 35% were postmenopausal. A first- or second-degree family history of breast cancer was present in 29%. In 24% there was prior benign breast disease.

Local recurrence rates following various therapies were illuminating. Lumpectomy was undergone by 77%; these patients had a 19.2% ipsilateral local recurrence rate during a median follow-up of 8.2 years. Of particular concern was that 60% of these local recurrences were invasive.

## Antidepressants Ease Monthly Dysphoria

SAN FRANCISCO — Women with premenstrual dysphoric disorder usually respond rapidly to the first treatment cycle of an antidepressant, allowing intermittent dosing that follows the menstrual cycle, Dr. Andrea J. Singer said at Perspectives in Women's Health sponsored by OB.GYN. NEWS.

Three selective serotonin reuptake inhibitors (SSRIs) are approved for the treatment of premenstrual dysphoric disorder (PMDD)—fluoxetine, sertraline, and paroxetine. "You don't necessarily have to have people on these long term to see a benefit," said Dr. Singer, director of women's primary care at Georgetown University Medical Center, Washington.

Dr. Singer is on the speakers' bureau of Pfizer Inc., which makes sertraline. OB.GYN. NEWS is published by the International Medical News Group, a division of Elsevier.

PMDD causes severe premenstrual symptoms that result in significant impairment of normal function, usually during the last 6-7 days of the menstrual cycle. American women with PMDD experience the symptoms on average for 8 years during their reproductive lives, she said.

Because the disorder is intermittent and treatment with an SSRI brings rapid onset of improvement, intermittent therapy is sufficient, which lowers medication costs and limits side effects, compared with treatment of overt depression. In addition, low doses typically are effective for PMDD.

Diagnosis of PMDD requires the exclusion of underlying overt depression and the presence of at least five symptoms, including at least one of four core symptoms: anger/irritability; depressed mood; moodiness; and anxiety/edginess/nervousness. Other symptoms include fatigue or lethargy, decreased interest in usual activities, insomnia or hypersomnia, difficulty concentrating, food cravings or appetite changes, feeling overwhelmed, headache, breast tenderness, bloating, and joint or muscle pain.

For women with PMDD on oral contraceptives, a pill containing drospirenone and ethinylestradiol improved PMDD symptoms in several studies. Other data suggest shortening the hormone-free interval during oral contraceptive regimens to 3-4 days can improve some symptoms.

—Sherry Boschert

**FIRST**  
IN A NOVEL  
CLASS OF  
**SLEEP**  
AGENTS



**Rozerem is indicated for the treatment of insomnia characterized by difficulty with sleep onset. Rozerem can be prescribed for long-term use. Rozerem should not be used in patients with hypersensitivity to any components of the formulation, severe hepatic impairment, or in combination with fluvoxamine. Failure of insomnia to remit after a reasonable period of time should be medically evaluated, as this may be the result of an unrecognized underlying medical disorder. Hypnotics should be administered with caution to patients exhibiting signs and symptoms of depression. Rozerem has not been studied in patients with severe sleep apnea, severe COPD, or in children or adolescents. The effects in these populations are unknown. Avoid taking Rozerem with alcohol. Rozerem has been associated with decreased testosterone levels and increased prolactin levels. Health professionals should be mindful of any unexplained symptoms possibly associated with such changes in these hormone levels. Rozerem should not be taken with or immediately after a high-fat meal. Rozerem should be taken within 30 minutes before going to bed and activities confined to preparing for bed. The most common adverse events seen with Rozerem that had at least a 2% incidence difference from placebo were somnolence, dizziness, and fatigue.**

**Please see adjacent Brief Summary of Prescribing Information.**

This experience translated to a 10-year ipsilateral local recurrence rate of 24% following conservative surgery, with a median 4.4-year delay between surgery and diagnosis of recurrent disease.

Among the 11% of patients with LCIS treated via conservative surgery plus radiotherapy, the ipsilateral local recurrence rate was 5.4% during a median 12.4 years of follow-up. Mastectomy-treated patients had no ipsilateral local recurrences. The contralateral breast cancer rate was 7.8% in patients treated with breast-conserving surgery, 5% in those who got surgery plus radiation, and 10% following mastectomy.


This was not a randomized trial. At pre-

sent no well-defined criteria exist as to which LCIS subtypes are more aggressive and thus likely to benefit from radiotherapy. But the current thinking is that the subtypes most prone to local recurrence include extensive LCIS with more than 10 involved lobules, LCIS with necrosis, pleomorphic tumors, and lesions containing cells with large nuclei, Dr. Cutuli said.

He added that his own favorable experience with breast-conserving surgery and whole-breast radiotherapy in 25 LCIS patients—there was only a single local recurrence in 12 years of follow-up—has convinced him this is a therapeutic approach worth testing in randomized trials. ■

DATA WATCH

Percentage of Women Aged ≥40 Years Having Regular Mammograms Is Decreasing



Note: Based on data from the Behavioral Risk Factor Surveillance System, 2000-2004. Source: Centers for Disease Control and Prevention

ELSEVIER GLOBAL MEDICAL NEWS

# Start and stay with nonscheduled Rozerem—ZERO evidence of abuse or dependence

Clinical studies show no evidence of potential abuse, dependence, or withdrawal\*

- **First and only**—nonscheduled prescription insomnia medication... not a controlled substance and approved for long-term use<sup>1</sup>
- **First and only**—prescription insomnia medication that targets the normal sleep-wake cycle<sup>1</sup>
- **First and only**—prescription insomnia medication with no evidence of abuse potential in clinical studies<sup>1</sup>
- **First and only**—prescription insomnia medication that does not promote sleep by CNS depression<sup>1</sup>
- **Promote sleep with Rozerem**—patients who took Rozerem fell asleep faster than those who took placebo<sup>1</sup>
- **One simple 8-mg dose**<sup>1</sup>

\*Rozerem is not a controlled substance. A clinical abuse liability study showed no differences indicative of abuse potential between Rozerem and placebo at doses up to 20 times the recommended dose (N=14). Three 35-day insomnia studies showed no evidence of rebound insomnia or withdrawal symptoms with Rozerem compared to placebo (N=2082).<sup>1,2</sup>

Please visit [www.rozerem.com](http://www.rozerem.com)

**Rozerem**<sup>TM</sup>  
**ramelteon** 8-mg tablets

*Proven for sleep.  
 Nonscheduled for added safety.*