

CPAP Alternatives Gaining Steam for Sleep Apnea

Suboptimal CPAP compliance is spurring greater use of oral appliances, maxillofacial surgery.

BY BRUCE JANCIN

FROM THE ANNUAL MEETING OF THE ASSOCIATED PROFESSIONAL SLEEP SOCIETIES

SAN ANTONIO — The days when continuous positive airway pressure was the only arrow in the quiver for physicians targeting obstructive sleep apnea are long gone.

The single most popular session at the meeting—the one whose overflow crowds brought out the fire marshals in full force—was devoted to alternatives to CPAP that have come of age: oral appliances; maxillofacial surgery; and weight loss through diet and exercise, bariatric surgery, or drugs.

Session chair James K. Walsh, Ph.D., set the stage, citing studies showing that typically 50% of patients discontinue CPAP within 1 year. Moreover, the percentage of nights patients use their CPAP drops after a couple of months from 50% to 40% and even 30%. An average of about 3 hours of use per night is the norm in clinical practice.



devices that are adjustable in all planes in space, he stressed.

“The titration of an oral appliance is essential, and it takes weeks to months,” Dr. Lowe said. “You don’t just send patients home with a ‘boil and bite’ device and say, ‘Okay, off you go.’ You need to go through the titration phase. So physicians who are prescribing oral appliances and just giving them to their patients might as well give them CPAP with a pressure of 7 mm Hg and send them home and tell them to wear it. It’s absolutely useless to do that.”

Oral appliances that have been subjected to formal trials typically show roughly an 80% success rate in patients with a baseline apnea-hypopnea index (AHI) below 30 episodes per hour, with the success rate dropping off to 60% in those with more severe OSA. Responders

‘I’m not at all trying to suggest [CPAP] therapy is totally ineffective, but I would term it highly suboptimal.’

DR. WALSH

experience less daytime sleepiness, improved cognition, better results on simulated driving performance tests, and reductions in nighttime blood pressure and serum lipids.

When Dr. Lowe and coinvestigators gave patients who were adherent to CPAP a trial period on an oral appliance, 55% subsequently switched over, while 30% maintained a clear preference for CPAP.

“Oxygenation improvement is always greater with CPAP because it forces air into the lungs. Oral appliances simply make the tube bigger and take away the obstruction,” he explained.

Device titration needs to be done by a skilled dentist. The American Academy of Dental Sleep Medicine (www.aadsm.org/index.aspx) has approximately 1,600 dentist members and is a useful resource for physicians who are seeking a local dentist experienced with oral appliances.

The main side effect associated with oral appliances is that they cause subtle tooth movement. In a series of 70 patients with full polysomnograms and dental records, Dr. Lowe found that only 10 had no change in dentition over time. Of the other 60 patients, 29 had favorable changes in the fit and function of their teeth, whereas 31 had unfavorable changes.

“The issue is how we manage it. I have yet to stop a patient from wearing an oral appliance because of tooth movement that we couldn’t manage somehow. It’s not an issue of having to cease wear. When we weigh tooth movement

against adequate oxygen to the heart, tooth movement loses. I’m trying to train the profession to think that way—panic less about tooth movement and think more about what the treatment is doing for the sleep-disordered breathing,” Dr. Lowe continued.

Besides, his 3-year study of patients using classic CPAP masks showed that they, too, cause quantifiable changes in tooth position over time, he added.

Maxillofacial Surgery

Maxillomandibular advancement is a big operation, and it yields big results, said Dr. Kasey Li of Stanford (Calif.) University.

Dr. Li cited a recent meta-analysis that included a total of 627 patients who underwent maxillomandibular advancement (MMA).

Their mean AHI dropped from 63.9 to 9.5 events/hour. Treatment success, defined as an AHI below 20, occurred in 86% of patients. A surgical cure, meaning an AHI below 5, was obtained in 43% of patients (*Sleep Med. Rev.* 2010 [doi: 10.1016/j.smrv.2009.11.003]).

This parallels Dr. Li’s personal experience, which includes 302 patients with pre- and post-MMA sleep data. The operation typically takes about 3 hours, with a 2- to 3-day hospital stay and return to work in 4-5 weeks.

As in the meta-analysis, there have been no postoperative deaths in Dr. Li’s own series. The most common side effect is cranial nerve paresthesia, which typically resolves within 6-12 months. Four of Dr. Li’s patients had severe malocclusion requiring revision surgery.

DR. LOWE

Ninety percent of patients report being satisfied with their results.

A multivariate regression analysis done as part of the meta-analysis identified four significant predictors of increased likelihood of MMA success: younger age, lower body mass index, less severe sleep apnea, and greater degree of maxillary advancement. This mirrors Dr. Li’s experience as well.

“My enthusiasm for offering surgery to patients over age 60 goes down quite a bit. I’m fairly reluctant to offer surgery to patients with a BMI of 32-33 kg/m² or above, and you have to be able to advance the maxilla at least 10 mm in order to advance the airway. And the treatment success rate is much higher in patients with an AHI in the 20s-40s than the 80s,” the surgeon said.

Patients who have failed on oral appliances remain reasonable candidates for surgery.

“The average advancement with an

oral appliance is 4-7 mm, in comparison to 15-16 mm of maxillary advancement measured at the teeth level with surgery, in my experience,” Dr. Li noted.

When asked if it makes sense to perform a less morbid soft tissue surgical procedure such as tonsillectomy or uvulopalatopharyngoplasty as a first-line operation for patients with moderate to severe OSA, reserving MMA for the nonresponders, Dr. Li’s answer was emphatically no.

“In the majority of patients, the entire airway is involved; the obstruction is at multiple levels. Surgical procedures that focus on one site often will not be successful,” he said.

The mean apnea-hypopnea index dropped from 63.9 to 9.5 events/hour after maxillomandibular advancement.

DR. LI

“The data over the past 10 years are very clear that patients with severe sleep apnea are not going to respond very well to soft tissue surgery, period. I tell patients that unless they’re going to have MMA, they shouldn’t bother with surgery. That’s my personal bias, and I think it’s supported by the data,” he added.

A prospective study that will compare MMA with CPAP is in the planning stages at Stanford.

Weight Loss

Too many physicians are jaded about this well-established but seriously underused treatment for OSA, according to Dr. Ronald R. Grunstein, professor of sleep medicine at the University of Sydney.

“I think we need to have a less nihilistic view about weight loss. We in sleep medicine are often still thinking very much in silos,” he said.

Dr. Grunstein was first author of a large study with a 2-year follow-up period that demonstrated bariatric surgery to be a highly effective treatment for OSA in obese patients (*Sleep* 2007;30:703-10).

In addition, recent studies conducted in Finland (*Am. J. Respir. Crit. Care Med.* 2009;179:320-7) and Sweden (*BMJ* 2009;339:b4609) have shown substantial improvement in OSA with weight loss achieved through a very-low-calorie diet plus exercise followed by a maintenance diet.

The bigger the weight loss, the greater the improvement in OSA as reflected in the reduction in AHI.

Promising pharmacologic alternatives to CPAP are also in development, and not all are weight-loss drugs. ■

Disclosures: Dr. Lowe is the inventor of the Klearway oral appliance, the royalties for which are assigned to the University of British Columbia, where they pay for much of his research. Dr. Walsh is a consultant to Ventus Medical Inc., which markets the Provent sleep apnea therapy device, which uses nasal expiratory positive airway pressure. Dr. Li and Dr. Grunstein reported no financial conflicts.