

Mass. Coalition Launches Health Records Project

BY MARY ELLEN SCHNEIDER
Senior Writer

BOSTON — Three Massachusetts communities will soon be wired for electronic health record systems as part of a \$50 million pilot project.

The idea, which is being undertaken by the Massachusetts eHealth Collaborative, is to test out the implementation of interoperable EHRs within communities before attempting to connect physicians

across the entire state.

“We’re completely focused on practical solutions so we can get these things into physicians’ hands and health care professionals’ hands and keep them there,” Micky Tripathi, CEO of the Massachusetts eHealth Collaborative said at a congress sponsored by the American Medical Informatics Association.

The collaborative is a not-for-profit group that was founded by 34 health care institutions seeking to create a health in-

formation network that would connect providers statewide.

The collaborative was launched last fall and requested applications for its pilot project last December. It received a total of 35 applications from communities located across the state and chose three communities—greater Brockton, greater Newburyport, and Northern Berkshire. The pilot project is being funded through a grant from Blue Cross Blue Shield of Massachusetts.

Each community chosen for the pilot project was a relatively self-contained medical referral market, had strong local health care professional leadership, and demonstrated an openness to information technology innovation, Mr. Tripathi said.

The final selection of the three communities was based in part on location, patient diversity, and information technology maturity, he said. Members of the collaborative also wanted to choose communities that were at different points of the information technology adoption curve in order to see the different types of benefits.

The three communities cover a total of nearly 600 physicians treating approximately 500,000 patients. Overall, there are 182 primary care physicians and 410 specialists. The pilots will include almost 200 office sites, most of which have between one and five physicians, Mr. Tripathi said.

The pilot projects will include the purchase and installation of EHRs at all clinical care points, as well as connecting them to other systems within the community.

Although existing studies have shown the benefits that are conferred by the use of EHRs on a small scale, Mr. Tripathi said the pilot project is a chance to see what will happen in a larger, community-wide rollout.

The pilot will be aimed at determining the barriers to adoption, identifying the costs—both direct and indirect—of adoption, and analyzing the benefits. Officials at the collaborative will also be seeking to figure out how the costs and benefits are distributed across the various stakeholders.

Finally, they will be looking for the best ways to provide incentives and how that could be replicated going forward.

“This transition can’t be done to physicians,” he said. “It’s got to be an idea that we sell to them.”

The Massachusetts eHealth Collaborative plans to select EHR vendors by the end of May and be under contract by the end of the summer.

The pilot timeline calls for implementing systems in a clinical care setting before the end of the year. At the beginning of 2006, the collaborative expects to implement interoperability capabilities for the systems.

The pilot projects are slated to end in mid-2008.

In addition, the collaborative plans to work with the other 32 communities that were not selected for the pilot to help them implement EHR systems by sharing the infrastructure, expertise, and arrangements created through the pilots.

ENABLEX® (darifenacin)

Extended-release tablets

Rx only
BRIEF SUMMARY: Please see package insert for full prescribing information.

INDICATIONS AND USAGE
ENABLEX® (darifenacin) extended-release tablets are indicated for the treatment of overactive bladder with symptoms of urge urinary incontinence, urgency and frequency.

CONTRAINDICATIONS
ENABLEX® (darifenacin) extended-release tablets are contraindicated in patients with urinary retention, gastric retention or uncontrolled narrow-angle glaucoma and in patients who are at risk for these conditions. ENABLEX is also contraindicated in patients with known hypersensitivity to the drug or its ingredients.

PRECAUTIONS
General
Risk of Urinary Retention
ENABLEX® (darifenacin) extended-release tablets should be administered with caution to patients with clinically significant bladder outflow obstruction because of the risk of urinary retention.

Decreased Gastrointestinal Motility
ENABLEX should be administered with caution to patients with gastrointestinal obstructive disorders because of the risk of gastric retention. ENABLEX, like other anticholinergic drugs, may decrease gastrointestinal motility and should be used with caution in patients with conditions such as severe constipation, ulcerative colitis, and myasthenia gravis.

Controlled Narrow-Angle Glaucoma
ENABLEX should be used with caution in patients being treated for narrow-angle glaucoma and only where the potential benefits outweigh the risks.

Patients with Hepatic Impairment
There are no dosing adjustments for patients with mild hepatic impairment. The daily dose of ENABLEX should not exceed 7.5 mg for patients with moderate hepatic impairment. ENABLEX has not been studied in patients with severe hepatic impairment and therefore is not recommended for use in this patient population (see CLINICAL PHARMACOLOGY, Pharmacokinetics in Special Populations and DOSAGE AND ADMINISTRATION in the full prescribing information).

Information for Patients
Patients should be informed that anticholinergic agents, such as ENABLEX, may produce clinically significant adverse effects related to anticholinergic pharmacological activity including constipation, urinary retention and blurred vision. Heat prostration (due to decreased sweating) can occur when anticholinergics such as ENABLEX are used in a hot environment. Because anticholinergics, such as ENABLEX, may produce dizziness or blurred vision, patients should be advised to exercise caution in decisions to engage in potentially dangerous activities until the drug's effects have been determined. Patients should read the patient information leaflet before starting therapy with ENABLEX.

ENABLEX extended-release tablets should be taken once daily with liquid. They may be taken with or without food, and should be swallowed whole and not chewed, divided or crushed.

Drug Interactions
The daily dose of ENABLEX should not exceed 7.5 mg when coadministered with potent CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, ritonavir, nelfinavir, clarithromycin and nefazodone) (see CLINICAL PHARMACOLOGY and DOSAGE AND ADMINISTRATION in the full prescribing information).

Caution should be taken when ENABLEX is used concomitantly with medications that are predominantly metabolized by CYP2D6 and which have a narrow therapeutic window, such as flecainide, thioridazine and tricyclic antidepressants (see CLINICAL PHARMACOLOGY in the full prescribing information).

The concomitant use of ENABLEX with other anticholinergic agents may increase the frequency and/or severity of dry mouth, constipation, blurred vision and other anticholinergic pharmacological effects. Anticholinergic agents may potentially alter the absorption of some concomitantly administered drugs due to effects on gastrointestinal motility.

Drug Laboratory Test Interactions
Interactions between darifenacin and laboratory tests have not been studied.

Carcinogenesis/Mutagenesis/Impairment of Fertility
Carcinogenicity studies with darifenacin were conducted in mice and rats. No evidence of drug-related carcinogenicity was revealed in a 24-month study in mice at dietary doses up to 100 mg/kg/day or approximately 32 times the estimated human-free AUC₀₋₂₄ reached with 15 mg, the maximum recommended human dose (AUC at MRHD) and in a 24-month study in rats at doses up to 15 mg/kg/day or up to approximately 12 times the AUC at MRHD in female rats and approximately eight times the AUC at MRHD in male rats.

Darifenacin was not mutagenic in the bacterial mutation assays (Ames test) and the Chinese hamster ovary assay, and not clastogenic in the human lymphocyte assay, and the *in vivo* mouse bone marrow cytogenetics assay.

There was no evidence for effects on fertility in male or female rats treated at oral doses up to 50 mg/kg/day. Exposures in this study correspond to approximately 78 times the AUC at MRHD.

Pregnancy Category C
Darifenacin was not teratogenic in rats and rabbits at doses up to 50 and 30 mg/kg/day, respectively. At the dose of 50 mg/kg in rats, there was a delay in the ossification of the sacral and caudal vertebrae which was not observed at 10 mg/kg (approximately 13 times the AUC of free plasma concentration at MRHD). Exposure in this study at 50 mg/kg corresponds to approximately 59 times the AUC of free plasma concentration at MRHD. Dystocia was observed in dams at 10 mg/kg/day (17 times the AUC of free plasma concentration at MRHD). Slight developmental delays were observed in pups at this dose. At 3 mg/kg/day (five times the AUC of free plasma concentration at MRHD) there were no effects on dams or pups. At the dose of 30 mg/kg in rabbits, darifenacin was shown to increase post-implantation loss but not at 10 mg/kg (nine times the AUC of free plasma concentration at MRHD). Exposure to unbound drug at 30 mg/kg in this study corresponds to approximately 28 times the AUC at MRHD. In rabbits, dilated ureter and/or kidney pelvis was observed in offspring at 30 mg/kg/day and one case was observed at 10 mg/kg/day along with urinary bladder dilation consistent with pharmacological action of darifenacin. No effect was observed at 3 mg/kg/day (2.8 times the AUC of free plasma concentration at MRHD). There are no studies of darifenacin in pregnant women. Because animal reproduction studies are not always predictive of human response, ENABLEX should be used during pregnancy only if the benefit to the mother outweighs the potential risk to the fetus.

Nursing Mothers
Darifenacin is excreted into the milk of rats. It is not known whether darifenacin is excreted into human milk and therefore caution should be exercised before ENABLEX is administered to a nursing woman.

Pediatric Use
The safety and effectiveness of ENABLEX in pediatric patients have not been established.

Geriatric Use
In the Phase III fixed-dose, placebo-controlled, clinical studies, 30% of patients treated with ENABLEX were over 65 years of age. No overall differences in safety or efficacy were observed between these patients (n=207) and younger patients <65 years (n=464). No dose adjustment is recommended for elderly patients (see CLINICAL PHARMACOLOGY, Pharmacokinetics in Special Populations and CLINICAL STUDIES in the full prescribing information).

References: **1.** Chapple CR, Yamanishi T, Chess-Williams R. Muscarinic receptor subtypes and management of the overactive bladder. *Urology*. 2002;60(suppl 5A):82-89. **2.** Federal Interagency Forum on Aging-Related Statistics. Older Americans 2000: key indicators of well-being. August 2000. Available at: <http://www.agingstats.gov/chartbook2000/healthstatus.html#Indicator%2014>. Accessed November 4, 2004. **3.** Feinberg M. The problems of anticholinergic adverse effects in older patients. *Drugs Aging*. 1993;3:335-348.

ADVERSE REACTIONS
During the clinical development of ENABLEX® (darifenacin) extended-release tablets, a total of 7,363 patients and volunteers were treated with doses of darifenacin from 3.75 mg to 75 mg once daily.

The safety of ENABLEX was evaluated in Phase II and III controlled clinical trials in a total of 8,830 patients, 6,001 of whom were treated with ENABLEX. Of this total, 1,069 patients participated in three, 12-week, Phase III, fixed-dose efficacy and safety studies. Of this total, 337 and 334 patients received ENABLEX 7.5 mg daily and 15 mg daily, respectively. In all long-term trials combined, 1,216 and 672 patients received treatment with ENABLEX for at least 24 and 52 weeks, respectively.

In all placebo-controlled trials combined, the incidence of serious adverse events for 7.5 mg, 15 mg and placebo was similar.

In all fixed-dose Phase III studies combined, 3.3% of patients treated with ENABLEX discontinued due to all adverse events versus 2.6% in placebo. Dry mouth leading to study discontinuation occurred in 0%, 0.9%, and 0% of patients treated with ENABLEX 7.5 mg daily, ENABLEX 15 mg daily and placebo, respectively. Constipation leading to study discontinuation occurred in 0.6%, 1.2%, and 0.3% of patients treated with ENABLEX 7.5 mg daily, ENABLEX 15 mg daily and placebo, respectively.

Table 4 lists the adverse events reported (regardless of causality) in 2% or more of patients treated with 7.5-mg or 15-mg ENABLEX extended-release tablets and greater than placebo in the three, fixed-dose, placebo-controlled Phase III studies (Studies 1, 2 and 3). Adverse events were reported by 54% and 66% of patients receiving 7.5 mg and 15 mg once-daily ENABLEX extended-release tablets, respectively, and by 49% of patients receiving placebo. In these studies, the most frequently reported adverse events were dry mouth and constipation. The majority of adverse events in ENABLEX-treated subjects were mild or moderate in severity and most occurred during the first two weeks of treatment.

Table 4 Incidence of Adverse Events* Reported in 2.0% of Patients Treated with ENABLEX® Extended-Release Tablets and More Frequent with ENABLEX® than with Placebo in Three, Fixed-Dose, Placebo-Controlled, Phase III Studies (Studies 1, 2, and 3)				
Body System	Adverse Event	Percentage of Subjects with Adverse Event (%)		
		ENABLEX® 7.5 mg N = 337	ENABLEX® 15 mg N = 334	Placebo N = 388
Digestive	Dry Mouth	20.2	35.3	8.2
	Constipation	14.8	21.3	6.2
	Dyspepsia	2.7	8.4	2.6
	Abdominal Pain	2.4	3.9	0.5
	Nausea	2.7	1.5	1.5
	Diarrhea	2.1	0.9	1.8
	Urinary Tract Infection	4.9	4.5	2.6
Urogenital	Dizziness	1.5	2.1	1.3
Nervous	Headache	1.5	2.7	1.3
Body as a Whole	Dry Eyes	1.5	2.1	0.5
Eye				

*Regardless of causality

Other adverse events reported, regardless of causality, by ≥1% of ENABLEX patients in either the 7.5 mg or 15 mg once-daily darifenacin-dose groups in these fixed-dose, placebo-controlled Phase III studies include: abnormal vision, accidental injury, back pain, dry skin, flu syndrome, pain, hypertension, vomiting, peripheral edema, weight gain, arthralgia, bronchitis, pharyngitis, rhinitis, sinusitis, rash, pruritus, urinary tract disorder and vaginitis.

Study 4 was a 12-week, placebo-controlled, dose-titration regimen study in which ENABLEX was administered in accordance with dosing recommendations (see DOSAGE AND ADMINISTRATION in the full prescribing information). All patients initially received placebo or ENABLEX 7.5 mg daily, and after two weeks, patients and physicians were allowed to adjust upward to ENABLEX 15 mg if needed. In this study, the most commonly reported adverse events were also constipation and dry mouth. The incidence of discontinuation due to all adverse events was 3.1% and 6.7% for placebo and for ENABLEX, respectively. Table 5 lists the adverse events (regardless of causality) reported in >3% of patients treated with ENABLEX extended-release tablets and greater than placebo.

Table 5 Number (%) of Adverse Events* Reported in >3% of Patients Treated with ENABLEX® Extended-Release Tablets, and More Frequent with ENABLEX® than Placebo, in the Placebo-Controlled, Dose-Titration, Phase III Study (Study 4)		
Adverse Event	ENABLEX® 7.5 mg/15 mg N = 268	Placebo N = 127
Constipation	56 (20.9%)	10 (7.9%)
Dry Mouth	50 (18.7%)	11 (8.7%)
Headache	18 (6.7%)	7 (5.5%)
Dyspepsia	12 (4.5%)	2 (1.6%)
Nausea	11 (4.1%)	2 (1.6%)
Urinary Tract Infection	10 (3.7%)	4 (3.1%)
Accidental Injury	8 (3.0%)	3 (2.4%)
Flu Syndrome	8 (3.0%)	3 (2.4%)

*Regardless of causality

Acute urinary retention (AUR) requiring treatment was reported in a total of 16 patients in the ENABLEX Phase I-III clinical trials. Of these 16 cases, seven were reported as serious adverse events, including one patient with detrusor hyperreflexia secondary to a stroke, one patient with benign prostatic hypertrophy (BPH), one patient with irritable bowel syndrome (IBS) and four OAB patients taking darifenacin 30 mg daily. Of the remaining nine cases, none were reported as serious adverse events. Three occurred in OAB patients taking the recommended doses, and two of these required bladder catheterization for 1-2 days.

Constipation was reported as a serious adverse event in six patients in the ENABLEX Phase I-III clinical trials, including one patient with benign prostatic hypertrophy (BPH), one OAB patient taking darifenacin 30 mg daily, and only one OAB patient taking the recommended doses. The latter patient was hospitalized for investigation with colonoscopy after reporting nine months of chronic constipation that was reported as being moderate in severity.

Storage
Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature]. Protect from light.

T2004-49
5000213
5000214
5000215

DECEMBER 2004
Manufactured by:
Pfizer Inc.
Brooklyn, New York 11206

Printed in U.S.A.

Distributed by:
Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936
©Novartis



Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936-1080

©2005 Novartis

3/05

C-ENA-1012