

ACIP Meningococcal Working Group Suggests Waiting on Infant Vaccination

BY HEIDI SPLETE

ATLANTA — The meningococcal working group of the Centers for Disease Control and Prevention’s Advisory Committee on Immunization Practices “believes that the ACIP should consider not adding meningococcal conjugate vaccines to the routine infant vaccine

schedule at this time,” said working group member Dr. Amanda Cohn. At its fall meeting, the ACIP discussed safety and epidemiology data on meningococcal vaccines in development for infants. These products have not yet been licensed. The low burden of meningococcal disease in infants raises the question of

whether every vaccine that is shown to be safe and effective should be recommended if the burden of disease is low, said Dr. H. Cody Meissner, chair of the working group. The last ACIP recommendations for meningococcal vaccines were published in May 2005, and an update is planned for 2010, he noted. ■

Second HPV Vaccine Gets Green Light

BY HEIDI SPLETE

ATLANTA — The Centers for Disease Control and Prevention’s Advisory Committee on Immunization Practices recommended a bivalent human papillomavirus vaccine as an alternative to the quadrivalent vaccine for the prevention of cervical cancer and related precancerous conditions in women and girls aged 9-26 years.

ACIP made the recommendation at its annual fall meeting. The bivalent human papillomavirus (HPV) vaccine (GlaxoSmithKline’s Cervarix) was recently approved by the Food and Drug Administration. The vaccine provides clinicians with another option to vaccinate adolescent girls and young women against diseases caused by HPV types 16 and 18. But unlike the quadrivalent vaccine, the bivalent vaccine is not designed to protect against genital warts, noted Dr. Lauri Markowitz of the CDC, who presented the ACIP recommendations.

The quadrivalent vaccine (Merck & Co.’s Gardasil) protects against genital warts associated with HPV types 6 and 11, in addition to protecting against diseases caused by HPV types 16 and 18. ACIP recommended against a statement of no preference between the bivalent and quadrivalent vaccines after a lively debate. Instead, the recommendations will present the information about the two vaccines without a statement of preference or a statement of nonpreference.

The recommendations state that the two vaccines can be used interchangeably to complete the three-dose series, but that using the same vaccine for the entire series is preferable. The bivalent vaccine, like the quadrivalent vaccine, is not a live vaccine, and it can be given simultaneously with other vaccines.

ACIP also voted to harmonize the age ranges for the two vaccines, with first doses given at ages 11-12 years and recommended second and third doses at 1-2 months and 6 months after the first dose. The recommended minimum dosing intervals remained as 4 weeks between the first and second dose and 12 weeks between the second and third doses. The vaccine can be initiated as young as 9 years, and catch-up vaccination is recommended for females aged 13-26 years.

In addition, ACIP voted to move information about pregnancy to the precautions section. Pregnancy is not currently a contraindication for the vaccine. But Dr. Frank DeStefano of the CDC’s immunization safety office said that post-marketing safety surveillance studies would be conducted on the bivalent HPV vaccine using the Vaccine Adverse Event Reporting System (VAERS). The committee voted to add the bivalent vaccine to the CDC’s Vaccines for Children program. ■

ONGLYZA™ (saxagliptin) tablets **Rx ONLY**
Brief Summary of Prescribing Information. For complete prescribing information consult official package insert.

INDICATIONS AND USAGE

Monotherapy and Combination Therapy

ONGLYZA (saxagliptin) is indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus. [See *Clinical Studies* (14).]

Important Limitations of Use

ONGLYZA should not be used for the treatment of type 1 diabetes mellitus or diabetic ketoacidosis, as it would not be effective in these settings.

ONGLYZA has not been studied in combination with insulin.

CONTRAINDICATIONS

None.

WARNINGS AND PRECAUTIONS

Use with Medications Known to Cause Hypoglycemia

Insulin secretagogues, such as sulfonylureas, cause hypoglycemia. Therefore, a lower dose of the insulin secretagogue may be required to reduce the risk of hypoglycemia when used in combination with ONGLYZA. [See *Adverse Reactions* (6.1).]

Macrovascular Outcomes

There have been no clinical studies establishing conclusive evidence of macrovascular risk reduction with ONGLYZA or any other antidiabetic drug.

ADVERSE REACTIONS

Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Monotherapy and Add-On Combination Therapy

In two placebo-controlled monotherapy trials of 24-weeks duration, patients were treated with ONGLYZA 2.5 mg daily, ONGLYZA 5 mg daily, and placebo. Three 24-week, placebo-controlled, add-on combination therapy trials were also conducted: one with metformin, one with a thiazolidinedione (pioglitazone or rosiglitazone), and one with glyburide. In these three trials, patients were randomized to add-on therapy with ONGLYZA 2.5 mg daily, ONGLYZA 5 mg daily, or placebo. A saxagliptin 10 mg treatment arm was included in one of the monotherapy trials and in the add-on combination trial with metformin.

In a prespecified pooled analysis of the 24-week data (regardless of glycemic rescue) from the two monotherapy trials, the add-on to metformin trial, the add-on to thiazolidinedione (TZD) trial, and the add-on to glyburide trial, the overall incidence of adverse events in patients treated with ONGLYZA 2.5 mg and ONGLYZA 5 mg was similar to placebo (72.0% and 72.2% versus 70.6%, respectively). Discontinuation of therapy due to adverse events occurred in 2.2%, 3.3%, and 1.8% of patients receiving ONGLYZA 2.5 mg, ONGLYZA 5 mg, and placebo, respectively. The most common adverse events (reported in at least 2 patients treated with ONGLYZA 2.5 mg or at least 2 patients treated with ONGLYZA 5 mg) associated with premature discontinuation of therapy included lymphopenia (0.1% and 0.5% versus 0%, respectively), rash (0.2% and 0.3% versus 0.3%), blood creatinine increased (0.3% and 0% versus 0%), and blood creatine phosphokinase increased (0.1% and 0.2% versus 0%). The adverse reactions in this pooled analysis reported (regardless of investigator assessment of causality) in ≥5% of patients treated with ONGLYZA 5 mg, and more commonly than in patients treated with placebo are shown in Table 1.

Table 1: Adverse Reactions (Regardless of Investigator Assessment of Causality) in Placebo-Controlled Trials* Reported in ≥5% of Patients Treated with ONGLYZA 5 mg and More Commonly than in Patients Treated with Placebo

	Number (%) of Patients	
	ONGLYZA 5 mg N=882	Placebo N=799
Upper respiratory tract infection	68 (7.7)	61 (7.6)
Urinary tract infection	60 (6.8)	49 (6.1)
Headache	57 (6.5)	47 (5.9)

* The 5 placebo-controlled trials include two monotherapy trials and one add-on combination therapy trial with each of the following: metformin, thiazolidinedione, or glyburide. Table shows 24-week data regardless of glycemic rescue.

In patients treated with ONGLYZA 2.5 mg, headache (6.5%) was the only adverse reaction reported at a rate ≥5% and more commonly than in patients treated with placebo.

In this pooled analysis, adverse reactions that were reported in ≥2% of patients treated with ONGLYZA 2.5 mg or ONGLYZA 5 mg and ≥1% more frequently compared to placebo included: sinusitis (2.9% and 2.6% versus 1.6%, respectively), abdominal pain (2.4% and 1.7% versus 0.5%), gastroenteritis (1.9% and 2.3% versus 0.9%), and vomiting (2.2% and 2.3% versus 1.3%).

In the add-on to TZD trial, the incidence of peripheral edema was higher for ONGLYZA 5 mg versus placebo (8.1% and 4.3%, respectively). The incidence of peripheral edema for ONGLYZA 2.5 mg was 3.1%. None of the reported adverse reactions of peripheral edema resulted in study drug discontinuation. Rates of peripheral edema for ONGLYZA 2.5 mg and ONGLYZA 5 mg versus placebo were 3.6% and 2% versus 3% given as monotherapy, 2.1% and 2.1% versus 2.2% given as add-on therapy to metformin, and 2.4% and 1.2% versus 2.2% given as add-on therapy to glyburide.

The incidence rate of fractures was 1.0 and 0.6 per 100 patient-years, respectively, for ONGLYZA (pooled analysis of 2.5 mg, 5 mg, and 10 mg) and placebo. The incidence rate of fracture events in patients who received ONGLYZA did not increase over time. Causality has not been established and nonclinical studies have not demonstrated adverse effects of saxagliptin on bone.

An event of thrombocytopenia, consistent with a diagnosis of idiopathic thrombocytopenic purpura, was observed in the clinical program. The relationship of this event to ONGLYZA is not known.

Adverse Reactions Associated with ONGLYZA (saxagliptin) Coadministered with Metformin in Treatment-Naïve Patients with Type 2 Diabetes

Table 2 shows the adverse reactions reported (regardless of investigator assessment of causality) in ≥5% of patients participating in an additional 24-week, active-controlled trial of coadministered ONGLYZA and metformin in treatment-naïve patients.

Table 2: Initial Therapy with Combination of ONGLYZA and Metformin in Treatment-Naïve Patients: Adverse Reactions Reported (Regardless of Investigator Assessment of Causality) in ≥5% of Patients Treated with Combination Therapy of ONGLYZA 5 mg Plus Metformin (and More Commonly than in Patients Treated with Metformin Alone)

	Number (%) of Patients	
	ONGLYZA 5 mg + Metformin* N=320	Metformin* N=328
Headache	24 (7.5)	17 (5.2)
Nasopharyngitis	22 (6.9)	13 (4.0)

* Metformin was initiated at a starting dose of 500 mg daily and titrated up to a maximum of 2000 mg daily.

Hypoglycemia

Adverse reactions of hypoglycemia were based on all reports of hypoglycemia; a concurrent glucose measurement was not required. In the add-on to glyburide study, the overall incidence of reported hypoglycemia was higher for ONGLYZA 2.5 mg and ONGLYZA 5 mg (13.3% and 14.6%) versus placebo (10.1%). The incidence of confirmed hypoglycemia in this study, defined as symptoms of hypoglycemia accompanied by a fingerstick glucose value of ≤50 mg/dL, was 2.4% and 0.8% for ONGLYZA 2.5 mg and ONGLYZA 5 mg and 0.7% for placebo. The incidence of reported hypoglycemia for ONGLYZA 2.5 mg and ONGLYZA 5 mg versus placebo given as monotherapy was 4.0% and 5.6% versus 4.1%, respectively, 7.8% and 5.8% versus 5% given as add-on therapy to metformin, and 4.1% and 2.7% versus 3.8% given as add-on therapy to TZD. The incidence of reported hypoglycemia was 3.4% in treatment-naïve patients given ONGLYZA 5 mg plus metformin and 4.0% in patients given metformin alone.

Hypersensitivity Reactions

Hypersensitivity-related events, such as urticaria and facial edema in the 5-study pooled analysis up to Week 24 were reported in 1.5%, 1.5%, and 0.4% of patients who received ONGLYZA 2.5 mg, ONGLYZA 5 mg, and placebo, respectively. None of these events in patients who received ONGLYZA required hospitalization or were reported as life-threatening by the investigators. One saxagliptin-treated patient in this pooled analysis discontinued due to generalized urticaria and facial edema.

Vital Signs

No clinically meaningful changes in vital signs have been observed in patients treated with ONGLYZA.

Laboratory Tests

Absolute Lymphocyte Counts

There was a dose-related mean decrease in absolute lymphocyte count observed with ONGLYZA. From a baseline mean absolute lymphocyte count of approximately 2200 cells/microL, mean decreases of approximately 100 and 120 cells/microL with ONGLYZA 5 mg and 10 mg, respectively, relative to placebo were observed at 24 weeks in a pooled analysis of five placebo-controlled clinical studies. Similar effects were observed when ONGLYZA 5 mg was given in initial combination with metformin compared to metformin alone. There was no difference observed for ONGLYZA 2.5 mg relative to placebo. The proportion of patients who were reported to have a lymphocyte count ≤750 cells/microL was 0.5%, 1.5%, 1.4%, and 0.4% in the saxagliptin 2.5 mg, 5 mg, 10 mg, and placebo groups, respectively. In most patients, recurrence was not observed with repeated exposure to ONGLYZA although some patients had recurrent decreases upon rechallenge that led to discontinuation of ONGLYZA. The decreases in lymphocyte count were not associated with clinically relevant adverse reactions.

The clinical significance of this decrease in lymphocyte count relative to placebo is not known. When clinically indicated, such as in settings of unusual or prolonged infection, lymphocyte count should be measured. The effect of ONGLYZA on lymphocyte counts in patients with lymphocyte abnormalities (e.g., human immunodeficiency virus) is unknown.

Platelets

ONGLYZA did not demonstrate a clinically meaningful or consistent effect on platelet count in the six, double-blind, controlled clinical safety and efficacy trials.

DRUG INTERACTIONS

Inducers of CYP3A4/5 Enzymes

Rifampin significantly decreased saxagliptin exposure with no change in the area under the time-concentration curve (AUC) of its active metabolite, 5-hydroxy saxagliptin. The plasma dipeptidyl peptidase-4 (DPP4) activity inhibition over a 24-hour dose interval was not affected by rifampin. Therefore, dosage adjustment of ONGLYZA is not recommended. [See *Clinical Pharmacology* (12.3).]

Inhibitors of CYP3A4/5 Enzymes

Moderate Inhibitors of CYP3A4/5

Diltiazem increased the exposure of saxagliptin. Similar increases in plasma concentrations of saxagliptin are anticipated in the presence of other moderate CYP3A4/5 inhibitors (e.g., amprenavir, aprepitant, erythromycin, fluconazole, fosamprenavir, grapefruit juice, and verapamil); however, dosage adjustment of ONGLYZA is not recommended. [See *Clinical Pharmacology* (12.3).]

Strong Inhibitors of CYP3A4/5

Ketoconazole significantly increased saxagliptin exposure. Similar significant increases in plasma concentrations of saxagliptin are anticipated with other strong CYP3A4/5 inhibitors (e.g., atazanavir, clarithromycin, indinavir, itraconazole, nefazodone, nelfinavir, ritonavir, saquinavir, and telithromycin). The dose of ONGLYZA should be limited to 2.5 mg when coadministered with a strong CYP3A4/5 inhibitor. [See *Dosage and Administration* (2.3) and *Clinical Pharmacology* (12.3).]

USE IN SPECIFIC POPULATIONS

Pregnancy

Pregnancy Category B

There are no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, ONGLYZA (saxagliptin), like other antidiabetic medications, should be used during pregnancy only if clearly needed.

Saxagliptin was not teratogenic at any dose tested when administered to pregnant rats and rabbits during periods of organogenesis. Incomplete ossification of the pelvis, a form of developmental delay, occurred in rats at a dose of 240 mg/kg, or approximately 1503 and 66 times human exposure to saxagliptin and the active metabolite, respectively, at the maximum recommended human dose (MRHD) of 5 mg. Maternal toxicity and reduced fetal body weights were observed at 7986 and 328 times the human exposure at the MRHD for saxagliptin and the active metabolite, respectively. Minor skeletal variations in rabbits occurred at a maternally toxic dose of 200 mg/kg, or approximately 1432 and 992 times the MRHD. When administered to rats in combination with metformin, saxagliptin was not teratogenic nor embryolethal at exposures 21 times the saxagliptin MRHD. Combination administration of metformin with a higher dose of saxagliptin (109 times the saxagliptin MRHD) was associated with craniorachischisis (a rare neural tube defect characterized by incomplete closure of the skull and spinal column) in two fetuses from a single dam. Metformin exposures in each combination were 4 times the human exposure of 2000 mg daily.

Saxagliptin administered to female rats from gestation day 6 to lactation day 20 resulted in decreased body weights in male and female offspring only at maternally toxic doses (exposures ≥1629 and 53 times saxagliptin and its active metabolite at the MRHD). No functional or behavioral toxicity was observed in offspring of rats administered saxagliptin at any dose.

Saxagliptin crosses the placenta into the fetus following dosing in pregnant rats.

Nursing Mothers

Saxagliptin is secreted in the milk of lactating rats at approximately a 1:1 ratio with plasma drug concentrations. It is not known whether saxagliptin is secreted in human milk. Because many drugs are secreted in human milk, caution should be exercised when ONGLYZA is administered to a nursing woman.

Pediatric Use

Safety and effectiveness of ONGLYZA in pediatric patients have not been established.

Geriatric Use

In the six, double-blind, controlled clinical safety and efficacy trials of ONGLYZA, 634 (15.3%) of the 4148 randomized patients were 65 years and over, and 59 (1.4%) patients were 75 years and over. No overall differences in safety or effectiveness were observed between patients ≥65 years old and the younger patients. While this clinical experience has not identified differences in responses between the elderly and younger patients, greater sensitivity of some older individuals cannot be ruled out.

Saxagliptin and its active metabolite are eliminated in part by the kidney. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection in the elderly based on renal function. [See *Dosage and Administration* (2.2) and *Clinical Pharmacology* (12.3).]

OVERDOSAGE

In a controlled clinical trial, once-daily, orally-administered ONGLYZA in healthy subjects at doses up to 400 mg daily for 2 weeks (80 times the MRHD) had no dose-related clinical adverse reactions and no clinically meaningful effect on QTc interval or heart rate.

In the event of an overdose, appropriate supportive treatment should be initiated as dictated by the patient’s clinical status. Saxagliptin and its active metabolite are removed by hemodialysis (23% of dose over 4 hours).

PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling.

Instructions

Patients should be informed of the potential risks and benefits of ONGLYZA and of alternative modes of therapy. Patients should also be informed about the importance of adherence to dietary instructions, regular physical activity, periodic blood glucose monitoring and A1C testing, recognition and management of hypoglycemia and hyperglycemia, and assessment of diabetes complications. During periods of stress such as fever, trauma, infection, or surgery, medication requirements may change and patients should be advised to seek medical advice promptly.

Physicians should instruct their patients to read the Patient Package Insert before starting ONGLYZA therapy and to reread it each time the prescription is renewed. Patients should be instructed to inform their doctor or pharmacist if they develop any unusual symptom or if any existing symptom persists or worsens.

Laboratory Tests

Patients should be informed that response to all diabetic therapies should be monitored by periodic measurements of blood glucose and A1C, with a goal of decreasing these levels toward the normal range. A1C is especially useful for evaluating long-term glycemic control. Patients should be informed of the potential need to adjust their dose based on changes in renal function tests over time.

Manufactured by:



Bristol-Myers Squibb
Princeton, NJ 08543 USA

Marketed by:

Bristol-Myers Squibb Company
Princeton, NJ 08543
and
AstraZeneca Pharmaceuticals LP
Wilmington, DE 19850

1256316 1256317 SA-B0001A-07-09 Iss July 2009