

POLICY & PRACTICE

Uninsured Projections

The plight of the uninsured isn’t likely to be resolved anytime soon. More than one in four American workers under the age of 65 will be uninsured in 2013—nearly 56 million people—driven by the increasing inability to afford health insurance, reports a Health Affairs Web-exclusive article. Because growth in per capita health spending is expected to outpace median personal income by 2.4% every year, health care coverage will continue to decline because more Americans will find it unaffordable. “It is unlikely that we will be

able to solve the problem of the uninsured without some form of universal health insurance requiring contributions from some combination of employers, employees, and taxpayers,” the study indicated. The researchers based the estimates of the uninsured on federal projections of health spending, personal income, and other population characteristics.

Seniors Don’t Take Their Medicine

Four in 10 seniors don’t take their medications as prescribed—either because the drug regimens were too complex and cost-

ly, they didn’t think the drugs were helping them, or they didn’t think they needed them, a survey of 17,685 Medicare beneficiaries aged 65 years and older revealed. The 2003 survey, which was conducted prior to the enactment of the Medicare Modernization Act, found that drug coverage made a substantial difference in adherence rates. Thirty-seven percent of seniors without drug coverage reported cost-related nonadherence, compared with 22% of seniors with drug coverage. “Urging doctors and patients to talk more about these issues and developing systems to monitor quality and safety” could improve compliance, said Com-

monwealth Fund President Karen Davis. The survey was conducted by the Kaiser Family Foundation, the Commonwealth Fund, and Tufts-New England Medical Center and was posted as a Health Affairs online article.

Unreadable Privacy Forms

Privacy forms used by major health care institutions are often lengthy and too difficult for patients to read, according to a study published in the Journal of the American Medical Association (2005;293:1593-4). The Health Insurance Portability and Accountability Act requires that health care institutions inform patients in detail about the use of information obtained during medical care, through a notice of privacy form “written in plain English.” The median length of the forms was six pages, and often used inappropriately small type, and complex language that was unlikely to be understood by a “considerable proportion” of the populations served by these institutions, the report said. Privacy practice notices from 185 institutions listed in the 2004 U.S. News & World Report “best hospitals” issue were collected for the study.

Obesity in African Americans

Officials at the Department of Health and Human Services have awarded \$1.2 million to fight obesity among African Americans. The money will be used by a coalition of national organizations to fund prevention, education, public awareness, and outreach activities aimed at promoting a greater understanding of the impact of obesity. Nearly two-thirds of Americans are considered overweight or obese but the problem appears to be more prevalent in the African American community. Adult African American women have age-adjusted obesity rates of 49%, compared with 31% for adult white women, according to 1999-2002 data from the Centers for Disease Control and Prevention. African American girls and boys also had higher rates of overweight than did white children in the same age groups. HHS plans a second phase of the initiative targeting the Hispanic community.

Bill on Livestock Antibiotics

Sen. Edward M. Kennedy (D-Mass.) and Sen. Olympia J. Snowe (R-Maine) have introduced a bill to cut down on the amount of antibiotics used in livestock, citing evidence that increased antibiotic use in animals leads to reduced effectiveness in humans. “Antibiotics are among the greatest miracles of modern medicine, yet we are destroying them faster than the pharmaceutical industry can create replacements,” Sen. Kennedy said in a statement. “If doctors lose these critical remedies, the most vulnerable among us will suffer the most—children, the elderly, and persons with HIV / AIDS, who are most in danger of resistant infections.” The measure would require the Food and Drug Administration to withdraw approval for nontherapeutic use of eight classes of antibiotics in food-producing animals after 2 years if the use has not been proved harmless during that time. It also requires manufacturers of animal drugs or drug-containing feed to make their sales records available to government regulators for tracking emerging antimicrobial resistance.

—Jennifer Silverman

Combunox®
(Oxycodone HCl and Ibuprofen) Tablets

5mg/400 mg

FOREST LABORATORIES CII Rx only

Brief Summary: For complete details, please see full prescribing information for Combunox.

INDICATIONS AND USAGE
Combunox tablets are indicated for the short term (no more than 7 days) management of acute, moderate to severe pain.

CONTRAINDICATIONS
Combunox should not be administered to patients who have previously exhibited hypersensitivity to oxycodone HCl, ibuprofen, or any of Combunox's components, or in any situation where opioids are contraindicated. This includes patients with significant respiratory depression (in unmonitored settings or the absence of resuscitative equipment) and patients with acute or severe bronchial asthma or hypercarbia. Combunox is contraindicated in any patient who has or is suspected of having paralytic ileus. Combunox should not be given to patients who have experienced asthma, urticaria, or allergic-type reactions after taking aspirin or other NSAIDs. Severe anaphylactoid reactions to NSAIDs, some of which were fatal, have been reported in such patients (see WARNINGS - Anaphylactoid Reactions, and PRECAUTIONS - Pre-existing Asthma). Patients known to be hypersensitive to other opioids may exhibit cross-sensitivity to oxycodone.

WARNINGS
Misuse Abuse and Diversion of Opioids
Combunox contains oxycodone, which is an opioid agonist, and a Schedule II controlled substance. Opioid agonists have the potential for being abused and are sought by abusers and people with addiction disorders, and are subject to diversion. Combunox can be abused in a manner similar to other opioid agonists, legal or illicit. This should be considered when prescribing or dispensing Combunox in situations where the physician or pharmacist is concerned about an increased risk of misuse, abuse or diversion (see DRUG ABUSE AND DEPENDENCE).

Respiratory Depression
Oxycodone may produce dose-related respiratory depression by acting directly on the brain stem respiratory centers. Oxycodone HCl also affects the center that controls respiratory rhythm, and may produce irregular and periodic breathing. Respiratory depression occurs most frequently in elderly or debilitated patients, usually following large initial doses in non-tolerant patients, or when opioids are given in conjunction with other agents that depress respiration. Combunox should be used with extreme caution in patients with significant chronic obstructive pulmonary disease or cor pulmonale, and in patients having substantially decreased respiratory reserve, hypoxia, hypercapnia, or pre-existing respiratory depression. In such patients, even usual therapeutic doses of Combunox may decrease respiratory drive to the point of apnea.

Hypotensive Effect
Combunox, like all opioid analgesics, may cause severe hypotension in an individual whose ability to maintain blood pressure has been compromised by a depleted blood volume, or after concurrent administration with drugs such as phenothiazines or other agents which compromise vasomotor tone. Combunox may produce orthostatic hypotension in ambulatory patients. Combunox, like all opioid analgesics, should be administered with caution to patients in circulatory shock, since vasodilation produced by the drug may further reduce cardiac output and blood pressure.

Head Injury and Increased Intracranial Pressure
The respiratory depressant effects of opioids and their capacity to elevate cerebrospinal fluid pressure may be markedly exaggerated in the presence of head injury, intracranial lesions or a pre-existing increase in intracranial pressure. Furthermore, opioids produce adverse reactions that may obscure the clinical course of patients with head injuries.

Acute Abdominal Conditions
The administration of opioids may obscure the diagnosis or clinical course of patients with acute abdominal conditions.

Gastrointestinal (GI) Effects
Serious gastrointestinal toxicity, such as inflammation, bleeding, ulceration, and perforation of the stomach, small intestine or large intestine, can occur at any time, with or without warning symptoms, in patients treated with non-steroidal anti-inflammatory drugs (NSAIDs) such as ibuprofen. Minor upper GI problems, such as dyspepsia, are common and may also occur at any time during NSAID therapy. Therefore, physicians and patients should remain alert for ulceration and bleeding even in the absence of previous GI tract symptoms. Even short term therapy is not without risk.

NSAIDs should be prescribed with extreme caution in those with a prior history of ulcer disease or gastrointestinal bleeding. Most spontaneous reports of fatal GI events are in elderly or debilitated patients and, therefore, special care should be taken in treating this population. To minimize the potential risk for an adverse GI event the treatment period should be of the shortest possible duration. For high risk patients, alternate therapies that do not involve NSAIDs should be considered.

In addition to a past history of ulcer disease, pharmacoepidemiological studies have identified several other co-therapies or co-medical conditions that may increase the risk for GI bleeding such as treatment with oral corticosteroids, treatment with anticoagulants, longer duration of NSAID therapy, smoking, and alcoholism.

Anaphylactoid Reactions
Anaphylactoid reactions may occur in patients without known prior exposure to Combunox. Combunox should not be given to patients with the aspirin triad or a history of angioedema. The triad typically occurs in asthmatic patients who experience rhinitis with or without nasal polyps, or who exhibit severe, potentially fatal bronchospasm after taking aspirin or other NSAIDs. Fatal reactions to NSAIDs have been reported in such patients (see CONTRAINDICATIONS and PRECAUTIONS - Pre-existing Asthma). Emergency help should be sought when anaphylactoid reaction occurs.

Advanced Renal Disease
In patients with advanced kidney disease, treatment with Combunox is not recommended. However, if Combunox therapy must be initiated, due to the NSAID component, close monitoring of the patient's kidney function is advisable (see PRECAUTIONS - Renal Effects).

Pregnancy
As with other NSAID-containing products, Combunox should be avoided in late pregnancy because it may cause premature closure of the ductus arteriosus.

Interactions with Alcohol and Drugs of Abuse
Oxycodone may be expected to have additive effects when used in conjunction with alcohol, other opioids, or illicit drugs that cause central nervous system depression.

PRECAUTIONS
General
Special Risk Patients
As with any opioid analgesic agent, Combunox tablets should be used with caution in elderly or debilitated patients and those with severe impairment of hepatic, pulmonary or renal function, hypothyroidism, Addison's disease, acute alcoholism, convulsive disorders, CNS depression or coma, delirium tremens, kyphoscoliosis associated with respiratory depression, toxic psychosis, prostatic hypertrophy or urethral stricture. The usual precautions should be observed and the possibility of respiratory depression, postural hypotension, and altered mental states should be kept in mind.

Use in Pancreatic/Biliary Tract Disease
Combunox may cause spasm of the sphincter of Oddi and should be used with caution in patients with biliary tract disease, including acute pancreatitis. Opioids like Combunox may cause increases in the serum amylase level.

Cough Reflex
Oxycodone suppresses the cough reflex; as with other opioid containing products, caution should be exercised when Combunox is used postoperatively and in patients with pulmonary disease.

Effect on Diagnostic Signs
The antipyretic and anti-inflammatory activity of ibuprofen may reduce fever and inflammation, thus diminishing their utility as diagnostic signs in detecting complications of presumed noninfectious, noninflammatory painful conditions.

Hepatic Effects
As with other NSAIDs, ibuprofen has been reported to cause borderline elevations of one or more liver enzymes; this may occur in up to 15% of patients. These abnormalities may progress, may remain essentially unchanged, or may be transient with continued therapy. Notable (3 times the upper limit of normal) elevations of SGPT (ALT) or SGOT (AST) occurred in controlled clinical trials in less than 1% of patients. A patient with symptoms and/or signs suggesting liver dysfunction, or in whom an abnormal liver test has occurred, should be evaluated for evidence of the development of more severe hepatic reactions while on therapy with Combunox. Severe hepatic reactions, including jaundice and cases of fatal hepatitis, have been reported with ibuprofen as with other NSAIDs. Although such reactions are rare, if abnormal liver tests persist or worsen, if clinical signs and symptoms consistent with liver disease develop, or if systemic manifestations occur (e.g. eosinophilia, rash, etc.), Combunox should be discontinued.

Renal Effects
Caution should be used when initiating treatment with Combunox in patients with considerable dehydration. It is advisable to rehydrate patients first and then start therapy with Combunox. Caution is also recommended in patients with pre-existing kidney disease (see WARNINGS - Advanced Renal Disease).

As with other NSAIDs, long-term administration of ibuprofen has resulted in renal papillary necrosis and other renal pathologic changes. Renal toxicity has also been seen in patients in which renal prostaglandins have a compensatory role in the maintenance of renal perfusion. In these patients, administration of a nonsteroidal anti-inflammatory drug may cause a dose-dependent reduction in prostaglandin formation and, secondarily, in renal blood flow, which may

precipitate overt renal decompensation. Patients at greatest risk of this reaction are those with impaired renal function, heart failure, liver dysfunction, those taking diuretics and ACE inhibitors, and the elderly. Discontinuation of nonsteroidal anti-inflammatory drug therapy is usually followed by recovery to the pretreatment state.

Ibuprofen metabolites are eliminated primarily by the kidneys. The extent to which the metabolites may accumulate in patients with renal failure has not been studied. Patients with significantly impaired renal function should be more closely monitored.

Hematological Effects
Ibuprofen, like other NSAIDs, can inhibit platelet aggregation but the effect is quantitatively less and of shorter duration than that seen with aspirin. Ibuprofen has been shown to prolong bleeding time in normal subjects. Because this prolonged bleeding effect may be exaggerated in patients with underlying hemostatic defects, Combunox should be used with caution in persons with intrinsic coagulation defects and those on anticoagulant therapy. Anemia is sometimes seen in patients receiving NSAIDs, including ibuprofen. This may be due to fluid retention, GI loss, or an incompletely described effect upon erythropoiesis.

Fluid Retention and Edema
Fluid retention and edema have been reported in association with ibuprofen; therefore, the drug should be used with caution in patients with a history of cardiac decompensation, hypertension or heart failure.

Pre-existing Asthma
Patients with asthma may have aspirin-sensitive asthma. The use of aspirin in patients with aspirin-sensitive asthma has been associated with severe bronchospasm, which may be fatal. Since cross-reactivity between aspirin and other NSAIDs has been reported in such aspirin-sensitive patients, Combunox should not be administered to patients with this form of aspirin sensitivity and should be used with caution in patients with pre-existing asthma.

Aspirin
Aspirin: Patients with severe hepatic or renal disease, effects of therapy should be monitored with liver and/or renal function tests.

Drug Interactions
Oxycodone
Oxycodone is metabolized in part to oxymorphone via the cytochrome P₄₅₀ isoenzyme CYP2D6. While this pathway may be blocked by a variety of drugs (e.g., certain cardiovascular drugs and antidepressants), such blockade has not yet been shown to be of clinical significance with this agent. However, clinicians should be aware of this possible interaction.

Anticholinergics
The concurrent use of anticholinergics with oxycodone preparations may produce paralytic ileus.

CNS Depressants
Patients receiving narcotic analgesics, general anesthetics, phenothiazines, other tranquilizers, sedative-hypnotics or other CNS depressants (including alcohol) concomitantly with oxycodone may exhibit an additive CNS depression. Interactive effects resulting in respiratory depression, hypotension, profound sedation, or coma may result if these drugs are taken in combination with the usual dosage of oxycodone. When such combined therapy is contemplated, the dose of one or both agents should be reduced.

Mixed Agonist/Antagonist Opioid Analgesics
Agonist/antagonist analgesics (i.e., pentazocine, nalbuphine, butorphanol and buprenorphine) should be administered with caution to patients who have received or are receiving a course of therapy with a pure opioid agonist analgesic such as oxycodone. In this situation, mixed agonist/antagonist analgesics may reduce the analgesic effect of oxycodone and/or may precipitate withdrawal symptoms in these patients.

Monooamine Oxidase Inhibitors (MAOIs)
MAOIs have been reported to intensify the effects of at least one opioid drug causing anxiety, confusion and significant depression of respiration or coma. The use of oxycodone is not recommended for patients taking MAOIs or within 14 days of stopping such treatment.

Neuromuscular Blocking Agents
Oxycodone, as well as other opioid analgesics, may enhance the neuromuscular blocking action of skeletal muscle relaxants and produce an increased degree of respiratory depression.

Ibuprofen
ACE-inhibitors: Reports suggest that NSAIDs may diminish the antihypertensive effect of ACE-inhibitors. This interaction should be given consideration in patients taking Combunox concomitantly with ACE-inhibitors.

Aspirin
As with other products containing NSAIDs, concomitant administration of Combunox and aspirin is not generally recommended because of the potential of increased adverse effects.

Diuretics
Ibuprofen has been shown to reduce the natriuretic effect of furosemide and thiazides in some patients. This response has been attributed to inhibition of renal prostaglandin synthesis. During concomitant therapy with Combunox the patient should be observed closely for signs of renal failure (see PRECAUTIONS - Renal Effects), as well as diuretic efficacy.

Lithium
Ibuprofen has been shown to elevate plasma lithium concentration and reduce renal lithium clearance. This effect has been attributed to inhibition of renal prostaglandin synthesis by ibuprofen. Thus, when Combunox and lithium are administered concurrently, patients should be observed for signs of lithium toxicity.

Methotrexate
Ibuprofen, as well as other NSAIDs, has been reported to competitively inhibit methotrexate accumulation in rabbit kidney slices. This may indicate that ibuprofen could enhance the toxicity of methotrexate. Caution should be used when Combunox is administered concomitantly with methotrexate.

Warfarin
The effects of warfarin and NSAIDs on GI bleeding are synergistic, such that users of both drugs together have a greater risk of serious GI bleeding than users of either drug alone.

Carcinogenicity, Mutagenicity and Impairment of Fertility
Studies to evaluate the potential effects of the combination of oxycodone and ibuprofen on carcinogenicity, mutagenicity or impairment of fertility have not been conducted.

Pregnancy
Teratogenic Effects
Pregnancy Category C
Animal studies to assess the potential effects of the combination of oxycodone and ibuprofen on embryo-fetal development were conducted in the rat and rabbit model.

Pregnant rats were treated by oral gavage with combination doses of oxycodone:ibuprofen mg/kg/day (0.25:20, 0.5:40, 1.0:80, or 2.0:160) on days 7-16 of gestation. There was no evidence for developmental toxicity or teratogenicity at any dose, although maternal toxicity was noted at doses of 0.5:40 and above. The highest dose tested in the rat (2.0:160 mg/kg/day) is equivalent to the maximum recommended human daily dose (20:1600 mg/kg) on a body surface area (mg/m²) basis. This dose was associated with maternal toxicity (death, clinical signs, decreased BW).

Pregnant rabbits were treated by oral gavage with combination doses of oxycodone:ibuprofen (0.38:30, 0.75:60, 1.50:120 or 3.00:240 mg/kg/day) on gestation days 7-19. Oxycodone:ibuprofen treatment was not teratogenic under the conditions of the assay. Maternal toxicity was noted at doses of 1.5:120 (reduced body weight and food consumption) and 3:240 mg/kg/day (mortality). The no adverse effect level (NOAEL) for maternal toxicity, 0.75:60 mg/kg/day, is 0.75 fold the proposed maximum daily human dose based upon the body surface area. Developmental toxicity, as evidenced by delayed ossification and reduced fetal body weights, was noted at the highest dose, which is approximately 3 times the MRHD on a mg/m² basis, and is likely due to maternal toxicity. The fetal NOAEL of 1.50:120 mg/kg/day is approximately 1.5 times the MRHD on a mg/m² basis.

There are no adequate and well-controlled studies in pregnant women. Combunox should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Because of the ibuprofen component, Combunox should not be used during the third trimester of pregnancy because it could cause problems in the unborn child (premature closure of the ductus arteriosus and pulmonary hypertension in the fetus/neonate).

Labor and Delivery
Combunox should not be used during the third trimester of pregnancy due to the potential for ibuprofen to inhibit prostaglandin synthetase which may prolong pregnancy and inhibit labor. Oxycodone is not recommended for use in women during and immediately prior to labor and delivery because oral opioids may cause respiratory depression in the newborn.

Nursing Mothers
Ibuprofen is not transferred to breast milk in significant quantities. The American Academy of Pediatrics classified ibuprofen as compatible with breastfeeding. In studies using a 1 mg/mL assay, ibuprofen was not detected in the milk of lactating mothers. Oxycodone is excreted in human milk. Withdrawal symptoms and/or respiratory depression have been observed in neonates whose mothers were taking narcotic analgesics during pregnancy. Although adverse

effects in the nursing infant have not been documented, withdrawal can occur in breast-feeding infants when maternal administration of an opioid analgesic is discontinued.

Because of the potential for serious adverse reactions in nursing infants from the oxycodone present in Combunox, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use
In the placebo-controlled, clinical studies of pain following dental surgery, 109 patients between the ages of 14 and 17 years were administered a single dose of Combunox. No apparent differences were noted in the safety of Combunox in patients below and above 17 years of age. Combunox has not been studied in patients under 14 years of age.

Geriatric Use
Of the total number of subjects in clinical studies of Combunox, 89 patients were 65 and over, while 37 patients were 75 and over. No overall differences in safety were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out. However, because the elderly may be more sensitive to the renal and gastrointestinal effects of nonsteroidal anti-inflammatory agents as well as possible increased risk of respiratory depression with opioids, extra caution should be used when treating the elderly with Combunox.

ADVERSE REACTIONS
Listed below are the adverse event incidence rates from single dose analgesia trials in which a total of 2437 patients received either Combunox, ibuprofen (400 mg), oxycodone HCl (5 mg), or placebo. Adverse event information is also provided from an additional 334 patients who were exposed to Combunox in a multiple dose analgesia trial, without placebo or active component comparison arms, given up to four times daily for up to 7 days.

Adverse Events Which Occurred at a Frequency of ≥ 1% and at a Higher Incidence than in the Placebo Group in Single Dose Studies	5/400 mg (n=923)	400 mg Ibuprofen (n=913)	5 mg Oxycodone HCl (n=286)	Placebo (n=315)
Digestive				
Nausea	81 (8.8%)	44 (4.8%)	46 (16.1%)	21 (6.7%)
Vomiting	49 (5.3%)	16 (1.8%)	30 (10.5%)	10 (3.2%)
Flatulence	9 (1.0%)	7 (0.8%)	3 (1.0%)	0
Nervous System				
Somnolence	67 (7.3%)	38 (4.2%)	12 (4.2%)	7 (2.2%)
Dizziness	47 (5.1%)	21 (2.3%)	17 (5.9%)	8 (2.5%)
Skin and Appendages				
Sweat	15 (1.6%)	7 (0.8%)	4 (1.4%)	1 (0.3%)

Adverse events that were reported by at least 1% of patients taking Combunox but were observed at a greater incidence in the placebo treated patients were fever, headache and pruritus.

Adverse events that occurred in less than 1% and in at least two Combunox treated patients in Single Dose studies not listed above include the following: **Body as Whole:** abdominal pain, asthenia, chest pain, enlarged abdomen. **Cardiovascular System:** hypotension, syncope, tachycardia, vasodilation. **Digestive System:** constipation, dry mouth, dyspepsia, eructation, ileus. **Hemic and Lymphatic System:** anemia. **Metabolic and Nutritional Disorders:** edema. **Nervous System:** euphoria, insomnia, nervousness. **Respiratory System:** hypoxia, lung disorder, pharyngitis. **Urogenital System:** urinary retention.

Adverse events that occurred in the Multiple Dose study in at least 2% of patients treated with Combunox include the following: **Body as Whole:** asthenia (3.3%), fever (3.0%), headache (10.2%). **Cardiovascular System:** vasodilation (3.0%). **Digestive System:** constipation (4.5%), diarrhea (2.1%), dyspepsia (2.1%), nausea (25.4%), vomiting (4.5%). **Nervous System:** dizziness (12.2%), somnolence (17.4%).

Adverse events that occurred in less than 2% of and at least two Combunox treated patients in the Multiple Dose study not listed previously include the following: **Body as Whole:** back pain, chills, infection. **Cardiovascular System:** thrombophlebitis. **Hemic and Lymphatic System:** ecchymosis. **Metabolic and Nutritional Disorders:** hypokalemia. **Musculoskeletal System:** arthritis. **Nervous System:** abnormal thinking, anxiety, hyperkinesia, hypertension. **Skin and Appendages:** rash. **Special Senses:** amblyopia, taste perversion. **Urogenital System:** urinary frequency.

DRUG ABUSE AND DEPENDENCE
Combunox contains oxycodone, which is a mu-opioid agonist with an abuse liability similar to other opioid agonists and is a Schedule II controlled substance. Combunox, and other opioids used in analgesia, can be abused and are subject to criminal diversion.

Addiction is a primary, chronic, neurobiologic disease, with genetic, psychosocial, and environmental factors influencing its development and manifestations. It is characterized by behaviors that include one or more of the following: impaired control over drug use, compulsive use, continued use despite harm, and craving. Drug addiction is a treatable disease utilizing a multidisciplinary approach, but relapse is common.

"Drug seeking" behavior is very common in addicts and drug abusers. Drug-seeking tactics include emergency calls or visits near the end of office hours, refusal to undergo appropriate examination, testing or referral, repeated "loss" of prescriptions, tampering with prescriptions and reluctance to provide prior medical records or contact information for other treating physician(s). "Doctor shopping" to obtain additional prescriptions is common among drug abusers and people suffering from untreated addiction.

Abuse and addiction are separate and distinct from physical dependence and tolerance. Physical dependence usually assumes clinically significant dimensions after several days to weeks of continuous opioid use. Tolerance, in which increasingly large doses are required in order to produce the same degree of analgesia, is manifested initially by a shorter duration of analgesic effect, and subsequently by a decrease in the intensity of analgesia. The rate of development of tolerance varies among patients. Physicians should be aware that abuse of opioids can occur in the absence of true addiction and is characterized by misuse for non-medical purposes, often in combination with other psychoactive substances. Combunox, like other opioids, may be diverted for non-medical use. Record-keeping of prescribing information, including quantity, frequency, and renewal requests is strongly advised.

Proper assessment of the patient, proper prescribing practices, periodic re-evaluation of therapy, and proper dispensing and storage are appropriate measures that help to limit abuse of opioid drugs.

OVERDOSAGE
Following an acute overdosage, toxicity may result from oxycodone and/or ibuprofen.

Signs and Symptoms:
Acute overdosage with oxycodone may be manifested by respiratory depression, somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, constricted pupils, bradycardia, or hypotension. In severe cases death may occur.

The toxicity of ibuprofen overdose is dependent on the amount of drug ingested and the time elapsed since ingestion, although individual response may vary, necessitating individual evaluation of each case. Although uncommon, serious toxicity and death have been reported in the medical literature with ibuprofen overdosage. The most frequently reported symptoms of ibuprofen overdose include abdominal pain, nausea, vomiting, lethargy, and drowsiness. Other central nervous system symptoms include headache, tinnitus, CNS depression, and seizures. Cardiovascular toxicity, including hypotension, bradycardia, tachycardia, and atrial fibrillation, have also been reported.

Treatment:
In the treatment of opioid overdosage, primary attention should be given to the re-establishment of a patent airway and institution of assisted or controlled ventilation. Supportive measures (including oxygen and vasopressors) should be employed in the management of circulatory shock and pulmonary edema accompanying overdosage, as indicated. Cardiac arrest or arrhythmias may require cardiac massage or defibrillation. The narcotic antagonist naloxone hydrochloride is a specific antidote against respiratory depression, which may result from overdosage or unusual sensitivity to narcotics including oxycodone. An appropriate dose of naloxone hydrochloride should be administered intravenously with simultaneous efforts at respiratory resuscitation. Since the duration of action of oxycodone may exceed that of the naloxone, the patient should be kept under continuous surveillance and repeated doses of the antagonist should be administered as needed to maintain adequate respiration. Management of hypotension, acidosis and gastrointestinal bleeding may be necessary. In cases of acute overdosage, the stomach should be emptied through ipecac-induced emesis or gastric lavage. Orally administered activated charcoal may help in reducing the absorption and reabsorption of ibuprofen. Emesis is most effective if initiated within 30 minutes of ingestion. Induced emesis is not recommended in patients with impaired consciousness or overdoses greater than 400 mg/kg of the ibuprofen component in children because of the risk for convulsions and the potential for aspiration of gastric contents.

A Schedule CII Narcotic

© Forest Pharmaceuticals, Inc.
Combunox is a registered trademark of Forest Pharmaceuticals, Inc. • Manufactured under U.S. Patent Nos. 5,811,111 and 5,811,112

11/04
© 2004 Forest Laboratories, Inc.