

Common Symptoms Could Mask Hypothyroidism

BY DOUG BRUNK
San Diego Bureau

SAN DIEGO — Symptoms of hypothyroidism such as fatigue, depression, myalgias, cold intolerance, and menstrual abnormalities can be so common that they often blur the distinction between screening and case finding.

“Such symptoms are commonly seen in primary care, so you have to have a low threshold to find hypothyroidism,” Dr.

William A. Norcross said at the annual meeting of the American Academy of Family Physicians. He discussed other symptoms of hypothyroidism that can be easily overlooked:

► **Hyperlipidemia.** Lipid abnormalities are dependent on the degree of thyroid failure and are more pronounced in patients with thyroid-stimulating hormone levels above 10 mU/L. “So if you’re screening a patient for hypercholesterolemia and you detect it, you should give some con-

sideration to checking a TSH because replacement therapy not only will help the patient feel better, but will help the elevated LDL situation,” said Dr. Norcross of the department of family medicine at the University of California, San Diego.

► **Hyponatremia.** “There is a relationship between hypothyroidism and hyponatremia, especially in elderly patients,” he said. “It’s important to detect because the hyponatremia can be difficult to treat if you’re missing the underlying cause.”

► **Myalgias/myositis.** One of Dr. Norcross’s patients, a healthy-looking middle-aged man, presented with severe myalgias and a TSH level of 150 mU/L. “Every muscle in his body ached,” he recalled. “You couldn’t duplicate it with palpation. It was very vexing. Hypothyroidism turned out to be the cause of this muscle pain, and his symptoms went away with levothyroxine replacement.”

Such a case “is not an everyday occurrence, but it happens and is important to keep in mind,” said Dr. Norcross, who also directs the physician assessment and clinical education program at the university.

He also noted that certain drugs may affect thyroid function and thyroid test results. These include lithium, drugs that decrease TSH secretion (such as dopamine, glucocorticoids, and phenytoin), and drugs that increase TSH secretion (such as metoclopramide, domperidone, and amiodarone).

According to the United States National Health and Nutrition Examination Survey III, the overall incidence of hypothyroidism in the United States is 4.6%. It’s eight times more common in women, compared with men.

Dr. Norcross said that he had no conflicts to disclose. ■

novo nordisk
is changing
diabetesSM

Leadership... It’s time to think differently about what a pharmaceutical company can be. Novo Nordisk has dedicated itself to defeating diabetes, which is an interesting goal given that diabetes is the primary focus of our business. Success would undoubtedly force us to change, but then again, change is what we have always embraced.

Changing DiabetesSM means helping people with our actions, and not just with our products. It means staying true to our ideals and being respectful of the world we live in. And it means continuing to dedicate significant resources to diabetes research until a cure is found.

With leadership comes responsibility. We are committed to improving options for care now while partnering with others who share our mission and ultimate goal of defeating this disease.



Changing Diabetes is a service mark of Novo Nordisk A/S.

NovoMedLink.com
1-800-727-6500

Weekly Dose of Levothyroxine Is Safe, Preferred

CHICAGO — Once-weekly levothyroxine administration in hypothyroid women proved to be a safe and well-tolerated alternative to standard daily therapy in a Brazilian randomized trial.

Echocardiographic evaluation showed no differences between the two dosing regimens in terms of cardiovascular function. And most study participants preferred the convenience of once-weekly dosing, Dr. Gisah Carvalho said at the annual meeting of the American Thyroid Association.

She presented a 12-week, randomized crossover trial in which 19 hypothyroid women spent 6 weeks taking their normal daily levothyroxine dose and another 6 weeks in which they took seven times their regular daily dose once weekly.

Mean serum TSH was significantly higher after 6 weeks of weekly therapy, at 4.41 mIU/L, as compared with 3.38 mIU/L with daily therapy. Free thyroxine was lower: 1.0 ng/dL, as compared with 1.2 ng/dL with daily therapy.

Four hours after once-weekly levothyroxine administration the mean free thyroxine was 1.8 ng/dL, compared with 1.15 ng/dL 4 hours after daily dosing. Total triiodothyronine was unaffected, according to Dr. Carvalho of the Federal University of Paraná, Curitiba, Brazil.

The weekly dosing regimen is particularly appreciated by patients who find adherence to daily therapy challenging, said Dr. Carvalho, who disclosed no conflicts of interest.

—Bruce Jancin