Uncle Sam Needs YOU for Next Disaster Response

BY MARY ELLEN SCHNEIDER New York Bureau

DALLAS — Last year's Gulf Coast hurricanes and their devastating aftermath left many physicians wondering how they could help in future disasters, Dr. Joseph A. Scott, director of the division of prehospital and emergency health care at the University of Miami, said at the annual meeting of the National Medical Association.

There are a number of federal disaster

response teams for which physicians can volunteer, Dr. Scott said.

The National Disaster Medical System (NDMS) is a public-private partnership, based primarily in the Department of Homeland Security, which coordinates teams of medical providers to respond to storms, floods, airplane crashes, and even large-scale events like the Olympics or presidential inaugurations. The NDMS was set up to supplement state and local medical resources.

The disaster medical assistance teams (DMATs) make up one component of the NDMS. DMATs are 35-person teams that can be on the ground at a disaster anywhere around the country within 24 hours. A DMAT typically comprises physicians, nurses, paramedics, emergency medical technicians, pharmacists, respiratory therapists, psychologists, and social workers. All team members are cross-trained in medical, logistical, communications, and administrative aspects of the response, said Dr. Scott, who serves as medical officer for a DMAT based in south Florida.

These teams handle a variety of medical situations-from triage to primary medical care to trauma—once they are on the ground, he said. The DMAT usually rolls out the door with a cache of medications provided by the federal government. The medications include drugs for treatment as well as for dispensing purposes, because most pharmacies will be shut down in a disaster, Dr. Scott said. The NDMS also

OPANA® none Hydrochloride) (Oxymorp 5 mg and 10 mg

(For full DICATIONS AND USAGE

INDICATIONS

liability

uld result in ove. INGS: Drug roose an

agents OPANA, dministere

Impairment of OPANA ER (an extended-releas on of oxymorphone) in patients wit disease indicated oreater plasm than those with normal hepati CLINICAL PHARMACOLOGY

PRECAUTIONS

or upon st or mixed

Use with CNS Depressants The concomitant use of

effects resulting in res

and titrated slowly while side effects (see ADMINISTRATION).

Juring geste Noomental t of 25

in elderly (≥65 years of subjects (see CLINICAL

nan in younge

DOSAGE

orted in Placebo Is able lists adverse reactions that in at least 2% of patients in

4.4%

2.2%

iting

6.8% 6.5%

Adverse Reactions Reported in All Clinical A total of 591 patients were treated with OPANA in the Phase 2/3 controlled clinical

include

of life and

of OPANA ER, patients moderate to severe remains shown to have an increat ranging from 57-65% PHARMACOLOGY). These started cautionsly with low (see CLINICAL patients should be

Store at 25°C (77°F); excursions pe 15°-30°C (59°-86°F). [See USP

Rx Only DEA Order Form Required

Copyright © Endo Pharmaceuticals Inc. 2006 413242 E1/Rev. July, 200

te. Iral, OPANA should not be abruptly nued (see DOSAGE AND STRATION: Cessation of Therapy). rmation for Patients/Caregive e full prescribing information for rmation for patients).

sis, Impairment

supplies the equipment, but Dr. Scott said his group also accepts donated equipment to supplement that stock.

Emergency medicine training is ideal for these types of teams, he said. Experience with urgent care, trauma, and pediatrics is especially important for disaster deployments.

The usual deployment period for members of a DMAT is about 1-2 weeks, which includes predisaster staging. Physicians in the past were required to commit to a 2week deployment, but that has been changing to 1 week for physicians only, he said.

Volunteering to be part of an organized response team can forestall many of the

problems that can occur at a disaster site when individual, well-meaning volunteers flood the scene. An overabundance of volunteers can divert resources, Dr. Scott said. And unsolicited volunteers often lack appropriate training, equipment, and supplies. They may also be outside the formal accountability system and may lack proper credentials, he said.

A big fear when volunteers-especially physicians-show up without formal coordination is that no one will be at home to take care of the regular emergencies, he said. Thus, physicians who can't volunteer for a DMAT can still help in a disaster while staying home, Dr. Scott said. Not everyone has the flexibility-either at home or at work—to volunteer for a 1- to 2-week deployment. But physicians can make a difference by volunteering to cover shifts for a colleague who is a member of a response team.

There are also increasing opportunities for physicians to volunteer for similar teams at the state level. Physicians can check with their state emergency management offices to find out if their state has a team. he said.

Other federal teams that need physician volunteers include:

► National medical response teams. These are 50-member teams that deploy

OPANA[®] ER orphone Hydrochl nded-Release Tat ablets 5 mg, 10 mg, 20 mg and 40 mg

ity similar to other opioid e can be abused in a manne

esic. IA ER TABLETS are to be swallowed a and are not to be broken, chewed blved, or crushed. Taking broken ed, dissolved, or crushed OPANA EF

in patients requiring continuous, ck opioid treatment for an extended

ER is not intended for use as a prn

ER is not indicated for pain in the te post-operative period (12-24 hours surgery) for patients not previously taking eccause of the risk of oversedation and y depression requiring reversal with opioid te

During the second period of time. **ICATIONS** is contrained and in patients with a is contrained to patients with a is contrained by the oxymorphone rymorphine analogs such as codoine, rymorphine analogs such as other ingregations of CPANA EFF; in th moderate or severe hepatic rymorphine analogs such as red such as; patients with respiratory (in the absence of resuscilative x in unmonitored settings), acute or hall asthma, hypercabla, and in any has or is suspected of having paralytic

ER is not indicated for pain in the te post-operative period (the first 12-24 ate post-operation of the pain is mean plowing surgery), or if the pain is mean of to persist for an extended period of time. A Erit is only indicated for post-operative use batient is already receiving the drug prior to or or if the post-operative pain is expected to a persist for an extended in a difficult part of the post-operative pain is expected to a standard persist for an extended

re not to be proken, chewed, ssolved. Taking broken, chewed, issolved OPANA ER TABLETS the rapid release and absorption fatal dose of oxymorphone.

or diversion. ER tablets may be abused by crushing, , snorting or injecting the product. These s will result in the uncontrolled delivery of id and pose a significant risk to the abuser

A. ressio. Joensing L Authority for n. Detect abuse or dive. a with Alcohol and D. one may be expected to h. an used in conjunction with alco. # illicit drugs that cause central ne. depression because respirato. Interpret of the conjunction of the second depression because respirato. Interpret of the conjunction of the second of the conjunction of the second of the conjunction of the second a single does of 40 mg of OPANA sted volunteers. The results ted volumeters. The results debression of the second of revent the proper transgement of pain, professionals should contact their State I Licensing Board or State Controlled Authority for information on how to detect eluse or diversion of this product

of OPAINA ET and the thank the thank

subjects. Following the of 240 mL of 20% etha average by 31% and subjects. Following the of 240 mL of 4% ethanol, e concomitant administration and the C_{max} increased up to 260% in increased on concomitant administration of the concomitant administration d agonists and is a Sc OPANA ER and oth and Diversion of Opioids). diction is characterized by a preocu-procurement, hoarding, and abuse -medicinal purposes. Drug addi e, utilizing a multi-disciplinary appro . avior is very common to

shopping" (visiting mi additional prescription

Abuse of OPANA Erripoon death. This risk is increase of OPANA ER with alcohol addition, parenteral dru

is (see PRECAUTIONS: Usage in they and PRECAUTIONS: Labor and

ts of opioid analgesics and their rate cerebrospinal fluid pressure sodiation following CO₂ retention) exaggerated. Furthermore, opioid produce effects on pupillary resclusives, which may obscure of turner increases

aution to patients in circulatory shoci lation produced by the drug may cardiac output and blood pressure. ic Impairment: A study of OPANA is with hepatic disease indicated a concentrations than those with

with caution in patients with mild impairmen patients should be started with the lowest d titrated elowly while carefully monitoring effects. OPANA ER is contraindicated for with packate and source broatini impairm

JTIONS :Opioid analgesics should be used with properially when combined with other drugs, especially when combined will other driv ould be reserved for cases where of opioid analgesia outweigh the km risks of respiratory depression, alte tate and postural hypotension. OPANA be used with caution in elderly ad patients and in patients who are km sensitive to contral nervous sys ants, such as those with cardiovasci ury, renal, or heaptic disease. system aution in the

ER should be used with ca conditions: acute prtical insufficiency (e.g., CNS depression or com alcoholism, Addison's a; delirium

administration of oxymorphone may obscure diagnosis or clinical course in patients with e abdominal conditions. Oxymorphone may be added with convulsive NA ER is intended for use in patients who re more than several days continuous Ambulatory Surgery and Po-OPANA ER is not indicated oracional (administration pre-constance)

against the possible hazards

atic function (see CLINICAL PHARMACOLOGY), INA ER should be used with caution in patients mild impairment. These patients should be ted with the lowest dose and litrated slowly o carefully monitoring for side effects. OPANA is contraindicated for patients with moderate

and severe hepatic impairment CONTRAINDICATIONS, WARNINGS, DOSAGE AND ADMINISTRATION).

weight, gender differences were not observ CLINICAL PHARMACOLOGY). In clinical

COSAGE AND ADMINISTRATION). Renal Impairment: In a study of OPANA ER, patients with moderate to severe renal impairment were shown to have an increase in bioavailability ranging from 57-65% (see CLINICAL PHARMACOLOGY). These patients should be started cautiously with lower doese of OPANA ER and titrated slowly while carefully monitored for side effects (see DOSAGE AND ADMINISTRATION). Cender Differences: When normalized the termination termination.

were slightly higher among females than for both OPANA ER subjects and placebo

Adverse Reactions Reported in All Clinical Trials: A total of 2011 patents were treated with OPANA ER in the Phase 23 controlled and open-label aincial Insias. The aliated Iralis consisted of patients with moderate to severe chronic pain and post surgical pain. The adverse reaching are presented in the blowing mariner: most common, common and

already receiving OPANA ER as

like other opioids, du is is a common , especially after i opioid analgesia. Cau nitor for decreased b

neral, OPANA ER should not be abruptly intinued. However, OPANA ER, like other development of withdrawal symptoms by slowly tapering the daily dose (see DOSAGE AND ADMINISTRATION: Cessation of Therapy)

use in Drug and Alcohol Addiction: OPANA ER s not approved for use in detoxification or t approved for use in decompanying analysis of a point of addiction enance treatment of opioid addiction ever, the history of an addictive disorder does cessarily preclude the use of this medication treatment of chronic pain. These patients will a intensive monitoring for signs of misuse

abuse, or addiction. Drug-Drug Interactions: Oxymorphone is highly metabolized principally in the liver and undergoes reduction or conjugation with glucuronic acid to form both active and inactive metabolites (see

CNS depressants: The conc. CNS depressants: The conc. CNS depressants including s, tranquilizers, general a atzines, other opioids, and al additive CNS depressant effect

reported at least once by part OPANA ER in the clinical tri hydrochloride tested positive in both the rat and mouse in vivo micronucleus assays. Impairment of fertility: The dose of oxymorphone that produced no adverse effects on reproductive findings in lemale rats is 0.6 fold the human dose of 40 mg every 12 hours on a body surface area basis. **Programery:** The safety of using expresphere pregnancy has not basen established with regard to possible adverse effects on telad development. The hancy has no been still development. The of OPANA ER in pregnancy, in nursing mothers, women of child-bearing potential requires that possible benefits of the drug be weighed to the mother and the Eye dis

Effects: Pregnancy Category C: Effects: Pregnancy Category C: hydrochloride administration did not rmations at any doses evaluated during tal toxicity studies in rats (<25 or rabbits (<50 mg/kg/day).

istention, agital central nervou

e but are not pathognomoni of hemorrhagic or ischem similar findings). Marked m

cautiously. Of the total number of subjects in clinical studies of OFANA ER, 27 percent were 65 and over, while 9 percent were 75 and over. No overall differences in effectiveness were observed between these subjects and younger subjects. There were several adverse events that were more frequently observed

AND HANDLING

call Board, State Board of F rol Board for information on ant diversion of this product. at 25°C (77°F); excursions permitted to 15°-(59°-86°F). [See USP Controlled Room

Dispense in tight container as defined in the USP, with a child-resistant closure (as required).

iticals In lds Ford, Pennsylvania 19317 ifactured by:

Novartis Consumer Treasures Lincoln, NE 68517 TIMERx[®]-N is a registered Trac Charmaceuticals Co., Danbury, er Health Inc

aceuticals Inc. 2006 413542 E1/July. 2006

less often than DMATs. They are generally tasked to respond to nuclear, biologic, and chemical incidents.

▶ Disaster mortuary operational response teams. These teams include pathologists, forensic pathologists, fingerprint experts, and dental assistants who help to identify victims' remains in a disaster. There are about 9 or 10 such teams in the country.

▶ International medical/surgical re**sponse teams.** These teams generally take care of American civilians overseas, Dr. Scott said. They are similar to DMATs, but generally deploy with a trauma surgeon and anesthesiologist; they can perform surgery in the field.

Urban search and rescue teams. These teams specialize in on-site medical treatment for victims trapped in confined spaces, such as those resulting from structural collapses. The physicians on these teams frequently care for both victims and responders, he said.

Physicians who are deployed as part of a federal team become federal employees during deployment and do collect a small paycheck, Dr. Scott said.

Physicians get other protections during their deployment, including federal liability protection.

But there are some disadvantages, Dr. Scott said. The initial credentialing process can take 6 months or more.

Overall, the experience can be very rewarding, Dr. Scott said. It's an opportunity to provide good care to appreciative patients with minimal paperwork. "You're actually really practicing emergency medicine the way most us went to medical school to do it," he said.

To learn more about volunteering for a disaster response team, visit www.ndms.fema.gov or contact Jack Beall of the Federal Emergency Management Agency at jack.beall@dhs.gov.

How to Prepare For Deployment

r. Joseph A. Scott, who serves on both a disaster medical assistance team and an international medical/surgical response team, offered tips for physicians preparing for deployment:

 Update your immunizations before heading into the field.

▶ Get packed. Physicians should carry cash, water, food, protective clothing, medications, flashlight, batteries, sleeping bag, gasoline, insect repellent, and their credentials, Dr. Scott said. Credit cards will be useless if the power is out. Physicians need to carry enough of their own supplies that they don't end up becoming victims themselves, he said. ► Take care of your own mental

health. This means making sure that your home and work situations are under control during deployment, he said. Make sure in advance that colleagues can cover your shifts. This will go a long way toward making you more effective during deployment, he said.

rrhea ahdr

and may produce respiratory depression and psycho-physiologic effects in neonates. OPANA EF is not recommended for use in women during and immediately prior to labor, when use of shorter acting analgesics or other analgesic techniques are more appropriate. A specific optical antagonist, such as natoxome or nalmetere, should be available for reversal of opticid-induced respiratory depression in the neonate. ric Use: Safety and effectiveness of OPANA pediatric patients below the age of 18 years ER in pediatric parents sector have not been established. Geriatric Use: OPANA ER should be used with caution in ekkery patients. The plasma levels of oxymorphone are about 40% higher in elderly (265