

BY KERRI WACHTER
Senior Writer

In a retrospective study of 12,538 women with postmenopausal osteoporosis,

Women were said to have postmenopausal osteoporosis if they had one or

Within 6 months of initiating therapy, 57% of the 9,117 women on weekly therapy and 62% of the 3,421 women on daily therapy were considered to have ad-

The research was funded by Amgen Inc., which is currently investigating a fully monoclonal antibody for osteoporosis. ■

FOR ORAL INHALATION ONLY

In general, the technique for administering XOPENEX HFA Inhalation Aerosol to children is similar to that for adults. Children should use XOPENEX HFA Inhalation Aerosol under adult supervision, as instructed by the patient's physician.

During marketing experience of racemic albuterol, various congenital anomalies, including cleft palate and limb defects, have been rarely reported in the offspring of patients being treated with racemic

mimetic agents, can cause adverse reactions such as angina, vertigo, central nervous system stimulation, slurred speech, headache, and drying or irritation of the oropharynx.

Rx only.

 SEPRACOR 12/05

©2005 SEPRACOR INC., MARLBOROUGH, MA 01752

Mean NTx levels measured prior to PTH infusion did not change between baseline and week 6 in the control group, but NTx levels increased by 24% in group 1, by 16% in group 2, and by 11% in group 3. Serum NTx increased during PTH infusion in all groups at all time points.

Pages 30a—30b◇