



**Benjamin H. Crenshaw, MD; Kortnee Y. Roberson, MD; James J. Stevermer, MD, MSPH**  
Department of Family and Community Medicine, University of Missouri-Columbia (Drs. Crenshaw and Stevermer); University of Chicago, Department of Family Medicine (Dr. Roberson)

**DEPUTY EDITOR**  
**Anne Mounsey, MD**  
University of North Carolina at Chapel Hill

# A new adjunctive Tx option for venous stasis ulcers

Adding simvastatin to standard wound care improves ulcer healing rates and times as well as the patient's quality of life.

## PRACTICE CHANGER

Consider adding simvastatin 40 mg/d to standard wound care and compression for patients with venous stasis ulcers.<sup>1</sup>

## STRENGTH OF RECOMMENDATION

**B:** Based on a high-quality randomized controlled trial (RCT).

Evangelista MT, Casintahan ME, Villafuerte LL. Simvastatin as a novel therapeutic agent for venous ulcers: a randomized, double-blind, placebo-controlled trial. *Br J Dermatol.* 2014;170:1151-1157.

## ILLUSTRATIVE CASE

A 74-year-old woman with chronic lower extremity edema seeks treatment for a nonhealing venous stasis ulcer. For the past 9 months, she's been wearing compression stockings and receiving intermittent home-based wound care, but nothing seems to help. She asks if there's anything else she can try.

**V**enous stasis ulcers affect 1% of US adults and lead to substantial morbidity and more than \$2 billion in annual health care expenditures.<sup>1,2</sup> Edema management—generally limb elevation and compression therapy—has been the mainstay of therapy. Treatment can be lengthy, and ulcer recurrences are common.<sup>2,3</sup>

Statins have been found to help wound healing through their diverse physiologic (pleiotropic) effects. Evidence shows they can be beneficial for treating diabetic foot ulcers,<sup>4</sup> pressure ulcers,<sup>5</sup> and ulcerations asso-

ciated with systemic sclerosis and Raynaud's phenomenon.<sup>6</sup> Evangelista et al<sup>1</sup> investigated whether adding a statin to standard wound care and compression could improve venous stasis ulcer healing.

## STUDY SUMMARY

### Ulcers are more likely to close when a statin is added to standard care

This randomized, double-blind, placebo-controlled trial was performed at a large medical center in the Philippines. It was designed to assess the efficacy and safety of simvastatin 40 mg/d for venous ulcer healing when combined with standard treatment (compression therapy, limb elevation, and standard wound care).<sup>1</sup>

Researchers randomized 66 patients ages 41 to 71 who'd had one or more venous ulcers for at least 3 months to receive either simvastatin 40 mg/d (N=32) or an identical appearing placebo (N=34). Patients were excluded if they were pregnant, had an ulcer that was infected or >10 cm in diameter, or were taking any medication that could interact with a statin. Patients were stratified according to ulcer diameter (≤5 cm and >5 cm). There was no statistically significant difference between the 2 groups in the duration of venous ulceration (3.80 years in the placebo group vs 3.93 years in the simvastatin group) or incidence of diabetes (5% in the placebo group vs 3% in the simvastatin group).

## INSTANT POLL

Do you ever prescribe a statin for the treatment of a venous stasis ulcer?

- Yes  
 No

jfponline.com

The primary outcome was the proportion of patients whose ulcers completely healed at 10 weeks. Secondary outcomes were measures of the total surface area healed and healing time, and Dermatology Life Quality Index (DLQI) scores. Baseline ulcer diameter and surface area and DLQI scores were obtained prior to therapy. The same dermatologist, who was blinded to the patients' assigned group, evaluated all patients every 2 weeks until wound closure or for a maximum of 10 weeks.

Overall, 90% of the patients who received simvastatin had complete ulcer closure at 10 weeks, compared with 34% of patients in the control group (relative risk [RR]=0.16; 95% confidence interval [CI], 0.05-0.47; number needed to treat [NNT]=2).

Among patients with ulcers  $\leq 5$  cm, 100% of the ulcers healed in the simvastatin group, compared to 50% in the control group (RR=0.10; 95% CI, 0.01-0.71; NNT=2). Perhaps more importantly, in patients with ulcers  $>5$  cm, 67% of the ulcers in the simvastatin group had closure with a mean healing time of 9 weeks, whereas none of the ulcers of this size closed in the control group (RR=0.33; 95% CI, 0.12-0.84; NNT=1.5), and the mean healed area was significantly larger in patients who received simvastatin (28.9 cm<sup>2</sup> vs 19.6 cm<sup>2</sup>;  $P=.03$ ).

In addition, in the simvastatin group, healing times were significantly reduced (7.53 $\pm$ 1.34 weeks vs 8.55 $\pm$ 1.13 weeks) and quality of life (as evaluated by DLQI scoring) significantly improved compared to the control group.

Study dropouts (8%; 2 in the placebo group and 3 in the intervention group) were minimal. Using intention-to-treat analysis and worst-case scenarios for dropouts did not affect the primary outcome. There were no withdrawals for adverse reactions.

#### WHAT'S NEW

### Statins offer significant benefits for treating venous stasis ulcers

This is the first human study to investigate the use of a statin in venous stasis ulcer healing. This intervention demonstrated significant improvements in healing rate and time, a

very small NNT for benefit, and improved patient quality of life compared to placebo.

#### CAVEATS

### Results were found in a carefully selected group of patients

Many wounds will heal with compression therapy alone, as occurred in this study, where 50% of ulcers  $\leq 5$  cm treated with standard therapy healed, albeit at a somewhat slower rate. Adding another medication to the regimen when these patients generally have multiple comorbidities should always prompt caution.

The study by Evangelista et al<sup>1</sup> was performed in a select population, and the exclusion criteria included the use of some commonly prescribed medications, such as angiotensin-converting enzyme inhibitors. No data were collected on patient body mass index, which is a risk factor for delayed healing. The prevalence of obesity is lower in the Philippines than in the United States, and it is uncertain what role this difference would have in the statin's effectiveness. Further studies, especially those conducted with a less selective population, would better clarify the generalizability of this intervention.

We found the results of this study impressive. The methods reported are rigorous and consistent with standard RCT methodologies. This is the only study of a statin in human venous stasis disease, but studies in animals—and studies of statins for other types of ulcers in humans—have consistently suggested benefit. It seems hard to argue against adding this low-cost, low-risk intervention.

#### CHALLENGES TO IMPLEMENTATION

There are no known barriers to implementing this practice.

JFP

#### ACKNOWLEDGEMENT

The PURLs Surveillance System was supported in part by Grant Number UL1RR024999 from the National Center For Research Resources, a Clinical Translational Science Award to the University of Chicago. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center For Research Resources or the National Institutes of Health.

Copyright © 2015. The Family Physicians Inquiries Network. All rights reserved.

CONTINUED

**>**  
Sixty-seven percent of ulcers  $>5$  cm in the simvastatin group had closure, while none of those in the control group did.

References

1. Evangelista MT, Casintahan ME, Villafuerte LL. Simvastatin as a novel therapeutic agent for venous ulcers: a randomized, double-blind, placebo-controlled trial. *Br J Dermatol*. 2014;170:1151-1157.
2. Collins L, Seraj S. Diagnosis and treatment of venous ulcers. *Am Fam Physician*. 2010;81:989-996.
3. The Australian Wound Management Association Inc, New Zealand Wound Care Society Inc. Australian and New Zealand clinical practice guideline for prevention and management of venous leg ulcers. 2011. Australian Government National Health and Medical Research Council Web site. Available at: [http://www.nhmrc.gov.au/\\_files\\_nhmrc/publications/attachments/ext003\\_venous\\_leg\\_ulcers\\_aust\\_nz\\_0.pdf](http://www.nhmrc.gov.au/_files_nhmrc/publications/attachments/ext003_venous_leg_ulcers_aust_nz_0.pdf). Accessed February 13, 2015.
4. Johansen OE, Birkeland KI, Jørgensen AP, et al. Diabetic foot ulcer burden may be modified by high-dose atorvastatin: A 6-month randomized controlled pilot trial. *J Diabetes*. 2009;1:182-187.
5. Farsaei S, Khalili H, Farhoud ES, et al. Efficacy of topical atorvastatin for the treatment of pressure ulcers: a randomized clinical trial. *Pharmacotherapy*. 2014;34:19-27.
6. Abou-Raya A, Abou-Raya S, Helmi M. Statins: potentially useful in therapy of systemic sclerosis-related Raynaud's phenomenon and digital ulcers. *J Rheumatol*. 2008;35:1801-1808.

Join the official job board of  
THE JOURNAL OF  
**FAMILY  
PRACTICE**

 **MEDJOBNETWORK.com**  
Physician • NP/PA Career Center

Search 1000s of Jobs and Apply in 1 Click  
And get FREE benefits including...

- Access to 30+ medical web sites
- Medical news and clinical advice
- E-Alert and Newsletters on your smart phone
- Online CME and MD-IQ Quizzes
- Coverage of over 200 meetings
- And much more!

**Julian Knight**  
Digital Sales Director, Classifieds  
Phone: 973-206-2317  
[jknight@frontlinemedcom.com](mailto:jknight@frontlinemedcom.com)  
[www.medjobnetwork.com](http://www.medjobnetwork.com)