



Thiazolidinediones and heart failure

(APRIL 2006)

TO THE EDITOR: Dr. Wilson Tang, in his critical review of the available evidence about thiazolidinediones (TZDs) and heart failure,¹ should surely have discussed the adverse effects outcomes in the largest TZD trial to date, PROactive.² This study, which included 5,238 patients with type 2 diabetes with evidence of macrovascular disease, randomized patients to pioglitazone or placebo to be taken in addition to their other glucose-lowering drugs and other medications. This study found a 39% increase in the relative risk of hospitalization for congestive heart failure, as well as a statistically significant increase in any report of heart failure in the pioglitazone group. The number needed to harm for pioglitazone was 33 (treat 33 patients for 34 months to cause one heart failure event). In Dr. Tang's article, this was listed as "not statistically significant," but clearly from the PROactive study this was statistically significant (with a *P* value of less than .001 for any report of heart failure, .003 for heart failure not needing hospital admission, and .007 for heart failure needing hospital admission). Either this was a misprint in Dr. Tang's article, or being on the speaker bureau for Takeda Pharmaceuticals (the maker of pioglitazone) has caused massive myopia.

In the PROactive study, both the primary and preplanned secondary end points were negative in the study. Main secondary outcomes, which were statistically significant, were compiled after the end of the study, but apparently before the data sets were opened. This is highly unusual.

Credible sources have both discounted the conclusions of the PROactive study and questioned its methodology.³⁻⁸ In fact, one commentary in *The Lancet* discussed using data from the United Kingdom Prospective Diabetes Study to simulate the same reductions in hemoglobin A_{1c} and blood pressure as in the pioglitazone group, and found the same outcomes in cardiovascular events but a risk reduction in heart failures.⁴

It is time to stop talking in *may's* and *might's* and to look at the data from an unbiased, non-pharmaceutical-sponsored perspective. The role of TZDs in type 2 diabetes remains unclear.

PONTUS L. JADERHOLM, PharmD
Pharmacist, Drug Information
Kaiser Permanente Northwest

REFERENCES

1. Tang WHW. Do thiazolidinediones cause heart failure? A critical review. *Cleve Clin J Med* 2006; 73:390-397.
2. Dormandy JA, Charbonnel B, Eckland DJ, et al. Secondary prevention of macrovascular events in patients with type 2 diabetes in the PROactive Study (PROspective pioglitAzone Clinical Trial in macroVascular Events): a randomised controlled trial. *Lancet* 2005; 366:1279-1289.
3. Goldstein MR. PROactive study [comment]. *Lancet* 2006; 367:23.
4. Guillausseau PJ. PROactive study [comment]. *Lancet* 2006; 367:23.
5. Gaede P, Parving H-H, Pedersen O. PROactive study [comment]. *Lancet* 2006; 367:23-24.
6. Yudkin JS, Freemantle N. PROactive study [comment]. *Lancet* 2006; 367:24-25.
7. Holman RR, Retnakaran R, Farmer A, Stevens R. PROactive study [comment]. *Lancet* 2006; 367:25-26.
8. Isley WL. Pioglitazone did not reduce a composite end-point of macrovascular complications and increased risk for heart failure in type 2 diabetes with macrovascular disease. *ACP Journal Club* 2006; 144:34.

IN REPLY: I thank Dr. Jaderholm for correctly pointing out the errors in my article regarding the risk of heart failure in the PROactive study.¹

In the PROactive study, after 3 years of follow-up, nonadjudicated heart failure events had occurred in 10.8% of patients in the pioglitazone group vs 7.5% in the placebo group; the difference was statistically significant (*P* < .0001). Patients taking pioglitazone had a 5.7% rate of hospitalization for heart failure vs 4.1% in the placebo group, which was also statistically significant (*P* = .003). Edema without heart failure occurred in 21.6% of patients in the pioglitazone group vs 13% in the placebo group. These data were emphasized in one of the summary key points at the beginning of my paper (page 390): "The Prospective Pioglitazone Clinical Trial in Macrovascular Events (PROactive; *Lancet* 2005; 366:1279-1289) found that the use of a TZD was associated with a 16% reduction in the combined end point of heart attack, stroke, and death, despite increased rates of edema and hospi-



talization for heart failure.”

The intent of my article was to provide a critical review of the literature in order to address the question, “Do TZDs cause heart failure?” This is an important scientific question, since the development of heart failure in patients with diabetes has been associated with poorer long-term survival.² It was my intention to point out that people who take TZDs clearly have a risk of developing fluid retention, but the risk of developing heart failure is less clear.

In fact, all glucose-lowering drugs have been implicated to some degree with regard to their heart failure risks, with insulin being the most prominent.³ However, the clinical diagnosis of heart failure is often subjective and nonadjudicated, and therefore definitive clinical evidence is limited.

Nevertheless, placebo-controlled echocardiographic studies have not directly linked fluid retention caused by TZDs with worsening cardiac performance.⁴ Careful analysis from the Kaiser Permanente Northern California Diabetes Registry of 23,440 patients with type 2 diabetes also did not find evidence that short-term TZD use was associated with an elevated risk of hospitalization from heart failure compared with other standard, first-line diabetes therapies.⁵ It is also reassuring that rates of death from heart failure in the PROactive study were similar between the TZD and the placebo groups (25/2605 vs 22/2633, $P = .634$), despite the aforementioned increase in reported rates of edema and hospitalizations for heart failure.⁶

In a recent analysis of 16,417 elderly diabetic patients discharged after hospitalization for heart failure, the use of insulin sensitizers (TZDs and metformin, both contraindicated in heart failure) was associated with a lower death rate compared with drugs other than insulin sensitizers.⁷ The discrepancy between the development of heart failure reported in patients with diabetes treated with TZDs and the potential benefit of fewer deaths is interesting and will require further insights from upcoming large clinical trials.

W.H. WILSON TANG, MD
Department of Cardiovascular
Medicine
Cleveland Clinic

■ REFERENCES

1. **Tang WHW.** Do thiazolidinediones cause heart failure? A critical review. *Clev Clin J Med* 2006; 73:390–397.
2. **Bertoni AG, Hundley WG, Massing MW, Bonds DE, Burke GL, Goff DC Jr.** Heart failure prevalence, incidence, and mortality in the elderly with diabetes. *Diabetes Care* 2004; 27:699–703.
3. **Nichols GA, Koro CE, Guillion CM, Ephross SA, Brown JB.** The incidence of congestive heart failure associated with antidiabetic therapies. *Diabetes Metab Res Rev* 2005; 21:51–57.
4. **St John Sutton M, Rendell M, Dandona P, et al.** A comparison of the effects of rosiglitazone and glyburide on cardiovascular function and glycemic control in patients with type 2 diabetes. *Diabetes Care* 2002; 25:2058–2064.
5. **Karter AJ, Ahmed AT, Liu J, Moffet HH, Parker MM.** Pioglitazone initiation and subsequent hospitalization for congestive heart failure. *Diabet Med* 2005; 22:986–993.
6. **Dormandy JA, Charbonnel B, Eckland DJ, et al.** Secondary prevention of macrovascular events in patients with type 2 diabetes in the PROactive Study (PROspective pioglitAzone Clinical Trial in macroVascular Events): a randomised controlled trial. *Lancet* 2005; 366:1279–1289.
7. **Masoudi FA, Inzucchi SE, Wang Y, Havranek EP, Foody JM, Krumholz HM.** Thiazolidinediones, metformin, and outcomes in older patients with diabetes and heart failure: an observational study. *Circulation* 2005; 111:583–590.

CORRECTION

Thiazolidinediones and heart failure

(APRIL 2006)

The article “Do thiazolidinediones cause heart failure? A critical review?” by Dr. Wilson Tang in the April 2006 issue (*Cleve Clin J Med* 2006; 73:390–397) contained several errors regarding the interpretation of the PROactive study (*Lancet* 2005; 366:1279–1289). The

CCJM article incorrectly stated that the differences between the pioglitazone and placebo groups were not statistically significant for any report of heart failure, for heart failure needing hospital admission, or for the occurrence of edema without heart failure. In fact, the pioglitazone group showed significantly higher rates of any report of heart failure and of heart failure needing hospital admission. The errors occurred as the article was being revised in the galley stage.