

# A Pharmacist-Led Transitional Care Program to Reduce Hospital Readmissions in Older Adults

Rebecca Rottman-Sagebiel, PharmD; Nicole Cupples, PharmD; Chen Pin Wang, PhD; Seth Cope; Stephanie Pastewait, PharmD; Hanna Braden; Daniel MacCarthy, MS; Alicia Conde, MA; Melody Moris; Eneida-Yvette Gonzalez; and Sara Espinoza, MD, MS, AGSF

Medication reconciliation and patient education during admission and after discharge helped older patients remain independent at home.

Author affiliations are listed at the end of this article.  
**Correspondence:** Sara Espinoza (espinozas2@uthscsa.edu)

There will be 53 million older adults in the US by 2020.<sup>1</sup> Increasing age often brings medical comorbidities and prescriptions for multiple medications. An increasing number of prescribed medications combined with age-related changes in the ability to metabolize drugs makes older adults highly vulnerable to adverse drug events (ADEs).<sup>2</sup> In addition, older adults often have difficulty self-managing their medications and adhering to prescribed regimens.<sup>3</sup> As a result, ADEs can lead to poor health outcomes, including hospitalizations, in older adults.

Medication errors and ADEs are particularly common during transitions from hospital to home and can lead to unnecessary readmissions, a major cause of wasteful health care spending in the US.<sup>4,5</sup> More than \$25 billion are estimated to be spent annually on hospital readmissions, with Medicare picking up the bill for \$17 billion of the total.<sup>6,7</sup> Researchers have found that the majority of ADEs following hospital discharge are either entirely preventable or at least ameliorable (ie, the negative impact or harm resulting from the ADE could have been reduced).<sup>8</sup>

To address these issues, we undertook a clinical demonstration project that implemented a new transitional care program to improve the quality of care for older veterans transitioning from the Audie L. Murphy Veterans Memorial Hospital of the

South Texas Veterans Health Care System (STVHCS) in San Antonio to home. The Geriatrics Medication Education at Discharge project (GMED) falls under the auspices of the San Antonio Geriatrics Research Education and Clinical Center (GRECC). Clinical demonstration projects are mandated for US Department of Veterans Affairs (VA) GRECCs to create and promote innovative models of care for older veterans. Dissemination of successful clinical demonstration projects to other VA sites is strongly encouraged. The GMED program was modeled after the Boston GRECC Pharmacological Intervention in Late Life (PILL) program.<sup>9</sup> The PILL program, which focuses on serving older veterans with cognitive impairment, demonstrated that a postdischarge pharmacist telephone visit for medication reconciliation leads to a reduction in readmission within 60 days of discharge.<sup>9</sup> The goals of the GMED program were to reduce polypharmacy, inappropriate prescribing and 30-day readmissions.

## METHODS

The project was conducted when a full-time clinical pharmacy specialist (CPS) was available (May-September 2013 and April 2014-March 2015). This project was approved as nonresearch/quality improvement by the University of Texas Health Science Center Institutional Review Board, which serves the STVHCS. Consent was not required.

## Eligibility

Patients were identified via a daily hospital database query of all adults aged  $\geq 65$  years admitted to the hospital through Inpatient Medicine, Neurology, or Cardiology services within the prior 24 hours. Patients meeting any of the following criteria based on review of the Computerized Patient Record System (CPRS) by the team geriatrician and CPS were considered eligible: (1) aged  $\geq 70$  years prescribed  $\geq 12$  outpatient medications; (2) aged  $\geq 65$  years with a medical history of dementia; (3) aged  $\geq 65$  years prescribed outpatient medications meeting Beers criteria<sup>10</sup>; (4) age  $\geq 65$  years with  $\geq 2$  hospital admissions (including the current, index admission) within the past calendar year; or (5) aged  $\geq 65$  years with  $\geq 3$  emergency department visits within the past calendar year. For the first polypharmacy criterion, patients aged  $\geq 70$  years were selected instead of aged  $\geq 65$  years so as not to exceed the capacity of 1 CPS. Twelve or more medications were used as a cutoff for polypharmacy based on prior quality improvement information gathered from our VA geriatrics clinic examining the average number of medications taken by older veterans in the outpatient setting.

Patients were excluded if they were expected to be discharged to any facility where the patient and/or the caregiver were not primarily responsible for medication administration after discharge. Patients who met eligibility criteria but were not seen by the transitional program pharmacist (due to staff capacity) were included in this analysis as a convenience comparison group of patients who received usual care. Patients were not randomized. All communication occurred in English, but this project did not exclude patients with limited English proficiency.

A program support assistant conducted the daily query of the hospital database. The pharmacist conducted the chart review

**TABLE 1** Group Characteristics

Variables	Intervention Group (N = 388)	Control Group (N = 1,189)	Total (N = 1,577)	P Value
Age, mean (SD), y	74.9 (7.6)	75.2 (8.35)	75.1 (8.2)	< .01
Female, No. (%)	14 (3.6)	26 (2.2)	40 (2.5)	.12
Ethnic group, No. (%)				.18
Non-Hispanic white	319 (82.2)	892 (75.0)	1,211 (76.8)	
Black	22 (5.7)	96 (8.1)	118 (7.5)	
Hispanic	112 (28.9)	334 (28.1)	446 (28.3)	
Frail 1 year prior to baseline, No. (%)	233 (60.1)	668 (57.9)	921 (58.4)	.45
Admitted 1 year prior to the first index admission, No. (%)	155 (40.0)	447 (37.6)	602 (38.2)	.41
Charlson Comorbidity Index, mean (SD)	4.4 (2.8)	4.27 (2.9)	4.30 (2.8)	.43


to determine eligibility and delivered the intervention. Eligible patients were selected at random for the intervention with the intention of providing the intervention to as many veterans as possible.

## THE GMED INTERVENTION

The GMED program included 2 phases, which were both conducted by a CPS with oversight from a senior CPS with geriatric pharmacology expertise and an internist/geriatrician. The CPS carrying out the transitional care program was involved in the planning and design of the project and met weekly with the geriatrician. The Figure provides an overview of the intervention.

The first phase of the transitional care program included an individual, face-to-face meeting between the CPS and the patient during the hospitalization. If a veteran was not present in the room at the time of an attempted visit, the pharmacist made 2 additional attempts (3 total) to include the patient in the transitional care program during the hospitalization. The CPS performed medication reconciliation and provided medication education regarding administration and usage of the patient's medications, using an open-ended format.<sup>11</sup> The caregiver, if any, was included in the discussion either at the bedside or by telephone following the face-to-face visit with the patient. The CPS communicated recommendations regarding appropriateness of therapy (including any

**FIGURE** Geriatrics Medication Education at Discharge



potential barriers to medication adherence) to the medical team (including the attending, resident[s], and interns) in person or by telephone and through documentation in the CPRS. The recommendations were based on the clinical expertise of the CPS as well as on guidelines for prescribing in older adults.<sup>10,12</sup> The CPS used a checklist to ensure all components of the intervention were completed (Appendices 1 and 2).

The second component of the transitional care program included a telephone visit within 2 to 3 days of discharge, conducted by the same CPS who performed the face-to-face visit. The purpose of the telephone visit was to perform medication reconciliation, identify and rectify medication errors, provide further patient education, and assist in facilitating appropriate follow-up by the patient's primary care provider (PCP), if required. At a minimum, veterans were asked a series of questions pertaining to their concerns about medication regimens, receipt of newly prescribed medications at discharge, additional education regarding medications after the CPS encounter during hospitalization, and whether the veteran required assistance with the medication regimen in the home setting. Follow-up questions were asked as needed to clarify and identify potential medication problems. All information from this telephone encounter was communicated to the PCP through CPRS documentation and

by telephone as needed.

#### Data Collection

A standardized questionnaire was used prospectively for patients in the transitional care program group to assess patient education, primary residence, presence of a caregiver, fall history, medication adherence, and cognitive status (using Mini-Cog).<sup>13</sup> Additional information (patient age, number of outpatient medications prior to and following the admission, presence of Beers criteria outpatient medications prior to and following the admission, new outpatient prescriptions, and changes to existing prescriptions as a result of the hospitalization) was gathered prospectively from patient interviews or from chart review.

For patients included in the comparison group, a retrospective administrative chart review was conducted to collect information such as age, sex, ethnic group, admission within 1 year prior to index admission, frailty, and Charlson Comorbidity Index (CCI) score, a method of categorizing comorbidities of patients based on the diagnosis codes found in administrative data.<sup>14</sup> Each comorbidity category has an associated weight (from 1 to 6), based on the adjusted risk of mortality or resource use, and the sum of all the weights results in a single comorbidity score for a patient (0 indicates no comorbidities; higher scores predict greater risk of mortality or increased resource use).

We used the index developed from 17 disease categories. The range for CCI was 0 to 25. Frailty was defined as the presence of any of the following frailty-related diagnoses: anemia; fall, head injury, other injury; coagulopathy; electrolyte disturbance; or gait disorder. These diagnoses are either primary frailty characteristics within the frailty phenotype or have been shown in prior studies to be associated with the frailty phenotype.<sup>15-18</sup> While more widely accepted frailty definitions exist, these other definitions require direct examination of the patient and could not be used in this project because we did not directly interact with the comparison group.<sup>16,19</sup> The frailty definition used has been previously identified as a predictor of health care utilization and 30-day readmission in a veteran population.<sup>20</sup> Whether or not the CPS detected a postdischarge medication error was recorded. All CPS recommendations were documented.

An index admission was defined as a hospital admission that occurred during the project period. Thirty-day readmission was defined as a hospital admission that occurred within 30 days of the discharge date of an index admission. Each index admission was considered individually for readmission (yes vs no) even if it occurred in the same patient over the project period. A 30-day readmission was not considered an index admission. An admission that occurred after a 30-day readmission was considered a subsequent index admission. Patients who died in the hospital were not included in this analysis, as they would not have participated in the entire intervention.

### Statistical Analysis

We compared characteristics between patients who received GMED and patients who never received GMED (comparison group). Generalized estimating equations (GEE) were used to determine whether the rate of 30-day readmission (yes vs no) in the transitional care program group differed from that of the comparison group. In our GEE analysis, we assumed a binomial distribution and the logit link to model the log-odds of readmission as a linear function of transitional care program status (yes vs no) and other covariates, including age, frailty, hos-

**TABLE 2** Patient Population Characteristics Seen by CPS (N = 378)

Characteristics	
Education, No. (%)	
Less than high school	61 (16.1)
High school graduate or some college	237 (62.7)
College degree or greater than college degree	80 (21.2)
Comorbid conditions	
Hypertension, No. (%)	350 (90.4)
Type 2 diabetes mellitus, No. (%)	225 (58.1)
Chronic obstructive pulmonary disease, No. (%)	113 (29.3)
Congestive heart failure, No. (%)	111 (28.7)
Glomerular filtration rate, mL/min, mean (SD)	53.3 (31.7)
Geriatric syndromes and medications	
Cognitive impairment or dementia noted, No. (%)	102 (27.0)
Caregiver assists with medications, No. (%)	124 (32.8)
Fall within the past year, No. (%)	167 (44.2)
Mini-Cog screens positive for possible cognitive impairment, No. (%)	168 (44.4)
Outpatient medications on admission, mean (SD), No.	15.5 (5.8)
Outpatient medications at discharge, mean (SD), No.	15.7 (5.9)
Patient received a prescription for a new outpatient medication at discharge, No. (%)	234 (61.9)
Patient had a change in the dose or frequency of $\geq 1$ of prescribed outpatient medications, No. (%)	180 (47.6)
Outpatient prescribed Beers medications at admission, No. (%)	166 (43.9)
GMED CPS recommendation at inpatient admission, No. (%)	240 (63.5)
A medication error was detected by GMED CPS follow-up, No. (%)	124 (32.8)

Abbreviations: CPS, clinical pharmacy specialist; GMED, Geriatrics Medication and Education.

pital admission within 1 year prior to the index admission, and CCI score as covariates. Thirty-day readmission status associated with each index admission was coded as 1 for a readmission within 30 days of the discharge date of the index admission, or 0 for no readmission within 30 days.

Transitional care program status was determined whether or not the individual received the transitional care program for each index admission. This analysis allowed us to model repeated measures of index admissions as a function of the project period and whether the patient was seen by the GMED CPS during the index admission. The patient identifier was used as a cluster variable in the GEE analysis. Inverse propensity scores of receiving GMED at the index admission were adjusted as weights in the GEE

**TABLE 3** Generalized Estimating Equations Analysis of Effect of Intervention Group on Readmission Adjusting for Propensity Scores<sup>a</sup>

Intervention Groups	Odds Ratio Estimate (SE)	95% Confidence Interval	P Value
Intervention group, unadjusted	0.74 (0.12)	0.54-1.01	.06
Intervention group, adjusted for follow-up time	0.53 (0.14)	0.32-0.88	.05
Intervention group, adjusted for follow-up time, age	0.53 (0.14)	0.32-0.88	.05
Intervention group, adjusted for follow-up time, age, prior admission within last year	0.54 (0.15)	0.32-0.90	.02
Intervention group, adjusted for follow-up time, age, prior admission within last year, frailty	0.55 (0.15)	0.33-0.90	.02
Intervention group, adjusted for follow-up time, age, prior admission within last year, frailty, ethnic group	0.54 (0.15)	0.32-0.90	.02

<sup>a</sup>All analyses shown are adjusted for length of follow-up.

analysis to minimize confounding and, hence, to strengthen the causal interpretation of the effect of the transitional care program. If there was ≥ 1 index admission, the GMED status (yes vs no) at the initial index admission was used as the dependent variable to calculate propensity scores. The propensity scores of transitional care program status were derived from the logistic regression analysis that modeled the log-odds of receiving the transitional care program at the index admission as a linear function of age, CCI, frailty, and prior hospitalization during the 1-year period prior to the index admission.

**RESULTS**

The GMED CPS saw 435 patients during the project period; 47 (10.8%) died prior to 30 days and were excluded, leaving 388 patients who received the transitional care program included in this evaluation. Another 1,189 patients met the eligibility criteria but were not included and were included in the comparison group. Patients in the transitional care program group were similar to those receiving usual care in the comparison group with regard to sex, ethnic group, frailty status, and CCI score (Table 1).

Data from the CPS-patient interviews and chart reviews were available for 378 of the 388 patients (Table 2). Patients were primarily

male, non-Hispanic white, with a high school education. More than half (65%) the patients were admitted for a new diagnosis or clinical condition. The majority of patients had diabetes mellitus, and about one-third had chronic obstructive pulmonary disease, congestive heart failure, or cognitive impairment. Although about 60% of patients were prescribed a new medication as a result of the hospital admission, the number of medications from admission to discharge did not differ significantly ( $15.4 \pm 5.5$  vs  $15.7 \pm 5.8$ ;  $P = .08$ ).

The 30-day readmission rate was 15.6% for the transitional care program group and 21.9% for the comparison group. Three hundred seventy-one patients received the transitional care program only once, 16 patients received the transitional care program twice (ie, they had 2 index admissions during the study period and received the intervention both times), and 1 patient received the transitional care program 3 times.

In an unadjusted GEE model, the odds ratio (OR) for readmission in the transitional care program group was 0.74 (95% CI, 0.54-1.0,  $P = .06$ ) compared with the usual care group (Table 3). After covariate adjustment, the OR for readmission was 0.54 (95% CI, 0.32-0.90,  $P = .02$ ).

Thirty-five percent of patients had ≥ 1 CPS-recommended change in their treatment at the time of the inpatient admission (Table 4). The most common recommendation was discontinuation of at least 1 medication (23.0%), followed by correcting the medication reconciliation list that was on record for the admission (17.8%). Thirty-nine percent of patients had ≥ 1 CPS-recommended change in their treatment at the time of the follow-up phone call. The most common recommendation was to clarify medication instructions for the patient and/or caregiver and provide medication education (33.7%). Other common

**TABLE 4** Summary of Recommendations Made by the Clinical Pharmacy Specialist

<b>Recommendations made at the time of inpatient visit (N = 326)<sup>a</sup></b>	<b>No.</b>	<b>%</b>
Discontinue any medication	75	23.0
Correct an incomplete or inaccurate medication reconciliation list	58	17.8
Start a new medication for new symptoms or condition	38	11.7
Recommend a change in medication(s) due to Beers Criteria	36	11.0
Discontinue a medication that the patient reported he was no longer taking (but was listed on the medication list and medication reconciliation on record)	34	10.4
Restart one or more medications that the patient reported he was not taking but is clinically indicated	14	4.3
Change or modify redundant medications	12	3.7
Decrease a medication dosage	12	3.7
Recommend a change in medication for patient's preference (ie, for dosing or ease of administration)	12	3.7
Recommend to resume medications that were held on admission while waiting evaluation but that had not been restarted	10	3.1
Recommend renal dose of medication(s)	8	2.5
Recommend laboratory assessment for medication monitoring	9	2.5
Correct an incorrect medication dose based on mg/kg	6	1.8
Suggest home health services	2	0.6
<b>Recommendations made at the time of discharge (N = 83)<sup>b</sup></b>	<b>No.</b>	<b>%</b>
Reeducation for patient not following recommendations from discharge	28	33.7
Medication reconciliation was incorrect	14	16.9
Poor communication between inpatient team and primary care physician	12	14.5
Limited medication supply (not given sufficient refill on discharge)	11	13.3
Inadequate follow-up for medications requiring frequent monitoring (eg, Coumadin)	11	13.3
Patient needed medical reevaluation, evaluation for drug-disease interaction, and/or lab follow-up	3	3.6
Drug-drug interaction	2	2.4
Discontinue a medication recommended at discharge or recommended a change to outpatient regimen	2	2.4

<sup>a</sup>35% of patients had a recommendation at the time of the inpatient pharmacist visit.

<sup>b</sup>39% of patients had a recommendation at the time of discharge.

recommendations were to correct a medication reconciliation (16.9%) and communicate pertinent information about the admission to the PCP (14.5%).

## DISCUSSION

We developed a transitional care program for hospitalized older veterans to improve the transition from hospital to home. After adjusting for clinical factors, GMED was associated with 26% lower odds of readmission within 30 days of discharge compared with that of the control group. The GMED CPS made changes to the medical regimen both during the inpatient admission as well as after discharge to correct medication errors and educate patients.

In addition, GMED led to a reduction in the number of prescribed medications, which impacts inappropriate polypharmacy—a significant problem in older adults, which contributes to ADEs.<sup>21</sup> Our intervention was patient centered, as all decisions and education regarding medication management were

tailored to each patient, taking into account medical and psychosocial factors.

Studies of similar programs have shown that a pharmacist-based program can improve outcomes in patients transitioning from hospital to home. A meta-analysis of 19 studies that evaluated the effectiveness of pharmacy-led medication reconciliation interventions at the time of a care transition showed that compared with usual care a pharmacist intervention led to reduced medication discrepancies.<sup>22</sup> In this meta-analysis, medication discrepancies of higher clinical impact were more easily identified through pharmacy-led interventions than with usual care, suggesting improved safety. Although not all studies have shown a clear reduction in readmission rates or other health care utilization, the addition of clinical pharmacist services in the care of inpatients has generally resulted in improved care with no evidence of harm.<sup>23</sup>

Based on these findings and collaboration with another GRECC, we designed our program to focus on older adults with

## APPENDIX 1 Checklist for Pharmacist Interaction With the Patient<sup>a</sup>

<p><b>1. Inquire about the patient's understanding of the medication renewal/refill process</b></p> <ul style="list-style-type: none"> <li>(a) Does the patient know how to request a refill or renewal of his/her medications?</li> <li>(b) Does the patient know how many days prior to needing the medication he/she should submit the refill or renewal request?</li> <li>(c) Does the patient receive all the medications requested reliably?</li> </ul>
<p><b>2. Inquire about the patient's support system and tools available for medication management</b></p> <ul style="list-style-type: none"> <li>(a) With whom does the patient reside?</li> <li>(b) What system, if any, does the patient use for medication management (ie, pill box, reminders)?</li> <li>(c) Does the patient require assistance from a caregiver for medication management?</li> </ul>
<p><b>3. Inquire about the patient's adherence to the prescribed medication regimen</b></p> <ul style="list-style-type: none"> <li>(a) Have you ever stopped taking your medication because you were feeling better?</li> <li>(b) Have you ever stopped taking your medication because you were feeling worse?</li> <li>(c) What do you do if you miss your medication (ie, question about doubling doses, taking doses too close together, etc)</li> <li>(d) How many times a week do you miss medications?</li> </ul>
<p><b>4. Review each medication with the patient and/or caregiver, including:</b></p> <ul style="list-style-type: none"> <li>(a) Understanding of the indication for each medication</li> <li>(b) Medication frequency</li> <li>(c) Potential adverse effects</li> <li>(d) Difficulties administering (ie, pill splitting, bottle opening, medication specific concerns)</li> </ul>
<p><b>5. Ask about all prescribed medications, including:</b></p> <ul style="list-style-type: none"> <li>(a) Active prescriptions from the VA</li> <li>(b) Expired prescriptions from the VA</li> <li>(c) Non-VA prescription medications (over-the-counter, herbal medications, and supplements)</li> </ul>
<p><b>6. Perform a brief geriatric assessment to assess the patient's abilities and risks with the current medication regimen and to determine needs for medication management at discharge</b></p> <ul style="list-style-type: none"> <li>(a) Cognitive screening (perform Mini-Cog assessment)</li> <li>(b) Falls screening (Has the patient fallen within the past year?)</li> </ul>
<p><b>7. Inform patient and provide resources</b></p> <ul style="list-style-type: none"> <li>(a) Inform patient and/or caregiver that medications may change during admission</li> <li>(b) Inform patient and/or caregiver to keep their medication list with them at all times</li> <li>(c) Provide patient with a pill box, if indicated, and instruct on its use</li> <li>(d) Inform patient and/or caregiver that the pharmacist will follow-up with a phone call a few days after discharge (inquire about the best telephone number to use)</li> </ul>

<sup>a</sup>This checklist uses an open-ended question format to encourage the most accurate information from the patient. Items do not need to be reviewed in this particular order.

polypharmacy, cognitive impairment, high-risk medication usage, and/or a history of high health care use.<sup>9</sup> Our findings add to the growing body of evidence that a CPS-led transitional care program results in reduced polypharmacy and reduced unnecessary hospital readmissions. Further, our findings have demonstrated the effectiveness of this type of program in a practical, clinical setting with veteran patients.

At the time of project inception, we believed that the majority of our interventions would occur postdischarge. We were some-

what surprised that a major component of GMED was suggested interventions by our pharmacist at the time of admission. We believe that because the CPS made suggestions during admission, we prevented postdischarge ADEs. A frequent intervention corrected the medication reconciliation on file at admission. This finding also was seen in another study by Gleason and colleagues, which examined medication errors at admission for 651 adult medicine inpatients.<sup>24</sup> This study found that more than one-third of patients had medication reconciliation errors.

Further, older age ( $\geq 65$  years) was associated with increased odds of medication errors in this study.

Of note, a survey of hospital-based pharmacists indicated medication reconciliation is the most important role of the pharmacist in improving care transitions.<sup>25</sup> The pharmacists stated that detection of errors at the time of admission is very important. The pharmacists further reported that additional education and counseling for patients with poor understanding of their medications was also important. Our findings support these findings and the use of a pharmacist as part of the medical team to improve medication reconciliation and education.

### Limitations

A limitation of GMED is that we monitored only admissions to our hospital; therefore, we did not account for any hospitalizations that may have occurred outside the STVHCS. Another limitation is that this was not a randomized controlled trial, and we used a convenience sample of patients who met our criteria for eligibility but were not seen due to time constraints. This introduces potential bias such that patients admitted and discharged on nights or weekends when the CPS was not available were not included in the transitional care program group, and these patients may fundamentally differ from those admitted and discharged Monday through Friday.

However, Khanna and colleagues found that night or weekend admission was not associated with 30-day readmission or other worse outcomes (such as length of stay, 30-day emergency department visit, or intensive care unit transfer) in 857 general medicine admissions at a tertiary care hospital.<sup>26</sup> Every effort was made to include as many eligible patients as possible in the transitional program group, and we were able to demonstrate that the patients in the 2 groups were similar. Frailty and prior hospital admission were more prevalent, although not significantly so, in the transitional program group, suggesting that any selection bias would have actually attenuated—not enhanced—the observed effect of the transitional program. Although the transitional program group patients

## APPENDIX 2 Checklist for Review of Hospital Admission Information and Pharmacist Interaction With Medical Team

### 1. Review the medical chart, keeping in mind information obtained from the patient and/or caregiver, and determine appropriateness of the prescribed medication regimen, including:

- (a) Renal dosing
- (b) Beers Criteria guidelines
- (c) START/STOP criteria
- (d) Applicable disease clinical guidelines
- (e) Benefit vs harm, including considerations of quality of life and patient preferences

### 2. Discuss considerations for potential changes in therapy with the inpatient medical team, including:

- (a) Stopping 1 or more medications
- (b) Starting 1 or more medications
- (c) Obtaining additional laboratory measurements, if needed, to determine whether therapy is still warranted

### 3. Inform inpatient medical team about the following:

- (a) The updated medication list if applicable
- (b) Ensuring that any medication changes made during the hospital admission are incorporated into the updated medication list and documented in the medical chart at discharge
- (c) Ensuring that an updated medication list is provided to the patient at discharge
- (d) That the patient will be contacted by the pharmacist a few days after discharge to review and clarify changes and ensure that the patient is taking medications appropriately
- (e) That a note will be included in the electronic medical record to reflect the findings and considerations discussed with the medical team

were slightly younger by 0.3 years, they were similar in frailty status and CCI score.

## CONCLUSION

The GMED program was associated with reduced 30-day hospital readmission, discontinuation of unnecessary medications, and corrected medication errors and discrepancies. We propose that a CPS-based transitional care program can improve the quality of care for older patients being discharged to home.

### Acknowledgments

Supported by funding from the Veterans Health Administration T21 Non-Institutional Long-Term Care Initiative and VA Office of Rural Health and the San Antonio Geriatrics Research, Education, and Clinical Center. The sponsor did not have any role in the design, methods, data collection, or analysis, and preparation.

### Author Contributions

R. Rottman-Sagebiel developed the transitional program concept and design and executed the program implementation, interpretation of data, and preparation of the manuscript. S. Pastewait, N. Cupples, A. Conde, M. Moris, and E. Gon-



zalez assisted with program design and implementation. S. Cope assisted with interpretation of data and preparation of the manuscript. H. Braden assisted with interpretation of data. D. MacCarthy assisted with data management and statistical analysis. C. Wang and S. Espinoza developed the program concept and design, performed statistical analysis and interpretation of data, and helped prepare the manuscript.

### Author disclosures

The authors report no actual or potential conflicts of interest with regard to this article.

### Disclaimer

The opinions expressed herein are those of the authors and do not necessarily reflect those of *Federal Practitioner*, Frontline Medical Communications Inc., the US Government, or any of its agencies.

### Author Affiliations

**Rebecca Rottman-Sagebiel**, **Nicole Cupples**, and **Stephanie Pastewart** are Clinical Pharmacy Specialists; **Chen Pin Wang** is a Biostatistician; **Seth Cope** and **Hanna Braden** are Medical Students; **Daniel MacCarthy** is a Data Analyst; **Melody Moris** is a Project Manager; **Eneida-Yvette Gonzalez** is a Program Support Assistant; **Alicia Conde** is a Research Assistant and **Sara Espinoza** is a Geriatrician at the University of Texas Health Science Center in San Antonio; all at the Geriatrics Research, Education and Clinical Center (GRECC) at the South Texas Veterans Health Care System (STVHCS) in San Antonio, Texas.

### Advances in Geriatrics

Advances in Geriatrics features articles focused on quality improvement/quality assurance initiatives, pilot studies, best practices, research, patient education, and patient-centered care written by health care providers associated with Veteran Health Administration Geriatric Research Education and Clinical Centers. Interested authors can submit articles at [editorialmanager.com/fedprac](mailto:editorialmanager.com/fedprac) or send a brief 2 to 3 sentence abstract to [fedprac@mdedge.com](mailto:fedprac@mdedge.com) for feedback and publication recommendations.

### References

1. Vincent GK, Wilkoff VA. The Next Four Decades: The Older Population in the United States: 2010 to 2050. US Department of Commerce, Economics and Statistics Administration, US Census Bureau; 2010.
2. Merle L, Laroche ML, Dantoine T, Charmes JP. Predicting and preventing adverse drug reactions in the very old. *Drugs Aging*. 2005;22(5):375-392.
3. Shi S, Mörike K, Klotz U. The clinical implications of ageing for rational drug therapy. *Eur J Clin Pharmacol*. 2008;64(2):183-199.
4. Coleman EA, Min Sj, Chomiak A, Kramer AM. Posthospital care transitions: patterns, complications, and risk identification. *Health Serv Res*. 2004;39(5):1449-1465.
5. Berwick DM, Hackbarth AD. Eliminating waste in US health care. *JAMA*. 2012;307(14):1513-1516.
6. Price Waterhouse Coopers Health Research Institute. The Price of Excess: Identifying Waste in Healthcare Spending. Price Waterhouse Coopers Health Research Institute; 2008.
7. Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. *N Engl J Med*. 2009;360(14):1418-1428.
8. Forster AJ, Murff HJ, Peterson JF, Gandhi TK, Bates DW. The incidence and severity of adverse events affecting patients after discharge from the hospital. *Ann Intern Med*. 2003;138(3):161-167.
9. Paquin AM, Salow M, Rudolph JL. Pharmacist calls to older

- adults with cognitive difficulties after discharge in a Tertiary Veterans Administration Medical Center: a quality improvement program. *J Am Geriatr Soc*. 2015;63(3):571-577.
10. The American Geriatrics Society 2015 Beers Criteria Update Expert Panel. American Geriatrics Society 2015 updated Beers Criteria for potentially inappropriate medication use in older adults. *J Am Geriatr Soc*. 2015;63(11):2227-2246.
11. Greenwald JL, Halasyamani L, Greene J, et al. Making inpatient medication reconciliation patient centered, clinically relevant and implementable: a consensus statement on key principles and necessary first steps. *J Hosp Med*. 2010;5(8):477-485.
12. Gallagher P, Ryan C, Byrne S, Kennedy J, O'Mahony D. STOPP (Screening Tool of Older Person's Prescriptions) and START (Screening Tool to Alert doctors to Right Treatment). Consensus validation. *Int J Clin Pharmacol Ther*. 2008;46(2):72-83.
13. Borson S, Scanlan J, Brush M, Vitaliano P, Dokmak A. The mini-cog: a cognitive 'vital signs' measure for dementia screening in multi-lingual elderly. *Int J Geriatr Psychiatry*. 2000;15(11):1021-1027.
14. Deyo RA, Cherkin DC, Ciol MA. Adapting a clinical comorbidity index for use with ICD-9-CM administrative databases. *J Clin Epidemiol*. 1992;45(6):613-619.
15. Chaves PH, Semba RD, Leng SX, et al. Impact of anemia and cardiovascular disease on frailty status of community-dwelling older women: the Women's Health and Aging Studies I and II. *J Gerontol A Biol Sci Med Sci*. 2005;60(6):729-735.
16. Fried LP, Tangen CM, Walston J, et al; Cardiovascular Health Study Collaborative Research Group. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci*. 2001;56(3):M146-M156.
17. Walston J, McBurnie MA, Newman A, et al; Cardiovascular Health Study. Frailty and activation of the inflammation and coagulation systems with and without clinical comorbidities: results from the Cardiovascular Health Study. *Arch Int Med*. 2002;162(20):2333-2341.
18. Stookey JD, Purser JL, Pieper CF, Cohen HJ. Plasma hypertonicity: another marker of frailty? *J Am Geriatr Soc*. 2004;52(8):1313-1320.
19. Rockwood K, Mitnitski A. Frailty in relation to the accumulation of deficits. *J Gerontol A Biol Sci Med Sci*. 2007;62(7):722-727.
20. Pugh JA, Wang CP, Espinoza SE, et al. Influence of frailty-related diagnoses, high-risk prescribing in elderly adults, and primary care use on readmissions in fewer than 30 days for veterans aged 65 and older. *J Am Geriatr Soc*. 2014;62(2):291-298.
21. Scott IA, Hilmer SN, Reeve E, et al. Reducing inappropriate polypharmacy: the process of deprescribing. *JAMA Intern Med*. 2015;175(5):827-834.
22. Mekonnen AB, McLachlan AJ, Brien JA. Pharmacy-led medication reconciliation programmes at hospital transitions: a systematic review and meta-analysis. *J Clin Pharm Ther*. 2016;41(2):128-144.
23. Kaboli PJ, Hoth AB, McClimon BJ, Schnipper JL. Clinical pharmacists and inpatient medical care: a systematic review. *Arch Int Med*. 2006;166(9):955-964.
24. Gleason KM, McDaniel MR, Feinglass J, et al. Results of the Medications at Transitions and Clinical Handoffs (MATCH) study: an analysis of medication reconciliation errors and risk factors at hospital admission. *J Gen Intern Med*. 2010;25(5):441-447.
25. Haynes KT, Oberne A, Cawthon C, Kripalani S. Pharmacists' recommendations to improve care transitions. *Ann Pharmacother*. 2012;46(9):1152-1159.
26. Khanna R, Wachsberg K, Marouni A, Feinglass J, Williams MV, Wayne DB. The association between night or weekend admission and hospitalization-relevant patient outcomes. *J Hosp Med*. 2011;6(1):10-14.