

ALICE KIM, MD

Department of Infective Diseases,
Cleveland Clinic

THOMAS KEYS, MD*

Department of Infective Diseases,
Cleveland Clinic

Infective endocarditis prophylaxis before dental procedures: New guidelines spark controversy

MANY FEWER PEOPLE will need to receive antibiotics as prophylaxis against infective endocarditis before undergoing dental procedures, according to new guidelines released by the American Heart Association.¹ Now, the only patients to receive antibiotics will be those at highest risk, ie, those with a prosthetic heart valve, a history of endocarditis, certain forms of congenital heart disease, or valvulopathy after heart transplantation, and only before certain dental procedures.

Unfortunately, these guidelines are still based largely on expert opinion, with very little hard evidence to show that antibiotic therapy actually prevents infective endocarditis. Nevertheless, the new guidelines appear reasonable, and we believe they should be followed.

■ A RARE BUT LIFE-THREATENING INFECTION

Infective endocarditis is a rare but life-threatening infection, with an incidence in the United States of 10,000 to 20,000 new cases per year. Mortality rates for both native-valve endocarditis and prosthetic-valve endocarditis range from 20% to 30%.^{2,3} For the past half-century, antibiotic prophylaxis for dental procedures has been recommended for patients judged to be at risk of infective endocarditis, in hopes of preventing this dreaded infectious disease.

*Dr. Keys has indicated that he has received royalties from Professional Communications, Inc., for editing a booklet on antimicrobial guidelines.

■ ENDOCARDIAL INJURY, THEN BACTERIAL SEEDING

A combination of events must occur to cause infective endocarditis. First, injury to the endocardial surface induces focal adherence of platelets and fibrin. Then, a bacteremic event seeds this aggregate with microorganisms, attracting more platelets and fibrin, allowing uninhibited microbial growth and the development of an inflammatory plaque or vegetation.

The magnitude and duration of bacteremia that produces this cascade of events is uncertain. Transient bacteremia occurs commonly, not only during procedures that cause trauma to mucosal surfaces or tissue but also with daily activities such as brushing teeth and chewing. The reported incidence of bacteremia during dental intervention ranges from 10% to 100%, and with daily brushing and flossing, from 20% to 68%.¹

■ STAPHYLOCOCCI OVERTAKING VIRIDANS STREPTOCOCCI AS CAUSE

While historically the viridans group of streptococci has been responsible for the largest percentage of cases of both native-valve endocarditis and late-onset prosthetic-valve endocarditis, times have changed. In more recently reported series, *Staphylococcus aureus* appears more common, and unlikely to be susceptible to antibiotics recommended for dental prophylaxis. Other causative pathogens include coagulase-negative staphylococci,

Under the new guidelines, only those at highest risk will get prophylactic antibiotics

TABLE 1

Patients for whom dental prophylaxis is recommended

Only those at highest risk, ie, those with:

A prosthetic cardiac valve

A history of infective endocarditis

Certain forms of congenital heart disease

Unrepaired cyanotic congenital heart disease, including palliative shunts and conduits

Completely repaired congenital heart defect with prosthetic material or device during the first 6 months after the procedure

Repaired congenital heart disease with residual defects at the site or adjacent to the site of prosthetic patch or prosthetic device

Valvulopathy after cardiac transplantation

WILSON W, TAUBERT KA, GEWITZ M, ET AL. PREVENTION OF INFECTIVE ENDOCARDITIS: GUIDELINES FROM THE AMERICAN HEART ASSOCIATION RHEUMATIC FEVER, ENDOCARDITIS, AND KAWASAKI DISEASE COMMITTEE, COUNCIL ON CARDIOVASCULAR DISEASE IN THE YOUNG, AND THE COUNCIL ON CLINICAL CARDIOLOGY, COUNCIL ON CARDIOVASCULAR SURGERY AND ANESTHESIA, AND THE QUALITY OF CARE AND OUTCOMES RESEARCH INTERDISCIPLINARY WORKING GROUP. CIRCULATION 2007; 116:1736–1754. REPRINTED WITH PERMISSION FROM LIPPINCOTT, WILLIAMS, AND WILKINS. ALL RIGHTS RESERVED.

enterococci, gram-negative microorganisms, and fungi.

■ PREVIOUS GUIDELINES—1997

Previous American Heart Association guidelines⁴ separated patients into three risk categories for infective endocarditis. High-risk patients were those with prosthetic heart valves, a history of infective endocarditis, complex cyanotic congenital heart disease, or surgically constructed systemic pulmonary shunts. Moderate-risk patients had other congenital cardiac defects, hypertrophic cardiomyopathy, or acquired valvular heart disease including mitral valve prolapse with regurgitation. Negligible-risk patients—ie, most patients—included those with coronary artery bypass grafts, a permanent pacemaker, or mitral valve prolapse without regurgitation. Antibiotic prophylaxis was recommended only for patients in the high-risk and moderate-risk groups.

■ THOUGHTS AND CHALLENGES

Although prophylaxis has been a standard practice for years, its efficacy and cost-effectiveness have never been proven, owing to a

lack of prospective randomized controlled trials. A sequential relationship between dental procedures and infective endocarditis can be demonstrated in only 4% to 7.5% of cases.⁵ Most cases of infective endocarditis are not preceded by dental procedures.

Furthermore, the data are limited and insufficient to substantiate the efficacy of antibiotics in preventing endocarditis in patients with high-risk cardiac conditions who undergo dental procedures.⁶ Failures have occurred even when the infecting microorganism was susceptible to the antibiotic given for prophylaxis. Since bacteremia occurs also during brushing and flossing of teeth, why give prophylaxis just for dental procedures? Moreover, the risks of causing adverse or anaphylactic reactions from antibiotics, as well as contributing to the nationwide antibiotic resistance problem, are issues not to be taken lightly.

Poor compliance with prophylaxis has been documented. Studies by Duval et al⁷ and others have shown that practitioners adhere to recommended dental prophylaxis programs only about 40% of the time, while only 22% of patients with predisposing cardiac conditions could recall taking their prescribed prophylactic antibiotics before an indicated procedure, as recommended.⁸

■ NEW GUIDELINES—2007

To address many of these concerns, the American Heart Association¹ released extensively revised guidelines in 2007. They are more pragmatic, narrowly focused for a selected group of patients who have a greater lifetime risk of illness and death from infective endocarditis.

The experts who wrote the guidelines agreed that evidence remains poor about which dental procedures increase the risk of infective endocarditis and the efficacy of antibiotic prophylaxis to prevent its development. They stress the importance of good oral hygiene and prevention of dental disease and argue persuasively that this will have a greater impact on decreasing the lifetime risk of infective endocarditis than will antibiotic prophylaxis.

Prophylaxis is now recommended only for patients with a prosthetic heart valve, a history of infective endocarditis, certain forms of

Although prophylaxis has been a standard practice for years, its efficacy and cost-effectiveness have never been proven

TABLE 2

Antibiotic prophylactic regimens

SITUATION	AGENT	ADULTS	CHILDREN
Can take oral medication	Amoxicillin	2 g	50 mg/kg
Cannot take oral medication	Ampicillin	2 g IM or IV	50 mg/kg IM or IV
	OR Cefazolin or ceftriaxone	1 g IM or IV	50 mg/kg IM or IV
Allergic to penicillins or ampicillin, can take oral medication	Cephalexin ^{a,b}	2 g	50 mg/kg
	OR Clindamycin	600 mg	20 mg/kg
	OR Azithromycin or clarithromycin	500 mg	15 mg/kg
Allergic to penicillins or ampicillin, cannot take oral medication	Cefazolin or ceftriaxone ^b	1 g IM or IV	50 mg/kg IM or IV
	OR Clindamycin	600 mg IM or IV	20 mg/kg IM or IV

IM = intramuscularly, IV = intravenously; all regimens given as a single dose 30 to 60 minutes before the procedure.

^aOr another first-generation or second-generation oral cephalosporin in an equivalent adult or pediatric dose.

^bCephalosporins should not be used in a patient with a history of anaphylaxis, angioedema, or urticaria with penicillins or ampicillin.

WILSON W, TAUBERT KA, GEWITZ M, ET AL. PREVENTION OF INFECTIVE ENDOCARDITIS: GUIDELINES FROM THE AMERICAN HEART ASSOCIATION RHEUMATIC FEVER, ENDOCARDITIS, AND KAWASAKI DISEASE COMMITTEE, COUNCIL ON CARDIOVASCULAR DISEASE IN THE YOUNG, AND THE COUNCIL ON CLINICAL CARDIOLOGY, COUNCIL ON CARDIOVASCULAR SURGERY AND ANESTHESIA, AND THE QUALITY OF CARE AND OUTCOMES RESEARCH INTERDISCIPLINARY WORKING GROUP. CIRCULATION 2007; 116:1736–1754. REPRINTED WITH PERMISSION FROM LIPPINCOTT, WILLIAMS, AND WILKINS. ALL RIGHTS RESERVED.

congenital heart disease, and valvulopathy after cardiac transplantation (TABLE 1), and only before procedures that involve manipulation of gingival tissue or the periapical region of teeth, or perforation of the oral mucosa. Excluded are routine dental cleaning and anesthetic injections through noninfected tissue, dental radiography, placement and adjustment of appliances, shedding of deciduous teeth, and bleeding from trauma to the lips.

Regimens for dental prophylaxis should always be given 30 to 60 minutes before the procedure. Oral amoxicillin remains the drug of choice, and, for patients unable to take oral medications, ampicillin, cefazolin, or ceftriaxone can be used, either intramuscularly or intravenously (TABLE 2). For penicillin-allergic patients, oral cephalexin, clindamycin, azithromycin, or clarithromycin is recommended, and, for patients unable to take oral antibiotics, cefazolin, ceftriaxone, or clindamycin, intramuscularly or intravenously.

■ CONTROVERSY WILL CONTINUE

The new guidelines for dental prophylaxis have been extensively revised and simplified. They are now focused only on patients who have a greater lifetime risk of illness and death from infective endocarditis. But what about patients who had previously been advised to take prophylaxis, such as those with mitral valve prolapse with regurgitation, who will not receive prophylaxis any more?

These guidelines will likely stir emotions, not only for practitioners who have strong desires to practice preventive medicine, but also for patients who have been taking prophylaxis in good faith per previous guidelines. They may feel abandoned. Unfortunately, funding for a prospective randomized clinical trial large enough to prove that antibiotic prophylaxis for dental procedures benefits patients is unlikely. That leaves us with the current recommendations, which are based on

Controversy will continue, but we think the new guidelines should be followed



2008

MARCH

FEMALE PELVIC FLOOR DISORDERS
March 7-10, 2008
Hyatt Regency Pier Sixty-Six
Fort Lauderdale, FL

DIAGNOSTIC AND THERAPEUTIC UPDATE: NEUROMUSCULAR DISORDERS, EPILEPSY, AND SLEEP DISORDERS
March 9-15, 2008
Cleveland Clinic
Cleveland, OH

PALLIATIVE MEDICINE 2008
March 13-15, 2008
The Westin-Kierland Resort & Spa
Scottsdale, AZ

AMBULATORY UROLOGY SYMPOSIUM
March 13-15, 2008
Cleveland Clinic
Cleveland, OH

TRIAL, MECHANISMS, AND NEW AGENTS IN B-CELL-DIRECTED THERAPIES
March 18 and 19, 2008
For locations, see www.clevelandclinicmeded.com

APRIL

PROSTATE CANCER UPDATE
April 2, 2008
Greenbriar Party and Conference Centre
Wooster, OH

CONTEMPORARY ISSUES IN PITUITARY DISEASE: CASE-BASED MANAGEMENT UPDATE
April 11, 2008
Cleveland Clinic
Cleveland, OH

ESOPHAGEAL SUMMIT
April 17-18, 2008
Cleveland Clinic
Cleveland, OH

MAY

A HANDS-ON APPROACH TO DIAGNOSTIC PATHOLOGY
May 17-18, 2008
Cleveland Clinic
Cleveland, OH

HIV
May 29-30, 2008
Cleveland Clinic
Cleveland, OH

JUNE

INTENSIVE REVIEW OF INTERNAL MEDICINE
June 1-6, 2008
Cleveland Clinic
Cleveland, OH

HEART-BRAIN SUMMIT
June 4-5, 2008
Cleveland Clinic
Cleveland, OH

ANNUAL RESEARCH, RESIDENTS, AND ALUMNI MEETING
June 12-13, 2008
Cleveland Clinic
Cleveland, OH

AUGUST

INTENSIVE REVIEW OF CARDIOLOGY
August 17-21, 2008
Cleveland Clinic
Cleveland, OH

SEPTEMBER

COLORECTAL CANCER SUMMIT
September 3-4, 2008
Cleveland Clinic
Cleveland, OH

MEETING OF COLLABORATIVE GROUP OF AMERICAS ON INHERITED COLORECTAL CANCER
September 5-6, 2008
Cleveland Clinic
Cleveland, OH

OBESITY SUMMIT
September 10-12, 2008
Cleveland Clinic
Cleveland, OH

FOR SCHEDULE UPDATES AND TO REGISTER ONLINE, VISIT: WWW.CLEVELANDCLINICMEDED.COM

scientific evidence that currently exists and on expert opinion.

The intention of the guidelines is laudable. Of course, there will continue to be controversies with the new rules. Nevertheless, we believe they should be followed until there is more persuasive evidence to the contrary.

REFERENCES

- 1. Wilson W, Taubert KA, Gewitz M, et al. Prevention of infective endocarditis: guidelines from the American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee, Council on Cardiovascular Disease in the Young, and the Council on Clinical Cardiology, Council on Cardiovascular Surgery and Anesthesia, and the Quality of Care and Outcomes Research Interdisciplinary Working Group. Circulation 2007; 116:1736-1754.
2. Hill EE, Herligers P, Claus P, Vanderschueren S, Herregods MC, Peetermans WE. Infective endocarditis: changing epidemiology and predictors of 6-month mortality: a prospective cohort study. Eur Heart J 2007; 28:196-203.
3. Wang A, Athan E, Pappas PA, et al; International Collaboration on Endocarditis-Prospective Cohort Study Investigators. Contemporary clinical profile and outcome of prosthetic valve endocarditis. JAMA 2007; 297:1354-1361.
4. Dajani AS, Taubert KA, Wilson W, et al. Prevention of bacterial endocarditis. Recommendations by the American Heart Association. JAMA 1997; 277:1794-1801.
5. Gendron R, Grenier D, Maheu-Robert LF. The oral cavity as a reservoir of bacterial pathogens for focal infections. Microbes Infect 2000; 2:897-906.
6. Strom BL, Abrutyn E, Berlin JA, et al. Dental and cardiac risk factors for infective endocarditis. A population-based, case-control study. Ann Intern Med 1998; 129:761-769.
7. Duval X, Alla F, Hoen B, et al. Estimated risk of endocarditis in adults with predisposing cardiac conditions undergoing dental procedures with or without antibiotic prophylaxis. Clin Infect Dis 2006; 42:e102-e107.
8. van der Meer JT, van Wijk W, Thompson J, Valkenburg HA, Michel MF. Awareness of need and actual use of prophylaxis: lack of patient compliance in the prevention of bacterial endocarditis. J Antimicrob Chemother 1992; 29:187-194.

ADDRESS: Alice Kim, MD, Department of Infectious Diseases, S32, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195.

Visit our web site at http://www.ccmj.org

Contact us by e-mail at ccmj@ccf.org