

THE CLINICAL PICTURE

NEZAM ALTOROK, MD

University of Toledo Medical Center,
Toledo, OH

AHMAD ABUGIAZIA, MD

University of Toledo Medical Center,
Toledo, OH

NEHAYA ABU SHANAB, MD

University of Toledo Medical Center,
Toledo, OH

ANITA CONWAY, MD

University of Toledo Medical Center,
Toledo, OH

The Clinical Picture

Genital lesions and acute urinary retention

A 36-YEAR-OLD MAN presents to the emergency department with generalized headache and subjective fevers over the past 3 days and painful vesicular lesions on the scrotum and penis (FIGURE 1). He complains of some sensory loss in the perianal region and in the areas of the scrotum and inner thighs. He has not urinated in the past 48 hours and currently has no urge to void.

He is sexually active. Nothing in his medical and surgical history would appear to contribute to his symptoms. He is not taking any medications.

His temperature is 38.5°C (101.3°F). His bladder is distended on palpation and percussion, and he has patches of vesicular and ulcerative lesions on an erythematous base over the scrotum and penis and the proximal inner thighs, lacking a dermatomal pattern. He also has decreased sensation to touch and pain perianally. Anal sphincter tone, deep tendon reflexes, and motor examination of the lower extremities are normal. The patient also lacks any positive meningeal findings. Bladder catheterization yields 1,250 mL of urine.

Q: What is the next step?

- Urine culture
- Immunofluorescence for chlamydial infection
- Cerebrospinal fluid (CSF) analysis
- Spinal myelography
- Urine toxicologic screen

A: CSF analysis is a vital diagnostic step in the evaluation of patients who present with skin rash around the genital area and acute urinary retention, to rule out acute sacral



FIGURE 1

myeloradiculitis. In this disease, CSF analysis typically shows an increase in lymphocytes and a mild increase in protein. Urinary tract infection, gonorrhea, central spinal cord compression, and drug intoxication are unlikely to have this presentation. Hence, CSF analysis is the correct answer.

The patient undergoes a spinal tap. CSF analysis reveals an elevated white cell count of 85 cells/ μ L (reference range 0–5), with 79% lymphocytes (reference range 60%–70%), a mildly elevated protein concentration at 80 mg/dL (reference range 15–50), and a normal glucose level. Polymerase chain reaction testing of the CSF is negative for herpes simplex virus (HSV).

doi:10.3949/cjcm.78a.10071

Magnetic resonance images of the brain and spinal cord are normal.

Polymerase chain reaction testing of specimens collected from the genital lesions detects HSV type 2 (HSV-2). Serum Venereal Disease Research Laboratory and serum human immunodeficiency virus tests are negative.

Diagnosis: HSV-2 sacral radiculitis (Elsberg syndrome).

■ **THE PATIENT'S COURSE AND TREATMENT**

The patient was initially treated symptomatically by draining the urinary bladder via a Foley catheter, followed by intermittent self-catheterization and topical application of lidocaine cream. He was also given intravenous acyclovir (Zovirax) 10 mg/kg every 8 hours for 2 days, followed by 400 mg orally every 8 hours for 10 days.

By 2 weeks, the skin lesions had resolved, and the patient reported regaining sensation in his anal and sacral areas. He also said he was able to void urine without any difficulty.

■ **ELSBERG SYNDROME**

Elsberg syndrome describes multiple disorders characterized by sacral myeloradiculitis, which can be secondary to viral infection, most commonly HSV type 2.^{1,2} CSF analysis and cultures should be performed. However,

cultures and HSV studies of the CSF are often negative, as in this patient.³ CSF lymphocytosis along with a slightly elevated protein level is typical of this disease. It is the latent HSV reactivation in the sacral sensory ganglia that results in sensory neuronal dysfunction and subsequent loss of bladder sensation along with areflexia.

Elsberg syndrome is self-limiting and has a generally good prognosis, but complications such as necrotizing myelitis have been described.⁴ The combination of acute urinary retention associated with vesicular skin lesions and sensory nerve dysfunction in the sacral nerve root distribution in a young, sexually active patient is a strong clue to the diagnosis of Elsberg syndrome, a serious but treatable disease. ■

■ **REFERENCES**

1. **Caplan LR, Kleeman FJ, Berg S.** Urinary retention probably secondary to herpes genitalis. *N Engl J Med* 1977; 297:920-921.
2. **Eberhardt O, Küker W, Dichgans J, Weller M.** HSV-2 sacral radiculitis (Elsberg syndrome). *Neurology* 2004; 63:758-759.
3. **Sakakibara R, Uchiyama T, Liu Z, et al.** Meningitis-retention syndrome. An unrecognized clinical condition. *J Neurol* 2005; 252:1495-1499.
4. **Koskiniemi ML, Vaeheri A, Manninen V, Nikki P.** Ascending myelitis with high antibody titer to herpes simplex virus in the cerebrospinal fluid. *J Neurol* 1982; 227:187-191.

ADDRESS: *Nezam Torok, MD, University of Toledo Medical Center, 3000 Arlington Avenue, Mailstop 1150, Toledo, OH 43614; e-mail nezam.torok@utoledo.edu.*

By 2 weeks, the lesions had resolved, and perianal sensation and normal voiding had returned

Dear Doctor:

As editors, we'd like you to look into every issue, every page of *the Cleveland Clinic Journal of Medicine*.
We'd like to know...

1 How many issues do you look into?

Here's our goal:

- All Most Half Few

2 How do you read the average issue?

Here's our goal:

- Cover-to-cover
 Most articles
 Selected articles

We put it in writing... please put it in writing for us.
We want to hear from you.

CLEVELAND CLINIC JOURNAL OF MEDICINE
The Cleveland Clinic Foundation
9500 Euclid Avenue, NA32
Cleveland, Ohio 44195

FAX 216.444.9385
E-MAIL ccjm@ccf.org