

THE CLINICAL PICTURE

JORGE A. BRENES-SALAZAR, MD
Department of Medicine, Division of Cardiovascular Diseases, Mayo Clinic, Rochester, MN

MARTHA GROGAN, MD
Department of Medicine, Division of Cardiovascular Diseases, Mayo Clinic, Rochester, MN

MARTHA Q. LACY, MD
Department of Medicine, Division of Hematology, Mayo Clinic, Rochester, MN

The Clinical Picture

A 66-year-old woman with an enlarged tongue



FIGURE 1

A 66-YEAR-OLD WOMAN was prompted by her dentist to seek medical attention for an unusually enlarged, smooth-appearing tongue (FIGURE 1). She also complained of fatigue, dyspnea on exertion, and tingling of her hands.

Basic laboratory tests showed normocytic anemia and renal insufficiency. Her thyrotropin level was within normal limits. Serum protein electrophoresis showed a monoclonal M-spike, which prompted a bone marrow biopsy that was diagnostic of multiple myeloma.

Transthoracic echocardiography revealed diffuse hypokinesia with a restrictive filling pattern, myocardial thickening, and moderate mitral and tricuspid regurgitation, highly suggestive of an infiltrative cardiomyopathy. Biopsy of the right ventricle confirmed cardiac amyloidosis of the amyloid immunoglobulin light chain (AL) subtype.

The patient underwent chemotherapy, followed

by autologous stem-cell transplantation. She achieved successful remission, and her cardiomyopathy was compensated.

■ AMYLOIDOSIS IS HETEROGENEOUS

Amyloidosis is a heterogeneous syndrome characterized by abnormal folding of proteins that deposit as insoluble fibrils in different tissues, impairing both structure and function. Virchow was the first to describe amyloid (from *amylon*, Greek for starch) as an abnormal material seen in postmortem examination of the liver. On Congo red staining, the extracellular proteins appear as salmon-red conglomerates, which also show apple-green birefringence under polarized light.

Amyloidosis can be localized but more often represents a systemic process, often associated with a plasma cell dyscrasia such as multiple myeloma.

Modern classification is based on the precursor protein,¹ eg:

- Light chains (AL)
- Acute-phase protein (AA)
- Beta-2-microglobulin (A β 2M)
- Transthyretin (ATTR; occurring in senile systemic amyloidosis)
- Other proteins (occurring in various forms of hereditary systemic amyloidosis).

■ AMYLOIDOSIS AND THE TONGUE

Macroglossia is defined as protrusion of the tongue beyond the alveolar ridge of the teeth at rest. When caused by amyloidosis, it is most often associated with the systemic AL variant and is present in 10% to 23% of patients with this subtype.²

On physical examination, tongue enlargement can present with lateral indentations, with a smooth contour or with nodular deposits. Less often, bullous lesions, vesicles, and ulcers can also be seen, particularly on the

lips. Infiltration of salivary glands can result in xerostomia. Functional symptoms, such as hypogeusia, dysarthria, dysphagia, dysphonia, and, in advanced cases, upper-airway dysfunction can result from restricted mobility of the tongue and tethering to deeper structures.

Surgical management may be necessary if severe obstructive symptoms are present, but infiltrative lesions tend to recur.

■ AMYLOIDOSIS AND THE HEART

Cardiac involvement in amyloidosis is currently the primary determinant of prognosis.³ It is more often seen in the AL, senile, and hereditary forms. It usually manifests as diastolic heart failure, but angina, orthostatic hypotension, dysrhythmias, and syncope can also occur. Systolic dysfunction is typically a late finding in the course of the disease.

Although an electrocardiographic pattern

of low voltage in the precordial and limb leads has been classically associated with cardiac amyloidosis, only 30% of patients with the senile and hereditary forms show this feature.⁴ Left ventricular hypertrophy on electrocardiography is thought to be uncommon, but it has been reported in 16% of patients with AL amyloidosis and biopsy-proven cardiac involvement.⁵ Cardiac troponin levels can be elevated, as seen in other infiltrative cardiomyopathies.³ The diagnosis can be established by endomyocardial biopsy or indirectly by cardiac imaging and a positive extracardiac biopsy.

Drug therapy is supportive and mainly involves diuretics, since angiotensin-converting enzyme inhibitors, beta-blockers, and calcium channel blockers may cause hypotension and exacerbate myocardial dysfunction. The specific treatment varies depending on the underlying cause of amyloidosis. ■

■ REFERENCES

1. **Westermark P, Benson MD, Buxbaum JN, et al.** A primer of amyloid nomenclature. *Amyloid* 2007; 14:179–183.
2. **Kyle RA, Gertz MA.** Primary systemic amyloidosis: clinical and laboratory features in 474 cases. *Semin Hematol* 1995; 32:45–59.
3. **Kapoor P, Thenappan T, Singh E, Kumar S, Greipp PR.** Cardiac amyloidosis: a practical approach to diagnosis and management. *Am J Med* 2011; 124:1006–1015.
4. **Ng B, Connors LH, Davidoff R, Skinner M, Falk RH.** Senile

systemic amyloidosis presenting with heart failure: a comparison with light chain-associated amyloidosis. *Arch Intern Med* 2005; 165:1425–1429.

5. **Murtagh B, Hammill SC, Gertz MA, Kyle RA, Tajik AJ, Grogan M.** Electrocardiographic findings in primary systemic amyloidosis and biopsy-proven cardiac involvement. *Am J Cardiol* 2005; 95:535–537.

ADDRESS: Jorge A. Brenes-Salazar, MD, Mayo Clinic, 200 First St. SW, Rochester, MN 55905;
e-mail: brenessalazar.jorge@mayo.edu

Functional symptoms can include hypogeusia, dysarthria, dysphagia, dysphonia, and, in advanced cases, upper-airway dysfunction



LET US HEAR FROM YOU

- Let us hear your opinions about the *Cleveland Clinic Journal of Medicine*.
- Do you like current articles and sections?
- What topics would you like to see covered and how can we make the *Journal* more useful to you?

PHONE 216.444.2661
FAX 216.444.9385
E-MAIL ccjm@ccf.org
WWW <http://www.ccjm.org>

CLEVELAND CLINIC JOURNAL OF MEDICINE
1950 Richmond Road, TR404
Lyndhurst, Ohio 44124