

LEO SHER, MD

James J. Peters Veterans' Administration Medical Center, and Department of Psychiatry, Mount Sinai School of Medicine, New York, NY

MARÍA DOLORES BRAQUEHAIS, MD, PhD

Department of Psychiatry and Legal Medicine, Vall d'Hebron University Hospital, Universitat Autònoma de Barcelona, CIBERSAM, Spain

MIQUEL CASAS, MD, PhD

Department of Psychiatry and Legal Medicine, Vall d'Hebron University Hospital, Universitat Autònoma de Barcelona, CIBERSAM, Spain

Posttraumatic stress disorder, depression, and suicide in veterans

ABSTRACT

Suicidal behavior is a critical problem in war veterans. Combat veterans are not only more likely to have suicidal ideation, often associated with posttraumatic stress disorder (PTSD) and depression, but they are more likely to act on a suicidal plan. Especially since veterans may be less likely to seek help from a mental health professional, non-mental-health physicians are in a key position to screen for PTSD, depression, and suicidal ideation in these patients. The authors discuss the association of PTSD, depression, and suicide in veterans, keys to assessment of suicide risk, and interventions.

KEY POINTS

The association of suicidal ideation with PTSD and depression and the prevalence of these conditions in combat veterans underline the importance of recognizing and treating these conditions.

In veterans with PTSD related to combat experience, combat-related guilt may be a significant predictor of suicidal ideation and attempts.

Research addressing PTSD, depression, and suicidal behavior in war veterans is critically needed to improve our understanding of the nature of these conditions and how best to treat them.

IN MILITARY VETERANS, depression, post-traumatic stress disorder (PTSD), and suicidal thoughts are common and closely linked. Veterans are less likely to seek care and more likely to act successfully on suicidal thoughts. Therefore, screening, timely diagnosis, and effective intervention are critical.¹

In this article, we review the signs and symptoms of depression and PTSD, the relationship of these conditions to suicidality in veterans, and the role of the non-mental-health clinician in detecting suicidal ideation early and then taking appropriate action. Early identification of suicidality may help save lives of those who otherwise may not seek care.

FROM IDEA TO PLAN TO ACTION

Suicide can be viewed as a process that begins with suicidal ideation, followed by planning and then by a suicidal act,²⁻⁹ and suicidal ideation can be prompted by depression or PTSD.

Suicidal ideation, defined as any thought of being the agent of one's own death,² is relatively common. Most people who attempt suicide report a history of suicidal ideation.¹⁰ In fact, current suicidal ideation increases suicide risk,^{11,12} and death from suicide is especially correlated with the worst previous suicidal ideation.³

Suicidal ideation is an important predictor of suicidal acts in all major psychiatric conditions.^{3,13-17} In a longitudinal study in a community sample, adolescents who had suicidal ideation at age 15 were more likely to have attempted suicide by age 30.⁵

The annual incidence of suicidal ideation in the United States is estimated to be 5.6%,¹⁸

while its estimated lifetime prevalence in Western countries ranges from 2.09% to 18.51%.¹⁹ A national survey found that 13.5% of Americans had suicidal ideation at some point during their lifetime.²⁰ About 34% of people who think about suicide report going from seriously thinking about it to making a plan, and 72% of planners move from a plan to an attempt.²⁰ In the European Study of the Epidemiology of Mental Disorders,²¹ the lifetime prevalence of suicidal ideation was 7.8%, and of suicide attempts 1.3%. Being female, younger, divorced, or widowed was associated with a higher prevalence of suicide ideation and attempts.

Although terms such as “acute suicidal ideation,” “chronic suicidal ideation,” “active suicidal ideation,” and “passive suicidal ideation” are used in the clinical and research literature, the difference between them is not clear. Regardless of the term one uses, any suicidal ideation should be taken very seriously.

HABITUATION IN VETERANS

Interestingly, according to the Interpersonal-Psychological Theory of Suicide,²² the suicidal process is related to feelings that one does not belong with other people, feelings that one is a burden on others or society, and an acquired capability to overcome the fear of pain associated with suicide.²² Veterans are likely to have acquired this capability as the result of military training and combat exposure, which may cause habituation to fear of painful experiences, including suicide.

FEATURES AND CAUSES OF PTSD

PTSD—a severe, multifaceted disorder precipitated by exposure to a psychologically distressing experience—first appeared in the *Diagnostic and Statistical Manual of Psychiatric Disorders* (DSM-III) in 1980,^{23,24} arising from studies of veterans of the Vietnam war and of civilian victims of natural and man-made disasters.^{44,45} However, the study of PTSD dates back more than 100 years. Before 1980, posttraumatic syndromes were recognized by various names, including railway spine, shell shock, traumatic (war) neurosis, concentra-

tion-camp syndrome, and rape-trauma syndrome.^{24,25} The symptoms described in these syndromes overlap considerably with what we now recognize as PTSD.

According to the most recent edition of the *Diagnostic and Statistical Manual*, DSM-IV-TR,²⁷ the basic feature of PTSD is the development of characteristic symptoms following exposure to a stressor event. Examples include:

- Direct personal experience of an event that involves actual or threatened death or serious injury, or other threat to one's physical integrity
- Witnessing an event that involves death, injury, or a threat to the physical integrity of another person
- Learning about unexpected or violent death, serious harm, or threat of death or injury experienced by a family member or other close associate.

People react to the event with fear and helplessness and try to avoid being reminded of it.

Traumatic events leading to PTSD include military combat, violent personal assault, being kidnapped or taken hostage, experiencing a terrorist attack, torture, incarceration, a natural or man-made disaster, or an automobile accident, or being diagnosed with a life-threatening illness.

PTSD is a potentially fatal disorder through suicide. There may be differences in the psychobiology of PTSD and suicidal behavior between war veterans and civilians.²⁸

PTSD often coexists with other psychiatric illnesses^{29,30}: the National Comorbidity Survey found that about 80% of patients with PTSD meet the criteria for at least one other psychiatric disorder.³⁰ Symptoms of PTSD and depression overlap significantly. Common features include diminished interest or participation in significant activities; irritability; sleep disturbance; difficulty concentrating; restricted range of affect; and social detachment.

PTSD also often coexists with traumatic brain injury and other neurologic and medical conditions.^{31,32} The clinician is more often than not faced with a PTSD patient with multiple diagnoses—psychiatric and medical.

Unfortunately, studies show that PTSD often goes unrecognized by non-mental-

Veterans may be habituated to the fear of painful experiences, including suicide

health practitioners.^{31,33} In a national cohort of primary care patients in Israel, 9% met criteria for current PTSD, but only 2% of actual cases were recognized by their treating physician.³³

■ SUICIDE RISK IN VETERANS

Suicidal behavior is a critical problem in war veterans. During the wars in Iraq and Afghanistan, the US Army's suicide rate has increased from 12.4 per 100,000 in 2003 to 18.1 per 100,000 in 2008.³⁴ In the United Kingdom, more veterans have committed suicide since the end of the 1982 Falklands War than the number of servicemen killed in action during the Falklands War.³⁵ The South Atlantic Medal Association, which represents and helps Falklands veterans, believes that 264 veterans had taken their own lives by 2002, a number exceeding the 255 who died in active service. The suicide rate in Falklands War veterans is about three times higher than the rate in those who left the UK armed forces from 1996 to 2005.^{36,37}

Observations have suggested a relatively high prevalence of suicide ideation and attempts in different generations of war veterans and in different countries.³⁸

Suicidal ideation is more dangerous in war veterans than in the general population because they know how to use firearms and they often own them. In other words, they often possess the lethal means to act on their suicidal thoughts.

And female veterans may be more likely to commit suicide with a firearm. A US study³⁹ observed that female veterans who committed suicide were 1.6 times more likely to have used a firearm and male veterans were 1.3 more likely, compared with nonveterans and adjusting for age, marital status, race, and region of residence.

■ DEPRESSION, PTSD, AND SUICIDE RISK

Suicidal ideation in war veterans is often associated with PTSD and depression, conditions that often coexist. And PTSD has been shown to be a risk factor for suicidal ideation in American veterans of the wars in Iraq and Afghanistan.⁴⁰ In a survey of 407 veterans,

those who screened positive for PTSD ($n = 202$) were more than four times as likely to endorse having suicidal ideation compared with veterans who screened negative for PTSD. In veterans who screened positive for PTSD, the risk of suicidal ideation was 5.7 times higher in those with two or more coexisting psychiatric disorders compared with veterans with PTSD alone.⁴⁰

Additional risk factors

Factors contributing to the risk of suicidal ideation and behavior in patients with PTSD include comorbid disorders (especially depression and substance abuse), impulsive behavior, feelings of guilt or shame, re-experiencing symptoms, and prewar traumatic experiences.⁴¹⁻⁴⁵

Recent studies have analyzed factors associated with suicidal ideation in US veterans of the wars in Iraq and Afghanistan. Pietrzak et al⁴⁶ surveyed 272 veterans, of whom 34 (12.5%) reported contemplating suicide in the 2 weeks prior to completing the survey. Screening positive for PTSD and depression and having psychosocial difficulties were associated with suicidal ideation, while postdeployment social support and a sense of purpose and control were negatively associated with it.

Other authors⁴⁷ found that only the “emotional numbing” cluster of PTSD symptoms and the “cognitive-affective” cluster of depression symptoms were distinctively associated with suicidal ideation. Maguen et al⁴⁸ recently reported that 2.8% of newly discharged US soldiers endorsed suicidal ideation. Prior suicide attempts, prior psychiatric medication, and killing in combat were each significantly associated with suicidal ideation, with killing exerting a mediated effect through depression and PTSD symptoms.

Another recent study⁴⁹ suggests that veterans reporting subthreshold PTSD (ie, having symptoms of PTSD but not meeting all the criteria for the diagnosis) were three times more likely to admit to having suicidal ideation compared with veterans without PTSD,⁴⁹ which indicates that subthreshold PTSD may increase suicide risk.

Lemaire and Graham⁵⁰ reported that prior exposure to physical or sexual abuse and having a history of a prior suicide attempt, a current diagnosis of a psychotic disorder, a depressive

PTSD with depression raises the risk of suicidality more than either condition alone

disorder, and PTSD were associated with current suicidal ideation. Other factors related to suicidal ideation were female sex, deployment concerns related to training (a protective factor—ie, it reduces suicide risk by enhancing resilience and by counterbalancing risk factors), the deployment environment, family concerns, postdeployment support (a protective factor), and postdeployment stressors.

PTSD and depression: An additive effect

These findings also suggest that the coexistence of PTSD and depression increases the risk of suicidal ideation more than PTSD or depression alone. This is consistent with the concept of posttraumatic mood disorder, ie, that when these diagnoses coexist, they are different than when they occur alone, and that the coexistence increases the risk of suicidal ideation and behavior.^{51,52}

■ HOW TO ASSESS SUICIDE RISK

Physicians are in a key position to screen for depression and PTSD in all their patients, including those who are veterans.^{31,53}

Traumatic events of adulthood can be asked about directly. For example, “Have you ever been physically attacked or assaulted? Have you ever been in an automobile accident? Have you ever been in a war or a disaster?” A positive response should alert the physician to inquire further about the relationship between the event and any current symptoms.

Traumatic childhood experiences require reassuring statements of normality to put the patient at ease. For example, “Many people continue to think about frightening aspects of their childhood. Do you?”

Physicians working with war veterans suffering from PTSD or depression should regularly inquire about suicidal ideation, and if the patient admits to having suicidal ideation, the physician should ask about the possession of firearms or other lethal means.

This type of screening has limitations. Fear of being socially stigmatized or of appearing weak may prevent veterans from disclosing thoughts of suicide. And one study⁵⁴ found little evidence to suggest that inquiring about suicide successfully identifies veterans most at risk of suicide.

Indirect indicators of suicidality

Identifying indirect indicators of suicidal thoughts is also important: these can include pill-seeking behavior; talking or writing about death, dying, or suicide; hopelessness; rage or uncontrolled anger; seeking revenge; reckless or risky behaviors or activities; feeling trapped; and saying or feeling there is no reason for living.⁵⁵

Other warning signs include depressed mood, anhedonia, insomnia, severe anxiety, and panic attacks.⁵⁶ A prior suicide attempt, a family history of suicidal behavior, and comorbidity of depression and alcoholism are associated with a high suicide risk.^{56–59}

Suicidal behavior is more common after recent, severe, stressful life events and in physical illnesses such as HIV/AIDS, Huntington disease, malignant neoplasm, multiple sclerosis, peptic ulcer, renal disease, spinal cord injury, and systemic lupus erythematosus. This is true in both veterans and nonveterans.⁶⁰

Useful questions

Useful questions in the assessment of suicidal risk can be formulated as follows⁶¹:

- How have you reacted to stress in the past, and how effective are your usual coping strategies?
- Have you contemplated or attempted suicide in the past? If so, how many times and under what circumstances? And how is your current situation compared with past situations when you considered or attempted suicide?
- Do you ever feel hopeless, helpless, powerless, or extremely angry?
- Do you ever have hallucinations or delusions?

The role of guilt

It is important to ask about guilt feelings. Hendin and Haas⁶² observed that in veterans with PTSD related to combat experience, combat-related guilt was the most significant predictor of suicide attempts and of preoccupation with suicide after discharge. Combat veterans may feel guilt about surviving when others have died, acts of omission and commission, and thoughts or feelings.⁶³ Some have suggested that guilt may be a mechanism through which violence is related to PTSD and major depressive disorder in combat veterans.⁶⁴

Suicidal ideation is more dangerous in war veterans than in the general population because they know how to use firearms and they often own them

INTERVENTIONS

Patients with comorbid depression, PTSD, and suicidal ideation are usually very sick and should be referred to a psychiatrist. They are usually treated with antidepressants, such as paroxetine (Paxil) or sertraline (Zoloft), and psychotherapy.⁶⁵ Patients who have a suicidal intent or a plan should be referred to an emergency department for evaluation or hospitalization. All veterans should be given the toll-free phone number of the Veterans Crisis Line (1-800-273-8255), a US Department of Veterans Affairs (VA) resource that connects veterans in crisis and their families and friends, with qualified VA professionals.

As with many illnesses, such as cancer, suicidal behavior is most treatable and yields the best outcome when diagnosed and treated early.⁶⁶ And the earliest manifestation of suicidal behavior is suicidal ideation.

The association of suicidal ideation with PTSD and depression underlines the importance of the timely diagnosis and effective treatment of these conditions among war veterans. Veterans experiencing subthreshold PTSD or depression may be less likely to receive mental health treatment. This indicates that non-mental-health clinicians should be educated about how to detect PTSD and depression symptoms. They may also help to detect suicidality early, which may help save lives.

Promoting social, emotional, and spiritual wellness

Our patients remind us every day that the work we do matters, that we have much more to learn, and that the more we understand suicidal behavior in veterans, the more we can do to reduce their suffering. We need to promote their social, emotional, and spiritual wellness. Encouraging resilience, optimism, and mental health can protect them from depression, suicidal ideation and behavior. Resilience can be promoted by teaching patients to:

- Build relationships with family members and friends who can provide support
- Think well about themselves and identify their areas of strength
- Invest time and energy in developing new skills
- Challenge negative thoughts; try to find optimistic ways of viewing any situation
- Look after their physical health and exercise regularly
- Get involved in community activities to help counter feelings of isolation
- Ask for assistance and support when they need it.⁶⁷

Our knowledge about what works and what does not work in suicide prevention in veterans is evolving. Research addressing combat-related PTSD, depression, and suicidal behavior in war veterans is critically needed to better understand the nature of these conditions.

**The Veterans
Crisis Line:
1-800-273-8255**

REFERENCES

1. Mann JJ. Searching for triggers of suicidal behavior. *Am J Psychiatry* 2004; 161:395–397.
2. American Psychiatric Association. Practice Guideline For The Assessment and Treatment of Patients with Suicidal Behaviors. Arlington, VA: American Psychiatric Publishing, Inc.; 2003.
3. Beck AT, Brown GK, Steer RA, Dahlsgaard KK, Grisham JR. Suicide ideation at its worst point: a predictor of eventual suicide in psychiatric outpatients. *Suicide Life Threat Behav* 1999; 29:1–9.
4. Beck AT, Steer RA, Kovacs M, Garrison B. Hopelessness and eventual suicide: a 10-year prospective study of patients hospitalized with suicidal ideation. *Am J Psychiatry* 1985; 142:559–563.
5. Reinherz HZ, Tanner JL, Berger SR, Beardslee WR, Fitzmaurice GM. Adolescent suicidal ideation as predictive of psychopathology, suicidal behavior, and compromised functioning at age 30. *Am J Psychiatry* 2006; 163:1226–1232.
6. Vilhjalmsson R, Kristjansdottir G, Sveinbjarnardottir E. Factors associated with suicide ideation in adults. *Soc Psychiatry Psychiatr Epidemiol* 1998; 33:97–103.
7. Miotto P, De Coppi M, Frezza M, Petretto D, Masala C, Preti A. Suicidal ideation and aggressiveness in school-aged youths. *Psychiatry Res* 2003; 120:247–255.
8. De Man AF, Leduc CP. Suicidal ideation in high school students: depression and other correlates. *J Clin Psychol* 1995; 51:173–181.
9. Chioqueta AP, Stiles TC. The relationship between psychological buffers, hopelessness, and suicidal ideation: identification of protective factors. *Crisis* 2007; 28:67–73.
10. Hatcher-Kay C, King CA. Depression and suicide. *Pediatr Rev* 2003; 24:363–371.
11. Brown GK, Beck AT, Steer RA, Grisham JR. Risk factors for suicide in psychiatric outpatients: a 20-year prospective study. *J Consult Clin Psychol* 2000; 68:371–377.
12. Fawcett J, Scheftner WA, Fogg L, et al. Time-related predictors of suicide in major affective disorder. *Am J Psychiatry* 1990; 147:1189–1194.
13. Bulik CM, Carpenter LL, Kupfer DJ, Frank E. Features associated with suicide attempts in recurrent major depression. *J Affect Disord* 1990; 18:29–37.
14. Drake RE, Gates C, Cotton PG, Whitaker A. Suicide among schizophrenics. Who is at risk? *J Nerv Ment Dis* 1984; 172:613–617.
15. Oquendo MA, Galfalvy H, Russo S, et al. Prospective study of clinical predictors of suicidal acts after a major depressive episode in patients with major depressive disorder or bipolar disorder. *Am J Psychiatry* 2004; 161:1433–1441.
16. Mann JJ, Ellis SP, Waternaux CM, et al. Classification trees distinguish suicide attempters in major psychiatric disorders: a model of clinical decision making. *J Clin Psychiatry* 2008; 69:23–31.
17. Galfalvy HC, Oquendo MA, Mann JJ. Evaluation of clinical prognostic models for suicide attempts after a major depressive episode. *Acta Psychiatr*

- Scand 2008; 117:244–252.
18. Crosby AE, Cheltenham MP, Sacks JJ. Incidence of suicidal ideation and behavior in the United States, 1994. *Suicide Life Threat Behav* 1999; 29:131–140.
19. Weissman MM, Bland RC, Canino GJ, et al. Prevalence of suicide ideation and suicide attempts in nine countries. *Psychol Med* 1999; 29:9–17.
20. Kessler RC, Borges G, Walters EE. Prevalence of and risk factors for lifetime suicide attempts in the National Comorbidity Survey. *Arch Gen Psychiatry* 1999; 56:617–626.
21. Bernal M, Haro JM, Bernert S, et al; ESEMED/MHEDEA Investigators. Risk factors for suicidality in Europe: results from the ESEMED study. *J Affect Disord* 2007; 101:27–34.
22. Selby EA, Anestis MD, Bender TW, et al. Overcoming the fear of lethal injury: evaluating suicidal behavior in the military through the lens of the Interpersonal-Psychological Theory of Suicide. *Clin Psychol Rev* 2010; 30:298–307.
23. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 3rd ed. Washington, DC: American Psychiatric Association; 1980:236–238.
24. Schnurr PP, Friedman MJ, Bernardy NC. Research on posttraumatic stress disorder: epidemiology, pathophysiology, and assessment. *J Clin Psychol* 2002; 58:877–889.
25. Saigh PA, Bremner JD. The history of posttraumatic stress disorder. In: Saigh PA, Bremner JD, eds. *Posttraumatic Stress Disorder. A Comprehensive Text*. Boston, MA: Allyn & Bacon; 1999:1–17.
26. Hageman I, Andersen HS, Jorgensen MB. Post-traumatic stress disorder: a review of psychobiology and pharmacotherapy. *Acta Psychiatr Scand* 2001; 104:411–422.
27. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. Text Revision. 4th ed. Washington, DC: American Psychiatric Association; 2000:463–468.
28. Sher L, Yehuda R. Preventing suicide among returning combat veterans: a moral imperative. *Mil Med* 2011; 176:601–602.
29. Davidson JR, Hughes D, Blazer DG, George LK. Post-traumatic stress disorder in the community: an epidemiological study. *Psychol Med* 1991; 21:713–721.
30. Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB. Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 1995; 52:1048–1060.
31. Sher L. Recognizing post-traumatic stress disorder. *QJM* 2004; 97:1–5.
32. Kaplan GB, Vasterling JJ, Vedak PC. Brain-derived neurotrophic factor in traumatic brain injury, post-traumatic stress disorder, and their comorbid conditions: role in pathogenesis and treatment. *Behav Pharmacol* 2010; 21:427–437.
33. Taubman-Ben-Ari O, Rabinowitz J, Feldman D, Vaturi R. Post-traumatic stress disorder in primary-care settings: prevalence and physicians' detection. *Psychol Med* 2001; 31:555–560.
34. Tanielian T, Jaycox LH, editors. *Invisible Wounds of War. Psychological and Cognitive Injuries, Their Consequences, and Services to Assist Recovery*. Santa Monica, CA: RAND Corporation; 2008.
35. Spooner MH. Suicide claiming more British Falkland veterans than fighting did. *CMAJ* 2002; 166:1453.
36. Kapur N, While D, Blatchley N, Bray I, Harrison K. Suicide after leaving the UK armed forces—a cohort study. *PLoS Med* 2009; 6:e26.
37. A brief history of the Falklands Islands. Part 7—The 1982 War and Beyond. <http://www.falklands.info/history/history7.html>. Accessed January 5, 2012.
38. Sher L, Vilens A, editors. *War and Suicide*. Hauppauge, New York: Nova Science Publishers; 2009.
39. Kaplan MS, McFarland BH, Huguet N. Firearm suicide among veterans in the general population: findings from the National Violent Death Reporting System. *J Trauma* 2009; 67:503–507.
40. Jakupcak M, Cook J, Imel Z, Fontana A, Rosenheck R, McFall M. Posttraumatic stress disorder as a risk factor for suicidal ideation in Iraq and Afghanistan War veterans. *J Trauma Stress* 2009; 22:303–306.
41. Tarrier N, Gregg L. Suicide risk in civilian PTSD patients—predictors of suicidal ideation, planning and attempts. *Soc Psychiatry Psychiatr Epidemiol* 2004; 39:655–661.
42. Bell JB, Nye EC. Specific symptoms predict suicidal ideation in Vietnam combat veterans with chronic post-traumatic stress disorder. *Mil Med* 2007; 172:1144–1147.
43. Kramer TL, Lindy JD, Green BL, Grace MC, Leonard AC. The comorbidity of post-traumatic stress disorder and suicidality in Vietnam veterans. *Suicide Life Threat Behav* 1994; 24:58–67.
44. Ferrada-Noli M, Asberg M, Ormstad K. Suicidal behavior after severe trauma. Part 2: The association between methods of torture and of suicidal ideation in posttraumatic stress disorder. *J Trauma Stress* 1998; 11:113–124.
45. Tiet QQ, Finney JW, Moos RH. Recent sexual abuse, physical abuse, and suicide attempts among male veterans seeking psychiatric treatment. *Psychiatr Serv* 2006; 57:107–113.
46. Pietrzak RH, Goldstein MB, Malley JC, Rivers AJ, Johnson DC, Southwick SM. Risk and protective factors associated with suicidal ideation in veterans of Operations Enduring Freedom and Iraqi Freedom. *J Affect Disord* 2010; 123:102–107.
47. Guerra VS, Calhoun PS; Mid-Atlantic Mental Illness Research, Education and Clinical Center Workgroup. Examining the relation between posttraumatic stress disorder and suicidal ideation in an OEF/OIF veteran sample. *J Anxiety Disord* 2011; 25:12–18.
48. Maguen S, Luxton DD, Skopp NA, et al. Killing in combat, mental health symptoms, and suicidal ideation in Iraq war veterans. *J Anxiety Disord* 2011; 25:563–567.
49. Jakupcak M, Hoerster KD, Varra A, Vannoy S, Felker B, Hunt S. Hopelessness and suicidal ideation in Iraq and Afghanistan War Veterans reporting sub-threshold and threshold posttraumatic stress disorder. *J Nerv Ment Dis* 2011; 199:272–275.
50. Lemaire CM, Graham DP. Factors associated with suicidal ideation in OEF/OIF veterans. *J Affect Disord* 2011; 130:231–238.
51. Sher L. The concept of post-traumatic mood disorder. *Med Hypotheses* 2005; 65:205–210.
52. Sher L. Suicide in war veterans: the role of comorbidity of PTSD and depression. *Expert Rev Neurother* 2009; 9:921–923.
53. Blank AS Jr. Clinical detection, diagnosis, and differential diagnosis of post-traumatic stress disorder. *Psychiatr Clin North Am* 1994; 17:351–383.
54. Denneson LM, Basham C, Dickinson KC, et al. Suicide risk assessment and content of VA health care contacts before suicide completion by veterans in Oregon. *Psychiatr Serv* 2010; 61:1192–1197.
55. US Department of Veterans Affairs. Mental Health. Suicide Prevention. http://www.mentalhealth.va.gov/suicide_prevention. Accessed December 8, 2011.
56. Gliatto MF, Rai AK. Evaluation and treatment of patients with suicidal ideation. *Am Fam Physician* 1999; 59:1500–1506.
57. Sher L, Oquendo MA, Mann JJ. Risk of suicide in mood disorders. *Clin Neurosci Res* 2001; 1:337–344.
58. Oquendo MA, Currier D, Mann JJ. Prospective studies of suicidal behavior in major depressive and bipolar disorders: what is the evidence for predictive risk factors? *Acta Psychiatr Scand* 2006; 114:151–158.
59. Sher L. Alcoholism and suicidal behavior: a clinical overview. *Acta Psychiatr Scand* 2006; 113:13–22.
60. Moscicki EK. Identification of suicide risk factors using epidemiologic studies. *Psychiatr Clin North Am* 1997; 20:499–517.
61. Goldman HH. *Review of General Psychiatry*, 5th ed. New York, NY: Lange Medical Books/McGraw-Hill; 2000.
62. Hendin H, Haas AP. Suicide and guilt as manifestations of PTSD in Vietnam combat veterans. *Am J Psychiatry* 1991; 148:586–591.
63. Henning KR, Frueh BC. Combat guilt and its relationship to PTSD symptoms. *J Clin Psychol* 1997; 53:801–808.
64. Marx BP, Foley KM, Feinstein BA, Wolf EJ, Kaloupek DG, Keane TM. Combat-related guilt mediates the relations between exposure to combat-related abusive violence and psychiatric diagnoses. *Depress Anxiety* 2010; 27:287–293.
65. Hetrick SE, Purcell R, Garner B, Parslow R. Combined pharmacotherapy and psychological therapies for post traumatic stress disorder (PTSD). *Cochrane Database Syst Rev* 2010; (7):CD007316.
66. Brent DA, Oquendo M, Birmaher B, et al. Familial pathways to early-onset suicide attempt: risk for suicidal behavior in offspring of mood-disordered suicide attempters. *Arch Gen Psychiatry* 2002; 59:801–807.
67. Australian Department of Health and Ageing. Fact sheet 6: Resilience, vulnerability, and suicide prevention. Living is for Everyone (LIFE) fact sheets. www.livingisforeveryone.com.au/LIFE-Fact-sheets.html. Accessed December 8, 2011.

ADDRESS: Leo Sher, MD, James J. Peters Veterans Administration Medical Center, 130 West Kingsbridge Road, Bronx, NY 10468; e-mail: Leo.Sher@mssm.edu.