

## THE CLINICAL PICTURE

**CRISTINA GÓMEZ-FERNÁNDEZ, MD**

Department of Dermatology, Hospital Universitario La Paz, Madrid, Spain

**BEATRIZ CASADO VERRIER, MD**

Department of Dermatology, Hospital Universitario La Paz, Madrid, Spain

**LETICIA SANZ CANALEJAS, MD**

Department of Dermatology, Hospital Universitario La Paz, Madrid, Spain

**RICARDO MORENO ALONSO DE CELADA, MD**

Department of Dermatology, Hospital Universitario La Paz, Madrid, Spain

**PEDRO HERRANZ PINTO, MD**

Department of Dermatology, Hospital Universitario La Paz, Madrid, Spain

# The Clinical Picture

## Cornflake-like scales on the ankles and feet



**FIGURE 1.** Brown, adherent scales on the dorsal aspect of her ankles and feet.

**A**N 81-YEAR-OLD WOMAN presents with slowly growing, asymptomatic lesions on both feet and ankles. She has a long history of neglecting personal hygiene. Physical examination notes patchy brown, waxy scales in a symmetric pattern, localized to the dorsal aspect of the ankles and feet (**FIGURE 1**).

**Q:** On the basis of the skin findings, which test should be ordered to establish a diagnosis?

- Biopsy of the lesions

- Hemoglobin A<sub>1c</sub>  
 Fungal culture  
 Brush with soap and water or alcohol  
 Bacterial culture

**A:** Patchy brown, waxy scales in a patient who neglects normal cleansing should raise the possibility of dermatosis neglecta. Hence, brushing with soap and water or alcohol is the correct answer.

In this patient, the plaques were easily removed using an alcohol-soaked swab, revealing underlying normal skin with bleeding spots. The patient's family was instructed to lightly scrub the area daily with a washcloth, soap, and water. The lesions resolved completely over the next 3 weeks.

### ■ DERMATOSIS NEGLECTA

Dermatosis neglecta is a condition that results from the accumulation in the epidermal stratum corneum of sebum, sweat, corneocytes, and bacterias in areas of unwashed skin. It affects all age groups, and it has a similar prevalence in both sexes. Sometimes, an underlying physical disability or emotional problem is the cause, other times it is due to prior trauma or hyperesthesia of the affected area. Dermatitis neglecta presents with asymptomatic areas of dark brown patches or plaque with cornflake-like scales and a verrucous surface.<sup>1-4</sup> The diagnosis is usually easy and is based on the characteristic findings and the simple test described above. For treatment, in some cases a keratolytic lotion containing lactic acid, glycolic acid, urea, or salicylic acid is helpful.<sup>4</sup>

doi:10.3949/ccjm.77a.10055

■ **MISDIAGNOSIS IS EASY**

Dermatosis neglecta could easily be misdiagnosed as verrucous epidermal nevus, terra firma-forme dermatosis, acanthosis nigricans, or another less common dermatosis of “dirty skin” lesions, such as confluent and reticulate papillomatosis (Gougerot-Carteaud syndrome).<sup>2-4</sup>

Verrucous epidermal nevus appears at birth or early childhood. Patients have verrucous papules and plaques that commonly follow a narrow linear distribution.

Acanthosis nigricans is characterized by broad, velvety, hyperpigmented smooth plaques on the neck, axillae, and flexures. It is related to insulin resistance, obesity, and underlying malignancy.

Confluent and reticulate papillomatosis is a rare dermatosis of unknown cause, with a pigmented reticular pattern distributed on the central trunk. Patients tend to practice normal hygiene.

Terra firma-forme dermatosis is characterized by dirt-like patches that are resistant to usual hygiene measures but that can be eliminated by rubbing with isopropyl alcohol. The cause is unknown, and it is not related to a lack of cleansing. It is considered a variant of dermatosis neglecta.

The medical literature contains few reports of dermatosis neglecta, but the true incidence is probably underestimated. Early recognition by a physician avoids the need for aggressive diagnostic and therapeutic procedures. ■

■ **REFERENCES**

1. Ruiz-Maldonado R, Durán-McKinster C, Tamayo-Sánchez L, Orozco-Covarrubias ML. Dermatitis neglecta: dirt crusts simulating verrucous nevi. *Arch Dermatol* 1999; 135:728-729.
2. Akkash L, Badran D, Al-Omari AQ. Terra firma forme dermatosis. Case series and review of the literature. *J Dtsch Dermatol Ges* 2009; 7:102-107.
3. Qadir SN, Ejaz A, Raza N. Dermatitis neglecta in a case of multiple fractures, shoulder dislocation and radial nerve palsy in a 35-year-old man: a case report. *J Med Case Reports* 2008; 2:347.
4. Lucas JL, Brodell RT, Feldman SR. Dermatitis neglecta: a series of case reports and review of other dirty-appearing dermatoses. *Dermatol Online J* 2006; 12:5.

**ADDRESS:** Cristina Gómez-Fernández, MD, Department of Dermatology, Hospital Universitario La Paz, Paseo de la Castellana 261, 28046 Madrid, Spain; e-mail [crisgomezfernandez@hotmail.com](mailto:crisgomezfernandez@hotmail.com).

# We Welcome Your Letters



WE ENCOURAGE YOU TO WRITE TO US, either to respond to an article published in the *Journal* or to address a clinical issue of importance to you. You may submit letters by mail, fax, or e-mail.

Please be sure to include your full address, phone number, fax number, and e-mail address. Please write concisely, as space is limited. Letters may be edited for style and length. We cannot return materials sent. Submission of a letter constitutes permission for the *Cleveland Clinic Journal of Medicine* to publish it in various editions and forms.

**MAILING ADDRESS**

Letters to the Editor  
*Cleveland Clinic Journal of Medicine*  
9500 Euclid Ave., NA32  
Cleveland, OH 44195  
**FAX** 216.444.9385  
**E-MAIL** [ccjm@ccf.org](mailto:ccjm@ccf.org)

