

Sponsored by and content developed in conjunction with Ortho Women's Health & Urology™, Division of Ortho-McNeil-Janssen Pharmaceuticals, Inc.

SUPPLEMENT TO

OBG
MANAGEMENT

May 2010

Available at obgmanagement.com

Counseling our patients regarding cycle control

Highlights from a randomized clinical trial comparing a 21-day and a 24-day oral contraceptive regimen

Andrew M. Kaunitz, MD

Professor and Associate Chairman

Department of Obstetrics and Gynecology

University of Florida College of Medicine-Jacksonville

Jacksonville, Florida

Catherine L. Dean, MD, MPH

Clinical Assistant Professor

Department of Obstetrics and Gynecology

Washington University School of Medicine

Managing Partner, Women's HealthPartners, Inc.

St. Louis, Missouri



ORTHO WOMEN'S HEALTH & UROLOGY™
DIVISION OF ORTHO-MCNEIL-JANSSEN PHARMACEUTICALS, INC.



Counseling our patients regarding cycle control

Highlights from a randomized clinical trial comparing a 21-day and a 24-day oral contraceptive regimen

KEY POINTS

- Unscheduled bleeding and an absence of scheduled bleeding are common reasons cited by women for discontinuing oral contraceptives (OCs).
- Women taking OCs with 20 mcg of estrogen report more bleeding issues (ie, irregular bleeding and bleeding disturbances) and have higher discontinuation rates than those taking OCs with more estrogen.¹
- Although newer regimens, including 24/4 formulations, have been developed to reduce side effects, unscheduled bleeding remains a primary reason for discontinuation.
- Data have been lacking to support a decrease in unscheduled bleeding with the use of active hormonal treatment for 24 days rather than 21 days.
- In the present 3-cycle trial, users of the 24/4-day drospirenone (DRSP)/ethinyl estradiol (EE) OC experienced more unscheduled bleeding and a greater absence of scheduled bleeding than users of the 21/7-day norgestimate (NGM)/EE OC at each cycle.
- With this study's findings, the clinician can counsel patients using the 24-day regimen studied to expect the following in the first 3 cycles: (1) The duration of withdrawal bleeding may be shorter; (2) There will likely be more unscheduled bleeding; and (3) There will likely be more "absence of withdrawal" bleeding as compared with the studied 21-day regimen.²

FACULTY DISCLOSURES

Dr Kaunitz reports the following:

Consultant and/or speakers bureau, and research support: Bayer HealthCare Pharmaceuticals, Merck & Co., Inc., Ortho Women's Health & Urology™, Division of Ortho-McNeil-Janssen Pharmaceuticals, Inc., and Teva Pharmaceuticals Industries Ltd. Dr Kaunitz has been compensated for his work on this publication.

Dr Dean reports the following:

Speakers bureau: Bayer HealthCare Pharmaceuticals, and Ortho Women's Health & Urology™, Division of Ortho-McNeil-Janssen Pharmaceuticals, Inc. Dr Dean has been compensated for her work on this publication.

BACKGROUND

Unscheduled bleeding (ie, spotting/break-through bleeding) and early, late, or absent withdrawal bleeding are common reasons cited by women for discontinuing use of oral contraceptives (OCs).³⁻⁵ A European survey of nearly 7000 current or past users of OCs found that women who experienced side effects such as unscheduled bleeding in the first 3 months of use were nearly twice as likely to discontinue use as those who reported no cycle control problems during this time.⁴

A recent Cochrane review of bleeding data from trials comparing OCs containing 20 mcg ethinyl estradiol (EE) and >20 mcg EE concluded that rates of early study discontinuation were higher among women taking OCs with 20 mcg EE than among those taking OCs with 30 to 35 mcg EE, reflecting a higher incidence of unscheduled bleeding with the 20 mcg EE formulations.³ These findings have prompted the development of novel regimens and/or OC formulations with a view toward improving bleeding patterns.

One approach employed with the goal of decreasing unscheduled bleeding is to use active hormonal treatment for 24 days rather than 21 days.⁶ In a comparative study of 21- and 24-day regimens of norethindrone acetate (NETA) 1 mg/EE 20 mcg, the pattern of unscheduled bleeding over 7 cycles of use appeared to be similar.⁶ However, in another comparative study, there was more unscheduled bleeding with the 21-day regimen of NETA 1 mg/EE 20 mcg than with a 21-day regimen of triphasic norgestimate (NGM)/EE 25 mcg.⁷

In a multicenter, noncomparative, 13-cycle trial of 24/4 day drospirenone (DRSP)/EE,

a favorable bleeding profile was suggested by the finding that less than 1% of women discontinued the OC due to unscheduled bleeding.⁸ Two comparative trials assessing bleeding patterns of this 24-day OC regimen have been published.^{9,10} One trial comparing the 24-day regimen of DRSP/EE and a 21-day regimen of desogestrel 150 mcg/EE 20 mcg found no significant difference in the number of bleeding/spotting episodes over 7 cycles of use.⁹ In the other trial comparing these 2 OC formulations, the number and duration of bleeding/spotting episodes was lower with the DRSP/EE 24/4-day regimen over 7 cycles of use. However, the incidence of unscheduled bleeding in the 2 groups was similar during cycles 2 to 6.¹⁰ A third trial comparing the 24/4-day regimen of DRSP/EE with an investigational OC formulation containing 2.5 mg nomegestrol acetate and 1.5 mg 17-beta estradiol (E2) was completed in July 2008, but the results have yet to be published.^{11,12} This OC formulation is not currently approved for use in either the United States or abroad.

A “dynamic” 26/2-day dosing regimen was developed for another new OC formulation containing estradiol valerate (E2V) and the progestin dienogest (DNG) (TABLE 1). Although marketed in some European countries, this OC is not currently FDA-approved.

In a 7-cycle, randomized, multicenter, double-blind, double-dummy trial comparing E2V/DNG and a 21/7-day regimen of EE

20 mcg and levonorgestrel 100 mcg (EE/LNG), women randomized to the E2V/DNG OC formulation reported shorter and lighter bleeding than those who received EE/LNG, and fewer bleeding/spotting days for each 90-day reference period.¹³ Intracyclic bleeding with the 2 OC formulations was similar. However, absence of scheduled withdrawal bleeding occurred more often among users of E2V/DNG than users of EE/LNG.¹³

BLEEDING PATTERNS WITH A 21-DAY VS 24-DAY OC REGIMEN—RANDOMIZED CLINICAL TRIAL

All of the aforementioned studies used various definitions of bleeding. The US Food and Drug Administration’s (FDA) Reproductive Health Drugs Advisory Committee (RHDAC) has now endorsed the definitions of measuring bleeding published by Mishell and colleagues.¹⁴⁻¹⁶ A recently published trial comparing bleeding patterns with a 21/7-day NGM/EE 25 mcg OC regimen and a 24/4-day DRSP/EE 20 mcg regimen incorporated these definitions, as well as the recommendation that an interactive voice-response system (IVRS) be used to collect bleeding data.¹⁶

Materials and Methods

This 3-cycle, randomized, open-label, active-controlled study was conducted at 20 centers in the United States. Inclusion and exclusion criteria are summarized in TABLE 2.²

Women in this study were randomized 1:1 to treatment with 21/7-day NGM/EE 25 mcg OC or 24/7-day DRSP/EE 20 mcg OC for three 28-day cycles, with randomization stratified by center and recent hormonal contraceptive exposure (ie, fresh starts vs switchers). Fresh starts were subjects who had not been on a hormonal contraceptive for 60 days or more, and switchers were those who had been on a hormonal contraceptive within the past 60 days.

Bleeding Definitions

Cycle control was evaluated using bleeding (blood loss) data recorded daily using an IVRS diary—one of the real-time methods

TABLE 1
“Dynamic” 26/2-day dosing regimen of estradiol valerate and dienogest¹³

Days	E2V	DNG	Placebo
1-2	3 mg	-	-
3-7	2 mg	2 mg	-
8-24	1 mg	3 mg	-
25-26	1 mg	-	-
27-28	-	-	✓

DNG, dienogest; E2V, estradiol valerate.

TABLE 2**Inclusion and exclusion criteria**

<p>Included:</p> <ul style="list-style-type: none"> • Aged 18 to 45 years (those aged 35 to 45 years must be nonsmokers) • Nonpregnant, nonlactating, sexually active women, seeking oral hormonal contraception • Healthy, based on medical history, physical examination, gynecologic history • Regular menstrual cycles, including at least one normal menstrual period (typical in duration and amount of flow for the subject) within 35 days of first study visit • Negative Chlamydia test (urine or cervical swab) • Pap smear without evidence of moderate to severe dysplasia, or any malignancy within preceding 12 months <p>Excluded:</p> <ul style="list-style-type: none"> • History or presence of disorders commonly accepted as contraindications to steroid hormone therapy • Body mass index >40 kg/m² • Subject previously discontinued triphasic NGM/EE 25 mcg or DRSP/EE 20 mcg due to breakthrough bleeding • Subjects who received depot medroxyprogesterone acetate or other hormonal injectables within 6 months of screening or other hormonal implants in place or removed within 60 days of screening • Used a steroid-containing intrauterine device within 3 months before screening <p>DRSP, drospirenone; EE, ethinyl estradiol; NGM, norgestimate.</p>
--

recommended by Mishell and colleagues.^{2,16} Bleeding was defined according to the RH-DAC-endorsed criteria with one exception—an episode of bleeding/spotting was defined using the Belsey/WHO definition (TABLE 3).¹⁷

Study Endpoints

The primary efficacy endpoint was the participant's total number of unscheduled bleeding days during cycles 1 to 3, as recorded daily using the IVRS diary. Secondary efficacy variables related to bleeding that were analyzed for each of cycles 1 to 3, and cumulatively, included: (1) incidence of unscheduled bleeding, (2) mean number of unscheduled bleeding episodes, (3) incidence of scheduled bleeding, and (4) mean number of scheduled bleeding days.

Consistent with the FDA-endorsed criteria, days 1 through 7 of cycle 1 were excluded from the analysis of bleeding. In addition, only participants who took study medication for at least 7 days and had bleeding data after day 7 were included in the analysis of cycle control.

These measures eliminate biased results related to early dropouts.

Evaluation of Bleeding Patterns

The evaluation of bleeding patterns was based on the intent-to-treat population, defined as all randomized subjects who took the study drug and for whom there was post-baseline bleeding data after day 7.

RESULTS

A total of 334 women took study OCs (167 in each treatment group) and were included in the safety analysis. Demographic and baseline characteristics of the 2 treatment groups are shown in TABLE 4.² Age, racial distribution, and previous contraceptive status were comparable in the 2 groups; however, mean weight and body mass index were greater in the 24/4-day DRSP/EE group (TABLE 4).

Bleeding Patterns

Efficacy data were evaluable for 165 of 167 participants in the 21/7-day NGM/EE group and all 167 participants in the 24/4-day DRSP/EE group.²

TABLE 3

Terminology and definitions for bleeding variables^{2,15,17}

Term	Definition
Bleeding	Evidence of blood loss that requires the use of sanitary protection with a tampon, pad, or pantyliner ¹⁵
Spotting	Evidence of minimal blood loss that does not require new use of any type of sanitary protection, including pantyliners ¹⁵
Episode of bleeding/spotting	One or more consecutive days during which blood loss (bleeding or spotting) has been recorded, each episode bounded by bleeding/spotting-free days ¹⁷
Scheduled bleeding	Any blood loss that occurs during a hormone-free interval, regardless of the duration of the regimen, and which may continue into days 1-4 of the next cycle of oral contraceptive pill therapy ¹⁵
Unscheduled bleeding	Any blood loss that occurs while taking active hormones, regardless of the duration of the regimen, with the exception of blood loss that begins during the hormone-free interval and continues through days 1-4 of the subsequent active cycle, and blood loss reported during days 1-7 of the first cycle of oral contraceptive pill therapy ¹⁵
Absence of all bleeding and spotting	No bleeding or spotting; replaces the term “amenorrhea” ¹⁵

Reprinted with permission from Kaunitz AM, Burkman RT, Fisher AC, et al. Cycle control with a 21-day compared with a 24-day oral contraceptive pill: a randomized controlled trial. *Obstet Gynecol.* 2009;114(6):1205-1212. Copyright © 2009 Lippincott Williams & Wilkins.

TABLE 4

Demographic and baseline characteristics

	21/7-day NGM/EE (n = 167)	24/4-day DRSP/EE (n = 167)
Age (y)	27.4 (±6.80)	27.3 (±6.59)
Race	128 (76.6)	126 (75.4)
White	20 (12.0)	28 (16.8)
Black or African	8 (4.8)	4 (2.4)
Asian	11 (6.6)	9 (5.4)
Other		
Weight, lb	147.6 (±30.10)	156.7(±35.38) ^a
Body mass index, kg/m ²	24.7 (±4.48)	26.6 (±5.47) ^b
Smoking status		
Smoker	11 (6.6)	22 (13.2)
Nonsmoker	156 (93.4)	145 (86.8)
Previous HC status		
Fresh start ^c	72 (43.1)	74 (44.3)
Switcher ^d	95 (56.9)	93 (55.7)

DRSP/EE, drospirenone/ethinyl estradiol; HC, hormonal contraceptive; NGM/EE, norgestimate/ethinyl estradiol. Data are mean (± standard deviation) or n (%). Statistically significant difference between groups: ^a*P* = 0.014; ^b*P* < 0.001. ^cNo HC within 60 days of study drug start. ^dUsed HC within 60 days of study drug start.

Reprinted with permission from Kaunitz AM, Burkman RT, Fisher AC, et al. Cycle control with a 21-day compared with a 24-day oral contraceptive pill: a randomized controlled trial. *Obstet Gynecol.* 2009;114(6):1205-1212. Copyright © 2009 Lippincott Williams & Wilkins.

Unscheduled bleeding

The mean number of days of unscheduled bleeding in cycles 2 and 3 and cumulatively was significantly less in the 21/7 NGM/EE group as compared with the 24/4 DRSP/EE group (**FIGURE 1**).² Over the 3 cycles, participants using the 21/7-day NGM/EE OC experienced approximately 1.5 fewer days of unscheduled bleeding than users of the 24/4-day DRSP/EE OC.

Across all 3 cycles, 55 (33.3%) participants in the 21/7-day NGM/EE group and 29 (17.4%) participants in the 24/4-day DRSP/EE group did not experience any unscheduled bleeding episodes (**FIGURE 2**).² There were significantly fewer unscheduled bleeding episodes in the 21/7-day NGM/EE group than in the 24/4-day DRSP/EE group (mean 1.47 vs 2.01; *P* = 0.001).

A post hoc analysis controlling separately for weight and smoking found no difference from the primary endpoint finding.²

Absence of scheduled bleeding

The percentage of patients reporting the absence of scheduled bleeding was significant-

ly lower in the 21/7-day NGM/EE group than in the 24/4-day DRSP/EE group in each cycle ($P < 0.001$ for each) (FIGURE 3).²

SAFETY AND TOLERABILITY

Safety was evaluated in all participants who took the study medication and provided post-baseline safety data after day 7 (N = 334; 167 in each study group). Treatment-emergent adverse events were reported by 34 of 167 participants in each group.

Serious adverse events reported (not considered related to study drug) were:


- Norgestimate/ethinyl estradiol: 2 subjects
 - Lumbar disc protrusion (1)
 - Uterine leiomyoma (fibroid) with ruptured ovarian cyst and pelvic hemorrhage (1)

Adverse events that led to discontinuation (possibly related to treatment):

- Norgestimate/ethinyl estradiol:
 - Nausea/vomiting
- Drospirenone/ethinyl estradiol:
 - Pulmonary embolism
 - Irritability
 - Mood swings

FIGURE 1

Mean number of days of unscheduled bleeding*



Other adverse events:

- Norgestimate/ethinyl estradiol:
 - Pregnancy

DISCUSSION

This study found that over the first 3 cycles of use, there was less unscheduled bleeding

FIGURE 2

Mean number of unscheduled bleeding episodes





FIGURE 3
Participants with absence of scheduled bleeding



among users of the 21/7-day NGM/EE OC compared with users of the 24/4-day DRSP/EE OC. This high incidence of absence of scheduled bleeding among participants in the 24/4-day DRSP/EE group was not expected. The greater number of scheduled bleeding days in the 21/7-day NGM/EE group than in the 24/4-day DRSP/EE group was expected based on the longer hormone-free interval. However, among those trial participants who reported scheduled bleeding, the duration of scheduled bleeding was similar for the 2 OC formulations.

Women using a cyclical OC regimen expect to experience bleeding during the hor-

mone-free interval, but do not expect to bleed while taking active hormones. Although many women can readily adjust to the experience of bleeding during the first few days of their next pill pack, this issue needs to be discussed when counseling women who are initiating therapy.

OC research and development continues to investigate novel dosing regimens that alter the length of the hormone-free period. Additionally, newer estrogen components are being evaluated as alternatives to EE. Consistent with past research, the goal for any novel OC formulation is that it will not only demonstrate contraceptive efficacy, but also improve on the safety and efficacy profiles of existing OCs. By meeting these goals, new OCs will afford clinicians the opportunity to offer, and women the option to select, OCs that meet their expectations and promote correct and consistent use.

CONCLUSIONS

In this 3-cycle study using FDA-recommended criteria to evaluate bleeding patterns, users of the 21/7-day NGM/EE OC experienced less unscheduled bleeding and more scheduled bleeding compared with users of the 24/4-day DRSP/EE OC. These findings can be a helpful guide for the clinician when counseling patients on what to expect when starting a 21-day or 24-day OC regimen.

INDICATION

ORTHO TRI-CYCLEN® LO (norgestimate/ethinyl estradiol) is indicated for the prevention of pregnancy in women who elect to use oral contraception as their method of contraception.

IMPORTANT SAFETY INFORMATION

Serious as well as minor side effects have been reported with the use of oral contraceptives. Serious risks include blood clots, stroke and

heart attacks. Cigarette smoking increases the risk of serious cardiovascular side effects, especially in women over 35. Women who use oral contraceptives are strongly advised not to smoke. The Pill does not protect against HIV or sexually transmitted diseases.

Please see full Prescribing Information, including Boxed WARNING in the back of the supplement.

References

1. Gallo MF, Nanda K, Grimes DA, et al. 20 µg versus >20 µg estrogen combined oral contraceptives for contraception. *Cochrane Database Syst Rev.* 2008;(4)CD003989.
2. Kaunitz AM, Burkman RT, Fisher AC, et al. Cycle control with a 21-day compared with a 24-day oral contraceptive pill: a randomized controlled trial. *Obstet Gynecol.* 2009;114:1205-1212.
3. Sabatini R, Cagiano R. Comparison profiles of cycle control, side effects and sexual satisfaction of three hormonal contraceptives. *Contraception.* 2006;74:220-223.
4. Rosenberg MJ, Waugh MS, Meehan TE. Use and misuse of oral contraceptives: risk indicators for poor pill taking and discontinuation. *Contraception.* 1995;51:283-288.
5. Rosenberg MJ, Waugh MS. Oral contraceptive discontinuation. A prospective evaluation of frequency and reasons. *Am J Obstet Gynecol.* 1998;179(3 pt 1):577-582.
6. Nakajima ST, Archer DF, Ellman H. Efficacy and safety of a new 24-day oral contraceptive regimen of norethindrone acetate 1 mg/ethinyl estradiol 20 µg (Loestrin® 24 Fe). *Contraception.* 2007;75:16-22.
7. Hampton RM, Short M, Bieber E, et al. Comparison of a novel norgestimate/ethinyl estradiol oral contraceptive (Ortho Tri-Cyclen Lo) with the oral contraceptive Loestrin Fe 1/20. *Contraception.* 2001;63:289-295.
8. Bachmann G, Sulak PJ, Sampson-Landers C, et al. Efficacy and safety of a low-dose 24-day combined oral contraceptive containing 20 µg ethinylestradiol and 3 mg drospirenone. *Contraception.* 2004;70:191-198.
9. Klipping C, Marr J. Effects of two combined oral contraceptives containing ethinyl estradiol 20 µg combined with either drospirenone or desogestrel on lipids, hemostatic parameters and carbohydrate metabolism. *Contraception.* 2005;71:409-416.
10. Anttila L, Kunz M, Marr J. Bleeding pattern with drospirenone 3 mg+ethinyl estradiol 20 mcg 24/4 combined oral contraceptive compared with desogestrel 150 mcg+ethinyl estradiol 20 mcg 21/7 combined oral contraceptive. *Contraception.* 2009;80:445-451.
11. Patent application: Oral contraceptive regimen [monophasic 1.5 mg 17-beta-estradiol and 2.5 mg nomegestrol for 24 days followed by a hormone-free period of 4 days]. <http://www.faqs.org/patents/app/20080242650>. Accessed February 15, 2010.
12. Efficacy and safety study of the combined oral contraceptive NOMAC-E2 compared to a COC containing DRSP/EE (292002) (Completed)(PO5722). <http://clinicaltrials.gov/ct2/show/NCT00413062>. Accessed December 7, 2009.
13. Ahrendt HJ, Makalova D, Parke S, et al. Bleeding pattern and cycle control with an estradiol-based oral contraceptive: a seven-cycle, randomized comparative trial of estradiol valerate/dienogest and ethinyl estradiol/levonorgestrel. *Contraception.* 2009;80:436-444.
14. Reproductive Health Drugs Advisory Committee. US Food and Drug Administration Center for Drug Evaluation and Research. Vol I and II. US Food and Drug Administration, January 23-24, 2007. <http://www.fda.gov/ohrms/dockets/ac/cder07.htm>.
15. Mishell DR Jr, Guillebaud J, Westhoff C, et al. Combined hormonal contraceptive trials: variable data collection and bleeding assessment methodologies influence study outcomes and physician perceptions. *Contraception.* 2007;75:4-10.
16. Mishell DR Jr, Guillebaud J, Westhoff C, et al. Recommendations for standardization of data collection and analysis of bleeding in combined hormone contraceptive trials. *Contraception.* 2007;75:11-15.
17. Belsey EM, Machin D, d'Arcangues D. The analysis of vaginal bleeding patterns induced by fertility regulating methods. *Contraception.* 1986;34:253-260.

ORTHO TRI-CYCLEN® Lo Tablets
(norgestimate/ethinyl estradiol)

Patients should be counseled that this product does not protect against HIV infection (AIDS) and other sexually transmitted diseases.

DESCRIPTION

ORTHO TRI-CYCLEN® Lo Tablets is a combination oral contraceptive containing the progestational compound norgestimate and the estrogenic compound ethinyl estradiol.


ORTHO TRI-CYCLEN® Lo Tablets

Each white tablet contains 0.180 mg of the progestational compound, norgestimate (+)-13-Ethyl-17-hydroxy-18,19-dinor-17 α -pregn-4-en-20-yn-3-one oxime acetate (ester) and 0.025 mg of the estrogenic compound, ethinyl estradiol (19-nor-17 α -pregna,1,3,5(10)-trien-20-yne-3,17-diol). Inactive ingredients include lactose, magnesium stearate, croscarmellose sodium, microcrystalline cellulose, carnauba wax, hypromellose, polyethylene glycol, titanium dioxide, and purified water.

Each light blue tablet contains 0.215 mg of the progestational compound norgestimate (+)-13-Ethyl-17-hydroxy-18,19-dinor-17 α -pregn-4-en-20-yn-3-one oxime acetate (ester) and 0.025 mg of the estrogenic compound, ethinyl estradiol (19-nor-17 α -pregna,1,3,5(10)-trien-20-yne-3,17-diol). Inactive ingredients include FD & C Blue No. 2 Aluminum Lake, lactose, magnesium stearate, croscarmellose sodium, microcrystalline cellulose, carnauba wax, hypromellose, polyethylene glycol, titanium dioxide, and purified water.

Each dark blue tablet contains 0.250 mg of the progestational compound norgestimate (+)-13-Ethyl-17-hydroxy-18,19-dinor-17 α -pregn-4-en-20-yn-3-one oxime acetate (ester) and 0.025 mg of the estrogenic compound, ethinyl estradiol (19-nor-17 α -pregna,1,3,5(10)-trien-20-yne-3,17-diol). Inactive ingredients include FD & C Blue No. 2 Aluminum Lake, lactose, magnesium stearate, croscarmellose sodium, microcrystalline cellulose, polysorbate 80, carnauba wax, hypromellose, polyethylene glycol, titanium dioxide, and purified water.

Each dark green tablet contains only inert ingredients, as follows: FD & C Blue No. 2 Aluminum Lake, lactose, magnesium stearate, pregelatinized starch, ferric oxide, hypromellose, polyethylene glycol, titanium dioxide, talc and purified water.



CLINICAL PHARMACOLOGY

Oral Contraception

Combination oral contraceptives act by suppression of gonadotropins. Although the primary mechanism of this action is inhibition of ovulation, other alterations include changes in the cervical mucus (which increase the difficulty of sperm entry into the uterus) and the endometrium (which reduce the likelihood of implantation).

Receptor binding studies, as well as studies in animals and humans, have shown that norgestimate and 17-deacetyl norgestimate, the major serum metabolite, combine high progestational activity with minimal intrinsic androgenicity.⁹⁰⁻⁹³ Norgestimate, in combination with ethinyl estradiol, does not counteract the estrogen-induced increases in sex hormone binding globulin (SHBG), resulting in lower serum testosterone.^{90,91,94}

PHARMACOKINETICS

Absorption

Norgestimate (NGM) and ethinyl estradiol (EE) are rapidly absorbed following oral administration. Norgestimate is rapidly and completely metabolized by first-pass (intestinal and/or hepatic) mechanisms to norelgestromin (NGMN) and norgestrel (NG), which are the major active metabolites of norgestimate. Mean pharmacokinetic parameters for NGMN, NG and EE during three cycles of administration of ORTHO TRI-CYCLEN® Lo are summarized in Table 1. These results indicate that: (1) Peak serum concentrations of NGMN and EE were generally reached by 2 hours after dosing; (2) Accumulation following multiple dosing of the 180 μ g NGM / 25 μ g dose is approximately 1.5 to 2 fold for NGMN and approximately 1.5 fold for EE compared with single dose administration, in agreement with that predicted based on linear kinetics of NGMN and EE; (3) The kinetics of NGMN are dose proportional following NGM doses of 180 to 250 μ g; (4) Steady-state conditions for NGMN following each NGM dose and for EE were achieved during the three cycle study; (5) Non-linear accumulation (4.5-14.5 fold) of norgestrel was observed as a result of high affinity

binding to SHBG, which limits its biological activity.¹⁰⁰ The effect of food on the pharmacokinetics of ORTHO TRI-CYCLEN® Lo has not been studied.

Table 1 provides a summary of norelgestromin, norgestrel and ethinyl estradiol pharmacokinetic parameters.

Table 1: Mean (SD) Pharmacokinetic Parameters of ORTHO TRI-CYCLEN® Lo During a Three Cycle Study

Analyte ¹	Cycle	Day	C _{max}	t _{max} (h)	AUC _{0-24h}	t _{1/2} (h)
NGMN ^(2,4)	1	1	0.91 (0.27)	1.8 (1.0)	5.86 (1.54)	NC
		7	1.42 (0.43)	1.8 (0.7)	11.3 (3.2)	NC
		14	1.57 (0.39)	1.8 (0.7)	13.9 (3.7)	NC
NG ^(2,4)	1	21	1.82 (0.54)	1.5 (0.7)	16.1 (4.8)	28.1 (10.6)
		1	0.32 (0.14)	2.0 (1.1)	2.44 (2.04)	NC
		3	1.64 (0.89)	1.9 (0.9)	27.9 (18.1)	NC
		14	2.11 (1.13)	4.0 (6.3)	40.7 (24.8)	NC
EE ^(2,3,5)	1	21	2.79 (1.42)	1.7 (1.2)	49.9 (27.6)	36.4 (10.2)
		1	55.6 (18.1)	1.7 (0.5)	421 (118)	NC
		3	91.1 (36.7)	1.3 (0.3)	782 (329)	NC
		14	96.9 (38.5)	1.3 (0.3)	796 (273)	NC
		21	95.9 (38.9)	1.3 (0.6)	771 (303)	17.7 (4.4)

¹ NGMN = Norelgestromin, NG = norgestrel, EE = ethinyl estradiol
² C_{max} = peak serum concentration, t_{max} = time to reach peak serum concentration, AUC_{0-24h} = area under serum concentration vs. time curve from 0 to 24 hours, t_{1/2} = elimination half-life.

³ units for all analytes; h = hours

⁴ units for NGMN and NG – C_{max} = ng/mL, AUC_{0-24h} = h.ng/mL

⁵ units for EE only – C_{max} = pg/mL, AUC_{0-24h} = h.pg/mL

NC = not calculated

Distribution

Norelgestromin and norgestrel (a serum metabolite of norelgestromin) are highly bound (>97%) to serum proteins. Norelgestromin is bound to albumin and not to SHBG, while norgestrel is bound primarily to SHBG. Ethinyl estradiol is extensively bound (> 97%) to serum albumin.

Metabolism

Norgestimate is extensively metabolized by first-pass mechanisms in the gastrointestinal tract and/or liver. Norgestimate's primary active metabolite is norelgestromin. Subsequent hepatic metabolism of norelgestromin occurs and metabolites include norgestrel, which is also active and various hydroxylated and conjugated metabolites. Ethinyl estradiol is also metabolized to various hydroxylated products and their glucuronide and sulfate conjugates.

Excretion

Following 3 cycles of administration of ORTHO TRI-CYCLEN® Lo, the mean (\pm SD) elimination half-life values, at steady-state, for norelgestromin, norgestrel and ethinyl estradiol were 28.1 (\pm 10.6) hours, 36.4 (\pm 10.2) hours and 17.7 (\pm 4.4) hours, respectively (Table 1). The metabolites of norelgestromin and ethinyl estradiol are eliminated by renal and fecal pathways.

Special Populations

Effects of Body Weight, Body Surface Area, and Age

The effects of body weight, body surface area, age and race on the pharmacokinetics of norelgestromin, norgestrel and ethinyl estradiol were evaluated in 79 healthy women using pooled data following single dose administration of NGM 180 or 250 μ g / EE 25 μ g tablets in four pharmacokinetic studies. Increasing body weight and body surface area were each associated with decreases in C_{max} and AUC_{0-24h} values for norelgestromin and ethinyl estradiol and increases in CL/F (oral clearance) for ethinyl estradiol. Increasing body weight by 10 kg is predicted to reduce the following parameters: NGMN C_{max} by 9% and AUC_{0-24h} by 19%, norgestrel C_{max} by 12% and AUC_{0-24h} by 46%, EE C_{max} by 13% and AUC_{0-24h} by 12%. These changes were statistically significant. Increasing age was associated with slight decreases (6% with increasing age by 5 years) in C_{max} and AUC_{0-24h} for norelgestromin and were statistically significant, but there was no significant effect for norgestrel or ethinyl estradiol. Only a small to moderate fraction (5-40%) of the overall variability in the pharmacokinetics of norelgestromin and ethinyl estradiol following ORTHO TRI-CYCLEN® Lo Tablets may be explained by any or all of the above demographic parameters.

In clinical studies involving 1673 subjects with a mean weight of 141 pounds, there was no association between pregnancy and weight.

Renal and Hepatic Impairment

No studies with ORTHO TRI-CYCLEN® Lo have been conducted in women with renal or hepatic impairment.

Drug-Drug Interactions

Although norelgestromin and its metabolites inhibit a variety of P450 enzymes in human liver microsomes, under the recommended dosing regimen, the in vivo concentrations of norelgestromin and its metabolites, even at the peak serum levels, are relatively low compared to the inhibitory constant (K_i).

Interactions between oral contraceptives and other drugs have been reported in the literature. No formal drug-drug interaction studies were conducted with ORTHO TRI-CYCLLEN® Lo (see PRECAUTIONS).

INDICATIONS AND USAGE

ORTHO TRI-CYCLLEN® Lo Tablets are indicated for the prevention of pregnancy in women who elect to use oral contraceptives as a method of contraception.

In an active controlled clinical trial 1,673 subjects completed 11,003 cycles of ORTHO TRI-CYCLLEN® Lo use and a total of 20 pregnancies were reported in ORTHO TRI-CYCLLEN® Lo users.⁹⁹ This represents an overall use-efficacy (typical user efficacy) pregnancy rate of 2.36 per 100 women-years of use.

Oral contraceptives are highly effective for pregnancy prevention. Table 2 lists the typical accidental pregnancy rates for users of combination oral contraceptives and other methods of contraception. The efficacy of these contraceptive methods, except sterilization, the IUD, and the Norplant® system, depends upon the reliability with which they are used. Correct and consistent use of methods can result in lower failure rates.

Method (1)	% of Women Experiencing an Unintended Pregnancy Within the First Year of Use		% of Women Continuing Use at One Year ³ (4)
	Typical Use ¹ (2)	Perfect Use ² (3)	
Chance ⁴	85	85	
Spermicides ⁵	26	6	40
Periodic abstinence	25		63
Calendar		9	
Ovulation Method		3	
Sympto-Thermal ⁶		2	
Post-Ovulation		1	
Withdrawal	19	4	
Cap ⁷			
Parous Women	40	26	42
Nulliparous Women	20	9	56
Sponge			
Parous Women	40	20	42
Nulliparous Women	20	9	56
Diaphragm ⁷	20	6	56
Condom ⁸			
Female (Reality®)	21	5	56
Male	14	3	61
Pill	5		71
Progestin Only		0.5	
Combined		0.1	
IUD			
Progesterone T	2.0	1.5	81
Copper T380A	0.8	0.6	78
LNg 20	0.1	0.1	81
Depo-Provera®	0.3	0.3	70
Norplant® and Norplant-2®	0.05	0.05	88
Female Sterilization	0.5	0.5	100
Male Sterilization	0.15	0.10	100

Emergency Contraceptives Pills: Treatment initiated within 72 hours after unprotected intercourse reduces the risk of pregnancy by at least 75%.⁹

Lactation Amenorrhea Method: LAM is a highly effective, temporary method of contraception.¹⁰

Source: Trussell J. Contraceptive efficacy. In Hatcher RA, Trussell J, Stewart F, Cates W, Stewart GK, Kowal D, Guest F, Contraceptive Technology: Seventeenth Revised Edition. New York NY: Irvington Publishers, 1998.

¹ Among *typical* couples who initiate use of a method (not necessarily for the first time), the percentage who experience an accidental pregnancy during the first year if they do not stop use for any other reason.

² Among couples who initiate use of a method (not necessarily for the first time) and who use it *perfectly* (both consistently and correctly), the percentage who experience an accidental pregnancy during the first year if they do not stop use for any other reason.

³ Among couples attempting to avoid pregnancy, the percentage who continue to use a method for one year.

⁴ The percents becoming pregnant in columns (2) and (3) are based on data from populations where contraception is not used and from women who cease using contraception in order to become pregnant. Among such populations, about 89% become pregnant within one

year. This estimate was lowered slightly (to 85%) to represent the percent who would become pregnant within one year among women now relying on reversible methods of contraception if they abandoned contraception altogether.

- ⁵ Foams, creams, gels, vaginal suppositories, and vaginal film.
- ⁶ Cervical mucus (ovulation) method supplemented by calendar in the pre-ovulatory and basal body temperature in the post-ovulatory phases.
- ⁷ With spermicidal cream or jelly.
- ⁸ Without spermicides.
- ⁹ The treatment schedule is one dose within 72 hours after unprotected intercourse, and a second dose 12 hours after the first dose. The FDA has declared the following brands of oral contraceptives to be safe and effective for emergency contraception: Ovral® (1 dose is 2 white pills), Alesse® (1 dose is 5 pink pills), Nordette® or Levlen® (1 dose is 4 yellow pills).
- ¹⁰ However, to maintain effective protection against pregnancy, another method of contraception must be used as soon as menstruation resumes, the frequency or duration of breastfeeds is reduced, bottle feeds are introduced, or the baby reaches 6 months of age.

ORTHO TRI-CYCLLEN® Lo has not been studied for and is not indicated for use in emergency contraception.

CONTRAINDICATIONS

Oral contraceptives should not be used in women who have any of the following conditions:

- Thrombophlebitis or thromboembolic disorders
- A past history of deep vein thrombophlebitis or thromboembolic disorders
- Cerebral vascular or coronary artery disease (current or history)
- Valvular heart disease with complications
- Severe hypertension
- Diabetes with vascular involvement
- Headaches with focal neurological symptoms
- Major surgery with prolonged immobilization
- Known or suspected carcinoma of the breast or personal history of breast cancer
- Carcinoma of the endometrium or other known or suspected estrogen-dependent neoplasia
- Undiagnosed abnormal genital bleeding
- Cholestatic jaundice of pregnancy or jaundice with prior pill use
- Hepatic adenomas or carcinomas
- Known or suspected pregnancy
- Hypersensitivity to any component of this product

WARNINGS

Cigarette smoking increases the risk of serious cardiovascular side effects from oral contraceptive use. This risk increases with age and with heavy smoking (15 or more cigarettes per day) and is quite marked in women over 35 years of age. Women who use oral contraceptives should be strongly advised not to smoke.

The use of oral contraceptives is associated with increased risks of several serious conditions including myocardial infarction, thromboembolism, stroke, hepatic neoplasia, and gallbladder disease, although the risk of serious morbidity or mortality is very small in healthy women without underlying risk factors. The risk of morbidity and mortality increases significantly in the presence of other underlying risk factors such as hypertension, hyperlipidemias, obesity and diabetes.

Practitioners prescribing oral contraceptives should be familiar with the following information relating to these risks.

The information contained in this package insert is principally based on studies carried out in patients who used oral contraceptives with higher formulations of estrogens and progestogens than those in common use today. The effect of long-term use of the oral contraceptives with lower formulations of both estrogens and progestogens remains to be determined.

Throughout this labeling, epidemiological studies reported are of two types: retrospective or case control studies and prospective or cohort studies. Case control studies provide a measure of the relative risk of a disease, namely, a ratio of the incidence of a disease among oral contraceptive users to that among nonusers. The relative risk does not provide information on the actual clinical occurrence of a disease. Cohort studies provide a measure of attributable risk, which is the difference in the incidence of disease between oral contraceptive users and nonusers. The attributable risk does provide information about the actual occurrence of a disease in the population (adapted from refs. 2 and 3 with the author's permission). For further information, the reader is referred to a text on epidemiological methods.

1. Thromboembolic Disorders and Other Vascular Problems

a. Myocardial Infarction

An increased risk of myocardial infarction has been attributed to oral contraceptive use. This risk is primarily in smokers or women with other underlying risk factors for coronary artery disease such as hypertension, hypercholesterolemia, morbid obesity, and diabetes. The relative risk of heart attack for current oral contraceptive users has been estimated to be two to six.⁴⁻¹⁰ The risk is very low under the age of 30.

Smoking in combination with oral contraceptive use has been shown to contribute substantially to the incidence of myocardial infarctions in women in their mid-thirties or older with smoking accounting for the majority of excess cases.¹¹ Mortality rates associated with circulatory disease have been shown to increase substantially in smokers, especially in those 35 years of age and older and in nonsmokers over the age of 40 among women who use oral contraceptives.

Figure 1: Circulatory Disease Mortality Rates Per 100,000 Women-Years By Age, Smoking Status And Oral Contraceptive Use




Figure 1 Adapted from P.M. Layde and V. Beral, Ref. #12.

Oral contraceptives may compound the effects of well-known risk factors, such as hypertension, diabetes, hyperlipidemias, age and obesity.¹³ In particular, some progestogens are known to decrease HDL cholesterol and cause glucose intolerance, while estrogens may create a state of hyperinsulinism.¹⁴⁻¹⁸ Oral contraceptives have been shown to increase blood pressure among users (see section 9 in WARNINGS). Similar effects on risk factors have been associated with an increased risk of heart disease. Oral contraceptives must be used with caution in women with cardiovascular disease risk factors.

Norgestimate has minimal androgenic activity (see CLINICAL PHARMACOLOGY), and there is some evidence that the risk of myocardial infarction associated with oral contraceptives is lower when the progestogen has minimal androgenic activity than when the activity is greater.⁹⁷

b. Thromboembolism

An increased risk of thromboembolic and thrombotic disease associated with the use of oral contraceptives is well established. Case control studies have found the relative risk of users compared to nonusers to be 3 for the first episode of superficial venous thrombosis, 4 to 11 for deep vein thrombosis or pulmonary embolism, and 1.5 to 6 for women with predisposing conditions for venous thromboembolic disease.^{2,3,19-24} Cohort studies have shown the relative risk to be somewhat lower, about 3 for new cases and about 4.5 for new cases requiring hospitalization.²⁵ The risk of thromboembolic disease associated with oral contraceptives is not related to length of use and disappears after pill use is stopped.²

A two- to four-fold increase in relative risk of post-operative thromboembolic complications has been reported with the use of oral contraceptives.⁹ The relative risk of venous thrombosis in women who have predisposing conditions is twice that of women without such medical conditions.²⁶ If feasible, oral contraceptives should be discontinued at least four weeks prior to and for two weeks after elective surgery of a type associated with an increase in risk of thromboembolism and during and following prolonged immobilization. Since the immediate postpartum period is also associated with an

increased risk of thromboembolism, oral contraceptives should be started no earlier than four weeks after delivery in women who elect not to breastfeed.

c. Cerebrovascular Diseases

Oral contraceptives have been shown to increase both the relative and attributable risks of cerebrovascular events (thrombotic and hemorrhagic strokes), although, in general, the risk is greatest among older (>35 years), hypertensive women who also smoke. Hypertension was found to be a risk factor for both users and nonusers, for both types of strokes, and smoking interacted to increase the risk of hemorrhagic stroke.²⁷⁻²⁹

In a large study, the relative risk of thrombotic strokes has been shown to range from 3 for normotensive users to 14 for users with severe hypertension.³⁰ The relative risk of hemorrhagic stroke is reported to be 1.2 for non-smokers who used oral contraceptives, 2.6 for smokers who did not use oral contraceptives, 7.6 for smokers who used oral contraceptives, 1.8 for normotensive users and 25.7 for users with severe hypertension.³⁰ The attributable risk is also greater in older women.³

d. Dose-Related Risk of Vascular Disease From Oral Contraceptives
A positive association has been observed between the amount of estrogen and progestogen in oral contraceptives and the risk of vascular disease.³¹⁻³³ A decline in serum high density lipoproteins (HDL) has been reported with many progestational agents.¹⁴⁻¹⁶ A decline in serum high density lipoproteins has been associated with an increased incidence of ischemic heart disease. Because estrogens increase HDL cholesterol, the net effect of an oral contraceptive depends on a balance achieved between doses of estrogen and progestogen and the activity of the progestogen used in the contraceptives. The activity and amount of both hormones should be considered in the choice of an oral contraceptive.

Minimizing exposure to estrogen and progestogen is in keeping with good principles of therapeutics. For any particular estrogen/progestogen combination, the dosage regimen prescribed should be one which contains the least amount of estrogen and progestogen that is compatible with a low failure rate and the needs of the individual patient. New acceptors of oral contraceptive agents should be started on preparations containing the lowest estrogen content which is judged appropriate for an individual patient.

e. Persistence of Risk of Vascular Disease

There are two studies which have shown persistence of risk of vascular disease for ever-users of oral contraceptives. In a study in the United States, the risk of developing myocardial infarction after discontinuing oral contraceptives persists for at least 9 years for women 40-49 years who had used oral contraceptives for five or more years, but this increased risk was not demonstrated in other age groups.⁸ In another study in Great Britain, the risk of developing cerebrovascular disease persisted for at least 6 years after discontinuation of oral contraceptives, although excess risk was very small.³⁴ However, both studies were performed with oral contraceptive formulations containing 50 micrograms or higher of estrogens.

2. Estimates of Mortality from Contraceptive Use

One study gathered data from a variety of sources which have estimated the mortality rate associated with different methods of contraception at different ages (Table 3). These estimates include the combined risk of death associated with contraceptive methods plus the risk attributable to pregnancy in the event of method failure. Each method of contraception has its specific benefits and risks. The study concluded that with the exception of oral contraceptive users 35 and older who smoke, and 40 and older who do not smoke, mortality associated with all methods of birth control is low and below that associated with childbirth. The observation of an increase in risk of mortality with age for oral contraceptive users is based on data gathered in the 1970's.³⁵ Current clinical recommendation involves the use of lower estrogen dose formulations and a careful consideration of risk factors. In 1989, the Fertility and Maternal Health Drugs Advisory Committee was asked to review the use of oral contraceptives in women 40 years of age and over.

The Committee concluded that although cardiovascular disease risks may be increased with oral contraceptive use after age 40 in healthy non-smoking women (even with the newer low-dose formulations), there are also greater potential health risks associated with pregnancy in older women and with the alternative surgical and medical procedures which may be necessary if such women do not have access to effective and acceptable means of contraception. The Committee recommended that the benefits of low-dose oral contraceptive use by healthy non-smoking women over 40 may outweigh the possible risks.

Of course, older women, as all women, who take oral contraceptives, should take an oral contraceptive which contains the least amount of estrogen and progestogen that is compatible with a low failure rate and individual patient needs.

Table 3: Annual Number of Birth-Related or Method-Related Deaths Associated with Control of Fertility Per 100,000 Nonsterile Women, By Fertility-Control Method and According to Age

Method of control and outcome	15-19	20-24	25-29	30-34	35-39	40-44
No fertility-control methods*	7.0	7.4	9.1	14.8	25.7	28.2
Oral contraceptives nonsmoker**	0.3	0.5	0.9	1.9	13.8	31.6
Oral contraceptives smoker**	2.2	3.4	6.6	13.5	51.1	117.2
IUD**	0.8	0.8	1.0	1.0	1.4	1.4
Condom*	1.1	1.6	0.7	0.2	0.3	0.4
Diaphragm/spermicide*	1.9	1.2	1.2	1.3	2.2	2.8
Periodic abstinence*	2.5	1.6	1.6	1.7	2.9	3.6

* Deaths are birth-related

** Deaths are method-related

Adapted from H.W. Ory, Family Planning Perspectives, Ref. #35.

3. Carcinoma of the Reproductive Organs and Breasts

Numerous epidemiological studies have been performed on the incidence of breast, endometrial, ovarian, and cervical cancer in women using oral contraceptives.

The risk of having breast cancer diagnosed may be slightly increased among current and recent users of combination oral contraceptives. However, this excess risk appears to decrease over time after discontinuation of combination oral contraceptives and by 10 years after cessation the increased risk disappears. Some studies report an increased risk with duration of use while other studies do not and no consistent relationships have been found with dose or type of steroid. Some studies have found a small increase in risk for women who first use combination oral contraceptives before age 20. Most studies show a similar pattern of risk with combination oral contraceptive use regardless of a woman's reproductive history or her family breast cancer history.

Breast cancers diagnosed in current or previous oral contraceptive users tend to be less clinically advanced than in nonusers.

Women who currently have or have had breast cancer should not use oral contraceptives because breast cancer is usually a hormonally-sensitive tumor.

Some studies suggest that oral contraceptive use has been associated with an increase in the risk of cervical intraepithelial neoplasia in some populations of women.⁴⁵⁻⁴⁸ However, there continues to be controversy about the extent to which such findings may be due to differences in sexual behavior and other factors.

In spite of many studies of the relationship between oral contraceptive use and breast and cervical cancers, a cause-and-effect relationship has not been established.

4. Hepatic Neoplasia

Benign hepatic adenomas are associated with oral contraceptive use, although the incidence of benign tumors is rare in the United States. Indirect calculations have estimated the attributable risk to be in the range of 3.3 cases/100,000 for users, a risk that increases after four or more years of use especially with oral contraceptives of higher dose.⁴⁹ Rupture of benign, hepatic adenomas may cause death through intra-abdominal hemorrhage.^{50,51}

Studies from Britain have shown an increased risk of developing hepatocellular carcinoma in long-term (>8 years) oral contraceptive users. However, these cancers are extremely rare in the U.S. and the attributable risk (the excess incidence) of liver cancers in oral contraceptive users approaches less than one per million users.

5. Ocular Lesions

There have been clinical case reports of retinal thrombosis associated with the use of oral contraceptives. Oral contraceptives should be discontinued if there is unexplained partial or complete loss of vision; onset of proptosis or diplopia; papilledema; or retinal vascular lesions. Appropriate diagnostic and therapeutic measures should be undertaken immediately.

6. Oral Contraceptive Use Before or During Early Pregnancy

Extensive epidemiological studies have revealed no increased risk of birth defects in women who have used oral contraceptives prior to pregnancy.^{56,57} The majority of recent studies also do not indicate a teratogenic effect, particularly in so far as cardiac anomalies and limb reduction defects are concerned,^{55,56,58,59} when taken inadvertently during early pregnancy.

The administration of oral contraceptives to induce withdrawal bleeding should not be used as a test for pregnancy. Oral contraceptives should not be used during pregnancy to treat threatened or habitual abortion.

It is recommended that for any patient who has missed two consecutive periods, pregnancy should be ruled out. If the patient has not adhered to the prescribed schedule, the possibility of pregnancy should be considered at the time of the first missed period. Oral contraceptive use should be discontinued if pregnancy is confirmed.

7. Gallbladder Disease

Earlier studies have reported an increased lifetime relative risk of gallbladder surgery in users of oral contraceptives and estrogens.^{60,61} More recent studies, however, have shown that the relative risk of developing gallbladder disease among oral contraceptive users may be minimal.⁶²⁻⁶⁴ The recent findings of minimal risk may be related to the use of oral contraceptive formulations containing lower hormonal doses of estrogens and progestogens.

8. Carbohydrate and Lipid Metabolic Effects

Oral contraceptives have been shown to cause a decrease in glucose tolerance in a significant percentage of users.¹⁷ This effect has been shown to be directly related to estrogen dose.⁶⁵ Progestogens increase insulin secretion and create insulin resistance, this effect varying with different progestational agents.^{17,66} However, in the non-diabetic woman, oral contraceptives appear to have no effect on fasting blood glucose.⁶⁷ Because of these demonstrated effects, prediabetic and diabetic women in particular should be carefully monitored while taking oral contraceptives.

A small proportion of women will have persistent hypertriglyceridemia while on the pill. As discussed earlier (see WARNINGS 1a and 1d), changes in serum triglycerides and lipoprotein levels have been reported in oral contraceptive users.

9. Elevated Blood Pressure

Women with significant hypertension should not be started on hormonal contraception.⁹⁸ An increase in blood pressure has been reported in women taking oral contraceptives⁶⁸ and this increase is more likely in older oral contraceptive users⁶⁹ and with extended duration of use.⁶¹ Data from the Royal College of General Practitioners¹² and subsequent randomized trials have shown that the incidence of hypertension increases with increasing progestational activity and concentrations of progestogens.

Women with a history of hypertension or hypertension-related diseases, or renal disease⁷⁰ should be encouraged to use another method of contraception. If women elect to use oral contraceptives, they should be monitored closely and if significant elevation of blood pressure occurs, oral contraceptives should be discontinued. For most women, elevated blood pressure will return to normal after stopping oral contraceptives, and there is no difference in the occurrence of hypertension between former and never users.⁶⁸⁻⁷¹

10. Headache

The onset or exacerbation of migraine or development of headache with a new pattern which is recurrent, persistent or severe requires discontinuation of oral contraceptives and evaluation of the cause.

11. Bleeding Irregularities

Breakthrough bleeding and spotting are sometimes encountered in patients on oral contraceptives, especially during the first three months of use. Non-hormonal causes should be considered and adequate diagnostic measures taken to rule out malignancy or pregnancy in the event of breakthrough bleeding, as in the case of any abnormal vaginal bleeding. If pathology has been excluded, time or a change to another formulation may solve the problem. In the event of amenorrhea, pregnancy should be ruled out.

Some women may encounter post-pill amenorrhea or oligomenorrhea, especially when such a condition was preexistent.

12. Ectopic Pregnancy

Ectopic as well as intrauterine pregnancy may occur in contraceptive failures.

PRECAUTIONS

1. General

Patients should be counseled that this product does not protect against HIV infection (AIDS) and other sexually transmitted diseases.

2. Physical Examination and Follow-Up

It is good medical practice for all women to have annual history and physical examinations, including women using oral contraceptives. The physical examination, however, may be deferred until after initiation of oral contraceptives if requested by the woman and judged appropriate by the clinician. The physical examination should include special reference to blood pressure, breasts, abdomen and pelvic organs, including cervical cytology, and relevant laboratory tests. In case of undiagnosed, persistent or recurrent abnormal vaginal bleeding, appropriate measures should be conducted to rule out malignancy. Women with a strong family history of breast cancer or who have breast nodules should be monitored with particular care.

3. Lipid Disorders

Women who are being treated for hyperlipidemias should be followed closely if they elect to use oral contraceptives. Some progestogens may elevate LDL levels and may render the control of hyperlipidemias more difficult.

4. Liver Function

If jaundice develops in any woman receiving oral contraceptives, the medication should be discontinued. Steroid hormones may be poorly metabolized in patients with impaired liver function.

5. Fluid Retention

Oral contraceptives may cause some degree of fluid retention. They should be prescribed with caution, and only with careful monitoring, in patients with conditions which might be aggravated by fluid retention.

6. Emotional Disorders

Women with a history of depression should be carefully observed and the drug discontinued if depression recurs to a serious degree.

7. Contact Lenses

Contact lens wearers who develop visual changes or changes in lens tolerance should be assessed by an ophthalmologist.

8. Drug Interactions

Changes in contraceptive effectiveness associated with co-administration of other products:

Contraceptive effectiveness may be reduced when hormonal contraceptives are coadministered with antibiotics, anticonvulsants, and other drugs that increase the metabolism of contraceptive steroids. This could result in unintended pregnancy or breakthrough bleeding. Examples include rifampin, barbiturates, phenylbutazone, phenytoin, carbamazepine, felbamate, oxcarbazepine, topiramate, griseofulvin and bosentan. Several cases of contraceptive failure and breakthrough bleeding have been reported in the literature with concomitant administration of antibiotics such as ampicillin and tetracyclines. However, clinical pharmacology studies investigating drug interaction between combined oral contraceptives and these antibiotics have reported inconsistent results.

Several of the anti-HIV protease inhibitors have been studied with co-administration of oral combination hormonal contraceptives; significant changes (increase and decrease) in the plasma levels of the estrogen and progestin have been noted in some cases. The safety and efficacy of oral contraceptive products may be affected with coadministration of anti-HIV protease inhibitors. Healthcare professionals should refer to the label of the individual anti-HIV protease inhibitors for further drug-drug interaction information.

Herbal products containing St. John's Wort (*hypericum perforatum*) may induce hepatic enzymes (cytochrome P450) and p-glycoprotein transporter and may reduce the effectiveness of contraceptive steroids. This may also result in breakthrough bleeding.

Concurrent use of bosentan and ethinyl estradiol containing products may result in decreased concentrations of these contraceptive hormones thereby increasing the risk of unintended pregnancy and unscheduled bleeding.

Increase in plasma ethinyl estradiol levels associated with co-administered drugs:

Co-administration of atorvastatin and certain oral contraceptives containing ethinyl estradiol increase AUC values for ethinyl estradiol by approximately 20%. Ascorbic acid and acetaminophen may increase plasma ethinyl estradiol levels, possibly by inhibition of conjugation. CYP 3A4 inhibitors such as itraconazole or ketoconazole may increase plasma hormone levels.

Changes in plasma levels of co-administered drugs:

Combination hormonal contraceptives containing some synthetic estrogens (e.g., ethinyl estradiol) may inhibit the metabolism of other compounds. Increased plasma concentrations of cyclosporin, prednisolone, and theophylline have been reported with concomitant administration of oral contraceptives. Decreased plasma concentrations of acetaminophen and increased clearance of temazepam, salicylic acid, morphine and clofibric acid, due to induction of conjugation, have been noted when drugs were administered with oral contraceptives.

Combined hormonal contraceptives have been shown to significantly decrease plasma concentrations of lamotrigine when co-administered due to induction of lamotrigine glucuronidation. This may reduce seizure control; therefore, dosage adjustments of lamotrigine may be necessary.¹⁰³

Healthcare professionals are advised to also refer to prescribing information of co-administered drugs for recommendations regarding management of concomitant therapy.

9. Interactions with Laboratory Tests

Certain endocrine and liver function tests and blood components may be affected by oral contraceptives:

a. Increased prothrombin and factors VII, VIII, IX, and X; decreased antithrombin 3; increased norepinephrine-induced platelet aggregability.

b. Increased thyroid binding globulin (TBG) leading to increased circulating total thyroid hormone, as measured by protein-bound iodine (PBI), T4 by column or by radioimmunoassay. Free T3 resin uptake is decreased, reflecting the elevated TBG, free T4 concentration is unaltered.

c. Other binding proteins may be elevated in serum.

d. Sex hormone binding globulins are increased and result in elevated levels of total circulating sex steroids; however, free or biologically active levels either decrease or remain unchanged.

e. Triglycerides may be increased and levels of various other lipids and lipoproteins may be affected.

f. Glucose tolerance may be decreased.

g. Serum folate levels may be depressed by oral contraceptive therapy. This may be of clinical significance if a woman becomes pregnant shortly after discontinuing oral contraceptives.

10. Carcinogenesis

See WARNINGS section.

11. Pregnancy

Pregnancy Category X. See CONTRAINDICATIONS and WARNINGS sections.

12. Nursing Mothers

Small amounts of oral contraceptive steroids have been identified in the milk of nursing mothers and a few adverse effects on the child have been reported, including jaundice and breast enlargement. In addition, oral contraceptives given in the postpartum period may interfere with lactation by decreasing the quantity and quality of breast milk. If possible, the nursing mother should be advised not to use combination oral contraceptives but to use other forms of contraception until she has completely weaned her child.

13. Pediatric Use

Safety and efficacy of ORTHO TRI-CYCLEN® Lo Tablets have been established in women of reproductive age. Safety and efficacy are expected to be the same for postpubertal adolescents under the age of 16 and for users 16 years and older. Use of this product before menarche is not indicated.

14. Geriatric Use

This product has not been studied in women over 65 years of age and is not indicated in this population.

INFORMATION FOR THE PATIENT

See Patient Labeling printed below.

ADVERSE REACTIONS

An increased risk of the following serious adverse reactions has been associated with the use of oral contraceptives (see WARNINGS section).

- Thrombophlebitis and venous thrombosis with or without embolism
- Arterial thromboembolism
- Pulmonary embolism
- Myocardial infarction
- Cerebral hemorrhage
- Cerebral thrombosis
- Hypertension
- Gallbladder disease
- Hepatic adenomas or benign liver tumors

There is evidence of an association between the following conditions and the use of oral contraceptives:

- Mesenteric thrombosis
- Retinal thrombosis

The following adverse reactions have been reported in patients receiving oral contraceptives and are believed to be drug-related:

- Nausea
- Vomiting
- Gastrointestinal symptoms (such as abdominal cramps and bloating)
- Breakthrough bleeding
- Spotting
- Change in menstrual flow
- Amenorrhea
- Temporary infertility after discontinuation of treatment
- Edema
- Melasma which may persist
- Breast changes: tenderness, enlargement, secretion
- Change in weight (increase or decrease)
- Change in cervical erosion and secretion
- Diminution in lactation when given immediately postpartum
- Cholestatic jaundice
- Migraine
- Allergic reaction, including rash, urticaria, and angioedema
- Mental depression
- Reduced tolerance to carbohydrates
- Vaginal candidiasis
- Change in corneal curvature (steepening)
- Intolerance to contact lenses

The following adverse reactions have been reported in users of oral contraceptives and the association has been neither confirmed nor refuted:

- Pre-menstrual syndrome
- Cataracts
- Changes in appetite
- Cystitis-like syndrome
- Headache
- Nervousness
- Dizziness
- Hirsutism
- Loss of scalp hair
- Erythema multiforme
- Erythema nodosum
- Hemorrhagic eruption
- Vaginitis
- Porphyria
- Impaired renal function
- Hemolytic uremic syndrome
- Acne
- Changes in libido
- Colitis
- Budd-Chiari Syndrome

OVERDOSAGE

Serious ill effects have not been reported following acute ingestion of large doses of oral contraceptives by young children. Overdosage may cause nausea and withdrawal bleeding may occur in females.

NON-CONTRACEPTIVE HEALTH BENEFITS

The following non-contraceptive health benefits related to the use of combination oral contraceptives are supported by epidemiological studies which largely utilized oral contraceptive formulations containing estrogen doses exceeding 0.035 mg of ethinyl estradiol or 0.05 mg mestranol.⁷³⁻⁷⁸

Effects on menses:

- increased menstrual cycle regularity
- decreased blood loss and decreased incidence of iron deficiency anemia
- decreased incidence of dysmenorrhea

Effects related to inhibition of ovulation:

- decreased incidence of functional ovarian cysts
- decreased incidence of ectopic pregnancies

Other effects:

- decreased incidence of fibroadenomas and fibrocystic disease of the breast
- decreased incidence of acute pelvic inflammatory disease
- decreased incidence of endometrial cancer
- decreased incidence of ovarian cancer

DOSAGE AND ADMINISTRATION

Oral Contraception

To achieve maximum contraceptive effectiveness, ORTHO TRI-CYCLEN® Lo Tablets must be taken exactly as directed and at intervals not exceeding 24 hours. The possibility of ovulation and conception prior to initiation of medication should be considered. ORTHO TRI-CYCLEN® Lo is available in the DIALPAK® Tablet Dispenser which is preset for a Sunday Start. Day 1 Start is also provided.

Sunday Start

When taking ORTHO TRI-CYCLEN® Lo the first white "active" tablet should be taken on the first Sunday after menstruation begins. If the menstrual period begins on Sunday, the first white "active" tablet should be taken that day. Take one white, light blue or dark blue "active" tablet daily for 21 days followed by one dark green "reminder" tablet daily for 7 days. After 28 tablets have been taken, a new course is started the next day (Sunday). For the first cycle of a Sunday Start regimen, another method of contraception should be used until after the first 7 consecutive days of administration.

If the patient misses one (1) white, light blue, or dark blue "active" tablet in Weeks 1, 2, or 3, the tablet should be taken as soon as she remembers. If the patient misses two (2) white or light blue "active" tablets in Week 1 or Week 2, the patient should take two (2) "active" tablets the day she remembers and two (2) "active" tablets the next day; and then continue taking one (1) tablet a day until she finishes the pack. The patient should be instructed to use a back-up method of birth control if she has sex in the seven (7) days after missing pills. If the patient misses two (2) dark blue "active" tablets in the third week or misses three (3) or more "active" tablets in a row, the patient should continue taking one tablet every day until Sunday. On Sunday the patient should throw out the rest of the pack and start a new pack that same day. The patient should be instructed to use a back-up method of birth control if she has sex in the seven (7) days after missing pills.

Complete instructions to facilitate patient counseling on proper pill usage may be found in the Detailed Patient Labeling ("How to Take the Pill" section).

Day 1 Start

The dosage of ORTHO TRI-CYCLEN® Lo for the initial cycle of therapy is one white, light blue or dark blue "active" tablet administered daily from the 1st day through the 21st day of the menstrual cycle, counting the first day of menstrual flow as "Day 1" followed by one dark green "reminder" tablet daily for 7 days. Tablets are taken without interruption for 28 days. After 28 tablets have been taken, a new course is started the next day.

If the patient misses one (1) white, light blue, or dark blue "active" tablet in Weeks 1, 2, or 3, the tablet should be taken as soon as she remembers. If the patient misses two (2) white or light blue "active" tablets in Week 1 or Week 2, the patient should take two (2) "active" tablets the day she remembers and two (2) "active" tablets the next day; and then continue taking one (1) tablet a day until she finishes the pack. The patient should be instructed to use a back-up method of birth control if she has sex in the seven (7) days after missing pills. If the patient misses two (2) dark blue "active" tablets in the third week or misses three (3) or more "active" tablets in a row, the patient should throw out the rest of the pack and start a new pack that same day. The patient should be instructed to use a back-up method of birth control if she has sex in the seven (7) days after missing pills.

Complete instructions to facilitate patient counseling on proper pill usage may be found in the Detailed Patient Labeling ("How to Take the Pill" section).

When switching from another oral contraceptive, ORTHO TRI-CYCLEN® Lo should be started on the same day that a new pack of the previous oral contraceptive would have been started.

The use of ORTHO TRI-CYCLEN® Lo for contraception may be initiated 4 weeks postpartum in women who elect not to breastfeed. When the tablets are administered during the postpartum period, the increased risk of thromboembolic disease associated with the postpartum period must be considered. (See CONTRAINDICATIONS and WARNINGS concerning thromboembolic disease. See also PRECAUTIONS for "Nursing Mothers.") The possibility of ovulation and conception prior to initiation of medication should be considered.

(See Discussion of Dose-Related Risk of Vascular Disease from Oral Contraceptives.)

ADDITIONAL INSTRUCTIONS FOR ALL DOSING REGIMENS

Breakthrough bleeding, spotting, and amenorrhea are frequent reasons for patients discontinuing oral contraceptives. In breakthrough bleeding, as in all cases of irregular bleeding from the vagina, nonfunctional causes should be borne in mind. In undiagnosed persistent or recurrent abnormal bleeding from the vagina, adequate diagnostic measures are indicated to rule out pregnancy or malignancy. If pathology has been excluded, time or a change to another formulation may solve the problem. Changing to an oral contraceptive with a higher estrogen content, while potentially useful in minimizing menstrual irregularity, should be done only if necessary since this may increase the risk of thromboembolic disease.

Use of oral contraceptives in the event of a missed menstrual period:

- 1.If the patient has not adhered to the prescribed schedule, the possibility of pregnancy should be considered at the time of the first missed period and oral contraceptive use should be discontinued if pregnancy is confirmed.
- 2.If the patient has adhered to the prescribed regimen and misses two consecutive periods, pregnancy should be ruled out before continuing oral contraceptive use.

HOW SUPPLIED

ORTHO TRI-CYCLEN® Lo Tablets are available in a DIALPAK® Tablet Dispenser (NDC 0062-1251-15) containing 28 tablets. Each of the 7 white, round, convex, coated tablets imprinted "O-M" on one side and "180" on the other side contains 0.180 mg of the progestational compound, norgestimate, together with 0.025 mg of the estrogenic compound, ethinyl estradiol. Each of the 7 light blue, round, convex, coated tablets imprinted "O-M" on one side and "215" on the other side contains 0.215 mg of the progestational compound, norgestimate, together with 0.025 mg of the estrogenic compound, ethinyl estradiol. Each of the 7 dark blue, round, convex, coated tablets imprinted "O-M" on one side and "250" on the other side contains 0.250 mg of the progestational compound, norgestimate, together with 0.025 mg of the estrogenic compound, ethinyl estradiol. Each of the 7 dark green, round, convex, coated tablets imprinted "O-M" on one side and "P" on the other side contains inert ingredients.

ORTHO TRI-CYCLEN® Lo Tablets are available for clinic usage in a VERIDATE® Tablet Dispenser (unfilled) and VERIDATE Refills (NDC 0062-1251-20).

Store at 25°C (77°F); excursions permitted to 15° - 30°C (59° - 86°F). Protect from light.

REFERENCES

1. Trussell J. Contraceptive efficacy. In Hatcher RA, Trussell J, Stewart F, Cates W, Stewart GK, Kowal D, Guest F, *Contraceptive Technology: Seventeenth Revised Edition*. New York NY: Irvington Publishers, 1998.
2. Stadel BV. Oral contraceptives and cardiovascular disease. (Pt.1). *N Engl J Med* 1981; 305:612-618.
3. Stadel BV. Oral contraceptives and cardiovascular disease. (Pt.2). *N Engl J Med* 1981; 305:672-677.
4. Adam SA, Thorogood M. Oral contraception and myocardial infarction revisited: the effects of new preparations and prescribing patterns. *Br J Obstet Gynaecol* 1981; 88:838-845.
5. Mann JI, Inman WH. Oral contraceptives and death from myocardial infarction. *Br Med J* 1975; 2(5965):245-248.
6. Mann JI, Vessey MP, Thorogood M, Doll R. Myocardial infarction in young women with special reference to oral contraceptive practice. *Br Med J* 1975; 2(5956):241-245.
7. Royal College of General Practitioners' Oral Contraception Study: Further analyses of mortality in oral contraceptive users. *Lancet* 1981; 1:541-546.
8. Slone D, Shapiro S, Kaufman DW, Rosenberg L, Miettinen OS, Stolley PD. Risk of myocardial infarction in relation to current and discontinued use of oral contraceptives. *N Engl J Med* 1981;305:420-424.
9. Vessey MP. Female hormones and vascular disease—an epidemiological overview. *Br J Fam Plann* 1980; 6(Supplement): 1-12.
10. Russell-Briefel RG, Ezzati TM, Fulwood R, Perlman JA, Murphy RS. Cardiovascular risk status and oral contraceptive use, United States, 1976-80. *Prevent Med* 1986; 15:352-362.
11. Goldbaum GM, Kendrick JS, Hogelin GC, Gentry EM. The relative impact of smoking and oral contraceptive use on women in the United States. *JAMA* 1987; 258: 1339-1342.
12. Layde PM, Beral V. Further analyses of mortality in oral contraceptive users; Royal College of General Practitioners' Oral Contraception Study. (Table 5) *Lancet* 1981; 1:541-546.
13. Knopp RH. Atherosclerosis risk: the roles of oral contraceptives and postmenopausal estrogens. *J Reprod Med* 1986; 31(9) (Supplement):913-921.
14. Krauss RM, Roy S, Mishell DR, Casagrande J, Pike MC. Effects of two low-dose oral contraceptives on serum lipids and lipoproteins: Differential changes in high-density lipoproteins subclasses. *Am J Obstet* 1983; 145:446-452.
15. Wahl P, Walden C, Knopp R, Hoover J, Wallace R, Heiss G, Rifkin B. Effect of estrogen/progestin potency on lipid/lipoprotein cholesterol. *N Engl J Med* 1983; 308:862-867.
16. Wynn V, Niththyananthan R. The effect of progestin in combined oral contraceptives on serum lipids with special reference to high density lipoproteins. *Am J Obstet Gynecol* 1982;142:766-771.
17. Wynn V, Godsland I. Effects of oral contraceptives on carbohydrate metabolism. *J Reprod Med* 1986;31(9)(Supplement):892-897.
18. LaRosa JC. Atherosclerotic risk factors in cardiovascular disease. *J Reprod Med* 1986;31(9) (Supplement):906-912.
19. Inman WH, Vessey MP. Investigation of death from pulmonary, coronary, and cerebral thrombosis and embolism in women of child-bearing age. *Br Med J* 1968;2(5599):193-199.
20. Maguire MG, Tonascia J, Sartwell PE, Stolley PD, Tockman MS. Increased risk of thrombosis due to oral contraceptives: a further report. *Am J Epidemiol* 1979;110(2):188-195.
21. Petitti DB, Wingerd J, Pellegrin F, Ramacharan S. Risk of vascular disease in women: smoking, oral contraceptives, noncontraceptive estrogens, and other factors. *JAMA* 1979;242:1150-1154.
22. Vessey MP, Doll R. Investigation of relation between use of oral contraceptives and thromboembolic disease. *Br Med J* 1968;2(5599):199-205.
23. Vessey MP, Doll R. Investigation of relation between use of oral contraceptives and thromboembolic disease. A further report. *Br Med J* 1969; 2(5658):651-657.
24. Porter JB, Hunter JR, Danielson DA, Jick H, Stergachis A. Oral contraceptives and non-fatal vascular disease—recent experience. *Obstet Gynecol* 1982;59(3):299-302.
25. Vessey M, Doll R, Peto R, Johnson B, Wiggins P. A long-term follow-up study of women using different methods of contraception: an interim report. *J Biosocial Sci* 1976;8:375-427.
26. Royal College of General Practitioners: Oral Contraceptives, venous thrombosis, and varicose veins. *J Royal Coll Gen Pract* 1978; 28:393-399.
27. Collaborative Group for the Study of Stroke in Young Women: Oral contraception and increased risk of cerebral ischemia or thrombosis. *N Engl J Med* 1973;288:871-878.
28. Petitti DB, Wingerd J. Use of oral contraceptives, cigarette smoking, and risk of subarachnoid hemorrhage. *Lancet* 1978;2:234-236.
29. Inman WH. Oral contraceptives and fatal subarachnoid hemorrhage. *Br Med J* 1979;2(6203):1468-1470.
30. Collaborative Group for the Study of Stroke in Young Women: Oral Contraceptives and stroke in young women: associated risk factors. *JAMA* 1975; 231:718-722.
31. Inman WH, Vessey MP, Westerholm B, Englund A. Thromboembolic disease and the steroidal content of oral contraceptives. A report to the Committee on Safety of Drugs. *Br Med J* 1970;2:203-209.
32. Meade TW, Greenberg G, Thompson SG. Progestogens and cardiovascular reactions associated with oral contraceptives and a comparison of the safety of 50- and 35-mcg oestrogen preparations. *Br Med J* 1980;280(6224):1157-1161.
33. Kay CR. Progestogens and arterial disease—evidence from the Royal College of General Practitioners' Study. *Am J Obstet Gynecol* 1982;142:762-765.
34. Royal College of General Practitioners: Incidence of arterial disease among oral contraceptive users. *J Royal Coll Gen Pract* 1983;33:75-82.
35. Ory HW. Mortality associated with fertility and fertility control: 1983. *Family Planning Perspectives* 1983;15:50-56.
36. The Cancer and Steroid Hormone Study of the Centers for Disease Control and the National Institute of Child Health and Human Development: Oral contraceptive use and the risk of breast cancer. *N Engl J Med* 1986;315:405-411.
37. Pike MC, Henderson BE, Krailo MD, Duke A, Roy S. Breast cancer in young women and use of oral contraceptives: possible modifying effect of formulation and age at use. *Lancet* 1983;2:926-929.
38. Paul C, Skegg DG, Spears GFS, Kaldor JM. Oral contraceptives and breast cancer: A national study. *Br Med J* 1986; 293:723-725.
39. Miller DR, Rosenberg L, Kaufman DW, Schottenfeld D, Stolley PD, Shapiro S. Breast cancer risk in relation to early oral contraceptive use. *Obstet Gynecol* 1986;68:863-868.
40. Olsson H, Olsson ML, Moller TR, Ranstam J, Holm P. Oral contraceptive use and breast cancer in young women in Sweden (letter). *Lancet* 1985; 1(8431):748-749.
41. McPherson K, Vessey M, Neil A, Doll R, Jones L, Roberts M. Early contraceptive use and breast cancer: Results of another case-control study. *Br J Cancer* 1987; 56:653-660.
42. Huggins GR, Zucker PF. Oral contraceptives and neoplasia; 1987 update. *Fertil Steril* 1987; 47:733-761.
43. McPherson K, Drife JO. The pill and breast cancer: why the uncertainty? *Br Med J* 1986; 293:709-710.
44. Shapiro S. Oral contraceptives—time to take stock. *N Engl J Med* 1987; 315:450-451.
45. Ory H, Naib Z, Conger SB, Hatcher RA, Tyler CW. Contraceptive choice and prevalence of cervical dysplasia and carcinoma in situ. *Am J Obstet Gynecol* 1976; 124:573-577.
46. Vessey MP, Lawless M, McPherson K, Yeates D. Neoplasia of the cervix uteri and contraception: a possible adverse effect of the pill. *Lancet* 1983; 2:930.
47. Brinton LA, Huggins GR, Lehman HF, Mallik K, Savitz DA, Trapido E, Rosenthal J, Hoover R. Long term use of oral contraceptives and risk of invasive cervical cancer. *Int J Cancer* 1986; 38:339-344.
48. WHO Collaborative Study of Neoplasia and Steroid Contraceptives: Invasive cervical cancer and combined oral contraceptives. *Br Med J* 1985; 290:961-965.
49. Rooks JB, Ory HW, Ishak KG, Strauss LT, Greenspan JR, Hill AP, Tyler CW. Epidemiology of hepatocellular adenoma: the role of oral contraceptive use. *JAMA* 1979; 242:644-648.
50. Bein NN, Goldsmith HS. Recurrent massive hemorrhage from benign hepatic tumors secondary to oral contraceptives. *Br J Surg* 1977; 64:433-435.
51. Klatskin G. Hepatic tumors: possible relationship to use of oral contraceptives. *Gastroenterology* 1977; 73:386-394.
52. Henderson BE, Preston-Martin S, Edmondson HA, Peters RL, Pike MC. Hepatocellular carcinoma and oral contraceptives. *Br J Cancer* 1983;48:437-440.
53. Neuberger J, Forman D, Doll R, Williams R. Oral contraceptives and hepatocellular carcinoma. *Br Med J* 1986; 292:1355-1357.
54. Forman D, Vincent TJ, Doll R. Cancer of the liver and oral contraceptives. *Br Med J* 1986; 292:1357-1361.
55. Harlap S, Eldor J. Births following oral contraceptive failures. *Obstet Gynecol* 1980; 55:447-452.
56. Savolainen E, Saksela E, Saxen L. Teratogenic hazards of oral contraceptives analyzed in a national malformation register. *Am J Obstet Gynecol* 1981; 140:521-524.
57. Janerich DT, Piper JM, Glebatis DM. Oral contraceptives and birth defects. *Am J Epidemiol* 1980; 112:73-79.
58. Ferencz C, Matanoski GM, Wilson PD, Rubin JD, Neill CA, Gutberlet R. Maternal hormone therapy and congenital heart disease. *Teratology* 1980; 21:225-239.
59. Rothman KJ, Fyler DC, Goldblatt A, Kreidberg MB. Exogenous hormones and other drug exposures of children with congenital heart disease. *Am J Epidemiol* 1979; 109:433-439.
60. Boston Collaborative Drug Surveillance Program: Oral contraceptives and venous thromboembolic disease, surgically confirmed gallbladder disease, and breast tumors. *Lancet* 1973; 1:1399-1404.
61. Royal College of General Practitioners: Oral contraceptives and health. New York, Pittman 1974.
62. Layde PM, Vessey MP, Yeates D. Risk of gallbladder disease: a cohort study of young women attending family planning clinics. *J Epidemiol Community Health* 1982; 36:274-278.
63. Rome Group for Epidemiology and Prevention of Cholelithiasis (GREPCO): Prevalence of gallstone disease in an Italian adult female population. *Am J Epidemiol* 1984; 119:796-805.
64. Storm BL, Tamragouri RT, Morse ML, Lazar EL, West SL, Stolley PD, Jones JK. Oral contraceptives and other risk factors for gallbladder disease. *Clin Pharmacol Ther* 1986; 39:335-341.
65. Wynn V, Adams PW, Godsland IF, Melrose J, Niththyananthan R, Oakley NW, Seedj A. Comparison of effects of different combined oral contraceptive formulations on carbohydrate and lipid metabolism. *Lancet* 1979; 1:1045-1049.
66. Wynn V. Effect of progesterone and progestins on carbohydrate metabolism. In: Progesterone and Progestin. Bardin CW, Milgrom E, Mauvis-Jarvis P. eds. New York, Raven Press 1983; pp. 395-410.
67. Perlman JA, Russell-Briefel RG, Ezzati TM, Lieberknecht G. Oral glucose tolerance and the potency of oral contraceptive progestogens. *J Chronic Dis* 1985;38:857-864.
68. Royal College of General Practitioners' Oral Contraception Study: Effect on hypertension and benign breast disease of progestogen component in combined oral contraceptives. *Lancet* 1977; 1:624.
69. Fisch IR, Frank J. Oral contraceptives and blood pressure. *JAMA* 1977; 237:2499-2503.
70. Laragh AJ. Oral contraceptive induced hypertension—nine years later. *Am J Obstet Gynecol* 1976; 126:141-147.
71. Ramcharan S, Peritz E, Pellegrin FA, Williams WT. Incidence of hypertension in the Walnut Creek Contraceptive Drug Study cohort:

In: Pharmacology of steroid contraceptive drugs. Garattini S, Berendes HW. Eds. New York, Raven Press, 1977; pp. 277-288, (Monographs of the Mario Negri Institute for Pharmacological Research Milan.)

72. Stockley I. Interactions with oral contraceptives. *J Pharm* 1976;216:140-143. **73.** The Cancer and Steroid Hormone Study of the Centers for Disease Control and the National Institute of Child Health and Human Development: Oral contraceptive use and the risk of ovarian cancer. *JAMA* 1983; 249:1596-1599. **74.** The Cancer and Steroid Hormone Study of the Centers for Disease Control and the National Institute of Child Health and Human Development: Combination oral contraceptive use and the risk of endometrial cancer. *JAMA* 1987; 257:796-800. **75.** Ory HW. Functional ovarian cysts and oral contraceptives: negative association confirmed surgically. *JAMA* 1974; 228:68-69. **76.** Ory HW, Cole P, MacMahon B, Hoover R. Oral contraceptives and reduced risk of benign breast disease. *N Engl J Med* 1976; 294:419-422. **77.** Ory HW. The noncontraceptive health benefits from oral contraceptive use. *Fam Plann Perspect* 1982; 14:182-184. **78.** Ory HW, Forrest JD, Lincoln R. Making choices: Evaluating the health risks and benefits of birth control methods. New York, The Alan Guttmacher Institute, 1983; p.1. **79.** Schlesselman J, Stadel BV, Murray P, Lai S. Breast cancer in relation to early use of oral contraceptives. *JAMA* 1988; 259:1828-1833. **80.** Hennekens CH, Speizer FE, Lipnick RJ, Rosner B, Bain C, Belanger C, Stampfer MJ, Willett W, Peto R. A case-control study of oral contraceptive use and breast cancer. *JNCI* 1984; 72:39-42. **81.** LaVecchia C, Decarli A, Fasoli M, Franceschi S, Gentile A, Negri E, Parazzini F, Tognoni G. Oral contraceptives and cancers of the breast and of the female genital tract. Interim results from a case-control study. *Br J Cancer* 1986; 54:311-317. **82.** Meirik O, Lund E, Adami H, Bergstrom R, Christoffersen T, Bergsjø P. Oral contraceptive use and breast cancer in young women. A Joint National Case-control study in Sweden and Norway. *Lancet* 1986; 11:650-654. **83.** Kay CR, Hannaford PC. Breast cancer and the pill—A further report from the Royal College of General Practitioners' oral contraception study. *Br J Cancer* 1988;58:675-680. **84.** Stadel BV, Lai S, Schlesselman JJ, Murray P. Oral contraceptives and premenopausal breast cancer in nulliparous women. *Contraception* 1988; 38:287-299. **85.** Miller DR, Rosenberg L, Kaufman DW, Stolley P, Warshauer ME, Shapiro S. Breast cancer before age 45 and oral contraceptive use: New Findings. *Am J Epidemiol* 1989; 129:269-280. **86.** The UK National Case-Control Study Group. Oral contraceptive use and breast cancer risk in young women. *Lancet* 1989; 1:973-982. **87.** Schlesselman JJ. Cancer of the breast and reproductive tract in relation to use of oral contraceptives. *Contraception* 1989; 40:1-38. **88.** Vessey MP, McPherson K, Villard-Mackintosh L, Yeates D. Oral contraceptives and breast cancer: latest findings in a large cohort study. *Br J Cancer* 1989; 59:613-617. **89.** Jick SS, Walker AM, Stergachis A, Jick H. Oral contraceptives and breast cancer. *Br J Cancer* 1989; 59:618-621. **90.** Anderson FD. Selectivity and minimal androgenicity of norgestimate in monophasic and triphasic oral contraceptives. *Acta Obstet Gynecol Scand* 1992; 156 (Supplement):15-21. **91.** Chapdelaine A, Desmaris J-L, Derman RJ. Clinical evidence of minimal androgenic activity of norgestimate. *Int J Fertil* 1989; 34(51):347-352. **92.** Phillips A, Demarest K, Hahn DW, Wong F, McGuire JL. Progestational and androgenic receptor binding affinities and in vivo activities of norgestimate and other progestins. *Contraception* 1989; 41(4):399-409. **93.** Phillips A, Hahn DW, Klimek S, McGuire JL. A comparison of the potencies and activities of progestogens used in contraceptives. *Contraception* 1987; 36(2):181-192. **94.** Janaud A, Rouffy J, Upmalis D, Dain M-P. A comparison study of lipid and androgen metabolism with triphasic oral contraceptive formulations containing norgestimate or levonorgestrel. *Acta Obstet Gynecol Scand* 1992; 156 (Supplement):34-38. **95.** Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and hormonal contraceptives: collaborative reanalysis of individual data on 53,297 women with breast cancer and 100,239 women without breast cancer from 54 epidemiological studies. *Lancet* 1996; 347:1713-1727. **96.** Palmer JR, Rosenberg L, Kaufman DW, Warshauer ME, Stolley P, Shapiro S. Oral Contraceptive Use and Liver Cancer. *Am J Epidemiol* 1989;130:878-882. **97.** Lewis M, Spitzer WO, Heinemann LAJ, MacRae KD, Bruppacher R, Thorogood M on behalf of Transnational Research Group on Oral Contraceptives and Health of Young Women. Third generation oral contraceptives and risk of myocardial infarction: an international case-control study. *Br Med J*, 1996;312:88-90. **98.** Improving access to quality care in family planning: Medical eligibility criteria for contraceptive use. Geneva, WHO, Family and Reproductive Health, 1996. **99.** Hampton RM, Short M, Bieber E, et al. Comparison of a novel norgestimate/ethinyl estradiol oral contraceptive (Ortho Tri-Cyclen Lo) with the oral contraceptive Loestrin Fe 1/20. *Contraception* 2001;63:289-295. **100.** Sitteri PK, Murai JT, Hammond GL, Nisker JA, Raymoure WJ, Huhn RW. The serum transport of steroid hormones. *Rec Prog Horm Res* 1982;38:457-510. **101.** Bork K, Fischer B, DeWald G. Recurrent episodes of skin angioedema and severe attacks of abdominal pain induced by oral contraceptives or hormone replacement therapy. *Am J Med* 2003;114:294-298. **102.** Van

Giersbergen PLM, Halabi A, Dingemans J. Pharmacokinetic interaction between bosentan and the oral contraceptives norethisterone and ethinyl estradiol. *Int J Clin Pharmacol Ther* 2006;44(3):113-118. **103.** Christensen J, Petrenaite V, Atterman J, et al. Oral contraceptives induce lamotrigine metabolism: evidence from a double-blind, placebo-controlled trial. *Epilepsia* 2007;48(3):484-489.

BRIEF SUMMARY PATIENT PACKAGE INSERT

Oral contraceptives, also known as "birth control pills" or "the pill," are taken to prevent pregnancy. When taken correctly without missing any pills, oral contraceptives are highly effective; however the typical failure rate of large numbers of pill users is 5% per year when women who miss pills are included. Forgetting to take pills considerably increases the chances of pregnancy. For most women oral contraceptives are also free of serious or unpleasant side effects.

For the majority of women, oral contraceptives can be taken safely. But there are some women who are at high risk of developing certain serious diseases that can be fatal or may cause temporary or permanent disability. The risks associated with taking oral contraceptives increase significantly if you:

- smoke
- have high blood pressure, diabetes, high cholesterol
- have or have had clotting disorders, heart attack, stroke, angina pectoris, cancer of the breast or sex organs, jaundice or malignant or benign liver tumors

Although cardiovascular disease risks may be increased with oral contraceptive use after age 40 in healthy, non-smoking women (even with the newer low-dose formulations), there are also greater potential health risks associated with pregnancy in older women.

You should not take the pill if you suspect you are pregnant or have unexplained vaginal bleeding.

Cigarette smoking increases the risk of serious cardiovascular side effects from oral contraceptive use. This risk increases with age and with heavy smoking (15 or more cigarettes per day) and is quite marked in women over 35 years of age. Women who use oral contraceptives are strongly advised not to smoke.

Most side effects of the pill are not serious. The most common such effects are nausea, vomiting, bleeding between menstrual periods, weight gain, breast tenderness, and difficulty wearing contact lenses. These side effects, especially nausea and vomiting, may subside within the first three months of use.

The serious side effects of the pill occur very infrequently, especially if you are in good health and are young. However, you should know that the following medical conditions have been associated with or made worse by the pill:

1. Blood clots in the legs (thrombophlebitis), lungs (pulmonary embolism), stoppage or rupture of a blood vessel in the brain (stroke), blockage of blood vessels in the heart (heart attack or angina pectoris) or other organs of the body. As mentioned above, smoking increases the risk of heart attacks and strokes and subsequent serious medical consequences.

2. In rare cases, oral contraceptives can cause benign but dangerous liver tumors. These benign liver tumors can rupture and cause fatal internal bleeding. In addition, some studies report an increased risk of developing liver cancer. However, liver cancers are rare.

3. High blood pressure, although blood pressure usually returns to normal when the pill is stopped.

The symptoms associated with these serious side effects are discussed in the detailed leaflet given to you with your supply of pills. Notify your healthcare professional if you notice any unusual physical disturbances while taking the pill. In addition, drugs such as rifampin, bosentan, as well as some anti-convulsants and some antibiotics, and herbal preparations containing St. John's Wort (hypericum perforatum) may decrease oral contraceptive effectiveness.

Oral contraceptives may interact with lamotrigine (LAMICTAL®), an anticonvulsant used for epilepsy. This may increase the risk of seizures so your healthcare professional may need to adjust the dose of lamotrigine.

Various studies give conflicting reports on the relationship between breast cancer and oral contraceptive use. Oral contraceptive use may slightly increase your chance of having breast cancer diagnosed, particularly after using hormonal contraceptives at a younger age. After you stop using hormonal contraceptives, the chances of having breast cancer diagnosed begin to go back down. You should have regular breast examinations by a healthcare professional and examine your own breasts monthly. Tell your healthcare professional if you have a family history of breast cancer or if you have had breast nodules or an abnormal mammogram. Women who currently have or have had breast cancer should not use oral contraceptives because breast cancer is usually a hormone-sensitive tumor.

Some studies have found an increase in the incidence of cancer of the cervix in women who use oral contraceptives. However, this finding may be related to factors other than the use of oral contraceptives. There is insufficient evidence to rule out the possibility that the pill may cause such cancers.

Taking the combination pill provides some important non-contraceptive benefits. These include less painful menstruation, less menstrual blood loss and anemia, fewer pelvic infections, and fewer cancers of the ovary and the lining of the uterus.

Be sure to discuss any medical condition you may have with your healthcare professional. Your healthcare professional will take a medical and family history before prescribing oral contraceptives and will examine you. The physical examination may be delayed to another time if you request it and the healthcare professional believes that it is a good medical practice to postpone it. You should be reexamined at least once a year while taking oral contraceptives. Your pharmacist should have given you the detailed patient information labeling which gives you further information which you should read and discuss with your healthcare professional.

ORTHO TRI-CYCLEN® Lo (like all oral contraceptives) is intended to prevent pregnancy. Oral contraceptives do not protect against transmission of HIV (AIDS) and other sexually transmitted diseases such as chlamydia, genital herpes, genital warts, gonorrhea, hepatitis B, and syphilis.

HOW TO TAKE THE PILL

IMPORTANT POINTS TO REMEMBER

BEFORE YOU START TAKING YOUR PILLS:

1. **BE SURE TO READ THESE DIRECTIONS:**
Before you start taking your pills.
Anytime you are not sure what to do.
2. **THE RIGHT WAY TO TAKE THE PILL IS TO TAKE ONE PILL EVERY DAY AT THE SAME TIME.**
If you miss pills you could get pregnant. This includes starting the pack late.
The more pills you miss, the more likely you are to get pregnant.
3. **MANY WOMEN HAVE SPOTTING OR LIGHT BLEEDING, OR MAY FEEL SICK TO THEIR STOMACH DURING THE FIRST 1-3 PACKS OF PILLS.** If you feel sick to your stomach, do not stop taking the pill. The problem will usually go away. If it doesn't go away, check with your healthcare professional.
4. **MISSING PILLS CAN ALSO CAUSE SPOTTING OR LIGHT BLEEDING,** even when you make up these missed pills.
On the days you take 2 pills to make up for missed pills, you could also feel a little sick to your stomach.
5. **IF YOU HAVE VOMITING OR DIARRHEA, or IF YOU TAKE SOME MEDICINES,** including some antibiotics, your pills may not work as well.
Use a back-up method (such as condoms or spermicides) until you check with your healthcare professional.
6. **IF YOU HAVE TROUBLE REMEMBERING TO TAKE THE PILL,** talk to your healthcare professional about how to make pill-taking easier or about using another method of birth control.
7. **IF YOU HAVE ANY QUESTIONS OR ARE UNSURE ABOUT THE INFORMATION IN THIS LEAFLET,** call your healthcare professional.

BEFORE YOU START TAKING YOUR PILLS

1. **DECIDE WHAT TIME OF DAY YOU WANT TO TAKE YOUR PILL.**
It is important to take it at about the same time every day.
2. The 28-pill pack has 21 white, light blue, and dark blue "active" pills (with hormones) to take for 3 weeks. This is followed by 1 week of dark green "reminder" pills (without hormones).
3. **ALSO FIND:**
 - 1) where on the pack to start taking pills,
 - 2) in what order to take the pills.

CHECK PICTURE OF PILL PACK AND ADDITIONAL INSTRUCTIONS FOR USING THIS PACKAGE IN THE BRIEF SUMMARY PATIENT PACKAGE INSERT.

4. **BE SURE YOU HAVE READY AT ALL TIMES:**
ANOTHER KIND OF BIRTH CONTROL (such as condoms or spermicide) to use as a back-up method in case you miss pills.
AN EXTRA, FULL PILL PACK.

WHEN TO START THE FIRST PACK OF PILLS

You have a choice of which day to start taking your first pack of pills. ORTHO TRI-CYCLEN® Lo is available in the DIALPAK® Tablet Dispenser which is preset for a Sunday Start. Day 1 Start is also provided. Decide with your healthcare professional which is the best day for you. Pick a time of day that will be easy to remember.

Sunday Start:

Take the first white "active" pill of the first pack on the Sunday after your period starts, even if you are still bleeding. If your period begins on Sunday, start the pack that same day.

Use another method of birth control (such as condoms or spermicide) as a back-up method if you have sex anytime from the Sunday you start your first pack until the next Sunday (7 days).

Day 1 Start:

Take the first white "active" pill of the first pack during the first 24 hours of your period.

You will not need to use a back-up method of birth control, since you are starting the pill at the beginning of your period.

WHAT TO DO DURING THE MONTH

1. Take One Pill At The Same Time Every Day Until The Pack Is Empty

Do not skip pills even if you are spotting or bleeding between monthly periods or feel sick to your stomach (nausea).

Do not skip pills even if you do not have sex very often.

2. When You Finish A Pack Or Switch Your Brand Of Pills

Start the next pack on the day after your last dark green "reminder" pill. Do not wait any days between packs.

WHAT TO DO IF YOU MISS PILLS

If you **MISS 1** white, light blue or dark blue "active" pill:

1. Take it as soon as you remember. Take the next pill at your regular time. This means you may take 2 pills in 1 day.
2. You do not need to use a back-up birth control method if you have sex.

If you **MISS 2** white or light blue "active" pills in a row in **WEEK 1 OR WEEK 2** of your pack:

1. Take 2 pills on the day you remember and 2 pills the next day.
2. Then take 1 pill a day until you finish the pack.
3. You **COULD BECOME PREGNANT** if you have sex in the 7 days after you miss pills. You **MUST** use another birth control method (such as condoms or spermicide) as a back-up method for those 7 days.

If you **MISS 2** dark blue "active" pills in a row in **THE 3RD WEEK**:

1. If you are a Sunday Starter:

Keep taking 1 pill every day until Sunday. On Sunday, **THROW OUT** the rest of the pack and start a new pack of pills that same day.

If you are a Day 1 Starter:

THROW OUT the rest of the pill pack and start a new pack that same day.

2. You may not have your period this month but this is expected. However, if you miss your period 2 months in a row, call your healthcare professional because you might be pregnant.
3. You **COULD BECOME PREGNANT** if you have sex in the 7 days after you miss pills. You **MUST** use another birth control method (such as condoms or spermicide) as a back-up method for those 7 days.

If you **MISS 3 OR MORE** white, light blue or dark blue "active" pills in a row (during the first 3 weeks):

1. If you are a Sunday Starter:

Keep taking 1 pill every day until Sunday. On Sunday, **THROW OUT** the rest of the pack and start a new pack of pills that same day.

If you are a Day 1 Starter:

THROW OUT the rest of the pill pack and start a new pack that same day.

2. You may not have your period this month but this is expected. However, if you miss your period 2 months in a row, call your healthcare professional because you might be pregnant.
3. You **COULD BECOME PREGNANT** if you have sex in the 7 days after you miss pills. You **MUST** use another birth control method (such as condoms or spermicide) as a back-up method for those 7 days.

If you forget any of the 7 dark green "reminder" pills in Week 4:

THROW AWAY the pills you missed.

Keep taking 1 pill each day until the pack is empty.

You do not need a back-up method.

FINALLY, IF YOU ARE STILL NOT SURE WHAT TO DO ABOUT THE PILLS YOU HAVE MISSED:

Use a **BACK-UP METHOD** anytime you have sex.

KEEP TAKING ONE "ACTIVE" PILL EACH DAY until you can reach your healthcare professional.


INSTRUCTIONS FOR USING YOUR DIALPAK® TABLET DISPENSER

Please Read Me!

 Sunday Start or Day 1 Start

There are two ways to start taking birth control pills: Sunday Start or Day 1 Start. Your healthcare provider will tell you which to use.

Save these instructions.




1 If this is the first time you are taking birth control pills, or if you have not taken birth control pills for 10 days or more, your first step is to **wait until the first day you get your menstrual period**. Then, follow these instructions for either Sunday Start or Day 1 Start.



Sunday Start
 Day 1 Start


When you get your period:

- You will use a **Sunday Start** if your doctor told you to take your first pill on a Sunday. Take pill "1" on the Sunday after your period starts. If your period starts on a Sunday, take pill "1" that day.
- You will use a **Day 1 Start** if your doctor told you to take pill "1" on the first day of your period.



SET THE DAY:

- Sunday Start:** the arrow on your *empty* Dialpak should point to SU (Sunday).
- Day 1 Start:** turn the dial on your *empty* Dialpak until the arrow points to the first day of your period (if your period starts on Tuesday, the arrow will point to TU).




4 Insert the new refill by lining up the "V" shape on the refill with the "V" shape at the top of your Dialpak. Snap the refill in place. You are ready to take pill "1." You should always begin your pill cycle with pill "1," as shown on the inner part of the refill ring.



5 Remove pill "1" by pushing down on the pill. The pill will come out through a hole in the back of the Dialpak.




6 Swallow the pill. You will take one pill each day. If you use a Sunday Start and you are taking the pill for the FIRST TIME, YOU MUST USE A BACK-UP METHOD OF BIRTH CONTROL FOR THE FIRST 7 DAYS. If you use a Day 1 Start, you are protected from becoming pregnant as soon as you take your first pill.



7 Wait 24 hours to take your next pill. To take pill "2," turn the dial on your Dialpak to the next day. Continue to take one pill each day until all the pills have been taken.



8 Take your pill at the same time every day. It is important to take the correct pill each day and not miss any pills. To help you remember, take your pill at the same time as another daily activity, like turning off your alarm clock or brushing your teeth.



9 When your refill is empty, keep your Dialpak case. You will start a new refill on the day after pill "28."



10 Turn the dial to the pill "1" position to remove the empty refill and insert a new refill. THE FIRST PILL IN EVERY REFILL WILL ALWAYS BE TAKEN ON THE SAME DAY OF THE WEEK, NO MATTER WHEN YOUR NEXT PERIOD STARTS.

DETAILED PATIENT LABELING

PLEASE NOTE: This labeling is revised from time to time as important new medical information becomes available. Therefore, please review this labeling carefully.

ORTHO TRI-CYCLEN® Lo Tablets

Each white tablet contains 0.180 mg norgestimate and 0.025 mg ethinyl estradiol.

Each light blue tablet contains 0.215 mg norgestimate and 0.025 mg ethinyl estradiol.

Each dark blue tablet contains 0.250 mg norgestimate and 0.025 mg ethinyl estradiol.

Each dark green tablet contains inert ingredients.

INTRODUCTION

Any woman who considers using oral contraceptives (the birth control pill or the pill) should understand the benefits and risks of using this form of birth control. This patient labeling will give you much of the information you will need to make this decision and will also help you determine if you are at risk of developing any of the serious side effects of the pill. It will tell you how to use the pill properly so that it will be as effective as possible. However, this labeling is not a replacement for a careful discussion between you and your healthcare professional. You should discuss the information provided in this labeling with him or her, both when you first start taking the pill and during your revisits. You should also follow your healthcare professional's advice with regard to regular check-ups while you are on the pill.

EFFECTIVENESS OF ORAL CONTRACEPTIVES FOR CONTRACEPTION

Oral contraceptives or "birth control pills" or "the pill" are used to prevent pregnancy and are more effective than most other non-surgical methods of birth control. When taken correctly without missing any pills, oral contraceptives are highly effective; however, typical failure rates are 5% per year. The chance of becoming pregnant increases with each missed pill during a menstrual cycle.

In comparison, typical failure rates for other non-surgical methods of birth control during the first year of use are as follows:

Implant: <1%	Male sterilization: <1%
Injection: <1%	Cervical Cap with spermicide: 20 to 40%
IUD: 1 to 2%	Condom alone (male): 14%
Diaphragm with spermicides: 20%	Condom alone (female): 21%
Spermicides alone: 26%	Periodic abstinence: 25%
Vaginal sponge: 20 to 40%	No methods: 85%
Female sterilization: <1%	Withdrawal: 19%

WHO SHOULD NOT TAKE ORAL CONTRACEPTIVES

Cigarette smoking increases the risk of serious cardiovascular side effects from oral contraceptive use. This risk increases with age and with heavy smoking (15 or more cigarettes per day) and is quite marked in women over 35 years of age. Women who use oral contraceptives are strongly advised not to smoke.

Some women should not use the pill. You should not use the pill if you have any of the following conditions:

- A history of heart attack or stroke
- Blood clots in the legs (thrombophlebitis), lungs (pulmonary embolism), or eyes
- A history of blood clots in the deep veins of your legs
- Chest pain (angina pectoris)
- Known or suspected breast cancer or cancer of the lining of the uterus, cervix or vagina
- Unexplained vaginal bleeding (until a diagnosis is reached by your healthcare professional)
- Yellowing of the whites of the eyes or of the skin (jaundice) during pregnancy or during previous use of the pill
- Liver tumor (benign or cancerous)
- Known or suspected pregnancy
- If you plan to have surgery with prolonged bedrest

Tell your healthcare professional if you have ever had any of these conditions. Your healthcare professional can recommend a safer method of birth control.

OTHER CONSIDERATIONS BEFORE TAKING ORAL CONTRACEPTIVES

Tell your healthcare professional if you have or have had:

- Breast nodules, fibrocystic disease of the breast, an abnormal breast x-ray or mammogram
- Diabetes
- Elevated cholesterol or triglycerides
- High blood pressure
- Migraine or other headaches or epilepsy

- Mental depression
- Gallbladder, liver, heart or kidney disease
- History of scanty or irregular menstrual periods

Women with any of these conditions should be checked often by their healthcare professional if they choose to use oral contraceptives.

Also, be sure to inform your healthcare professional if you smoke or are on any medications.

RISKS OF TAKING ORAL CONTRACEPTIVES

1. Risk of Developing Blood Clots

Blood clots and blockage of blood vessels are one of the most serious side effects of taking oral contraceptives and can cause death or serious disability. In particular, a clot in the legs can cause thrombophlebitis and a clot that travels to the lungs can cause a sudden blocking of the vessel carrying blood to the lungs. Rarely, clots occur in the blood vessels of the eye and may cause blindness, double vision, or impaired vision.

If you take oral contraceptives and need elective surgery, need to stay in bed for a prolonged illness or injury or have recently delivered a baby, you may be at risk of developing blood clots. You should consult your healthcare professional about stopping oral contraceptives four weeks before surgery and not taking oral contraceptives for two weeks after surgery or during bed rest. You should also not take oral contraceptives soon after delivery of a baby. It is advisable to wait for at least four weeks after delivery if you are not breastfeeding. If you are breastfeeding, you should wait until you have weaned your child before using the pill. (See also the section on Breastfeeding in General Precautions.)

The risk of circulatory disease in oral contraceptive users may be higher in users of high-dose pills and may be greater with longer duration of oral contraceptive use. In addition, some of these increased risks may continue for a number of years after stopping oral contraceptives. The risk of abnormal blood clotting increases with age in both users and nonusers of oral contraceptives, but the increased risk from the oral contraceptive appears to be present at all ages. For women aged 20 to 44 it is estimated that about 1 in 2,000 using oral contraceptives will be hospitalized each year because of abnormal clotting. Among nonusers in the same age group, about 1 in 20,000 would be hospitalized each year. For oral contraceptive users in general, it has been estimated that in women between the ages of 15 and 34 the risk of death due to a circulatory disorder is about 1 in 12,000 per year, whereas for nonusers the rate is about 1 in 50,000 per year. In the age group 35 to 44, the risk is estimated to be about 1 in 2,500 per year for oral contraceptive users and about 1 in 10,000 per year for nonusers.

2. Heart Attacks and Strokes

Oral contraceptives may increase the tendency to develop strokes (stoppage or rupture of blood vessels in the brain) and angina pectoris and heart attacks (blockage of blood vessels in the heart). Any of these conditions can cause death or serious disability.

Smoking greatly increases the possibility of suffering heart attacks and strokes. Furthermore, smoking and the use of oral contraceptives greatly increase the chances of developing and dying of heart disease.

3. Gallbladder Disease

Oral contraceptive users probably have a greater risk than nonusers of having gallbladder disease, although this risk may be related to pills containing high doses of estrogens.

4. Liver Tumors

In rare cases, oral contraceptives can cause benign but dangerous liver tumors. These benign liver tumors can rupture and cause fatal internal bleeding. In addition, some studies report an increased risk of developing liver cancer. However, liver cancers are rare.

5. Cancer of the Reproductive Organs and Breasts

Various studies give conflicting reports on the relationship between breast cancer and oral contraceptive use. Oral contraceptive use may slightly increase your chance of having breast cancer diagnosed, particularly after using hormonal contraceptives at a younger age. After you stop using hormonal contraceptives, the chances of having breast cancer diagnosed begin to go back down. You should have regular breast examinations by a healthcare professional and examine your own breasts monthly. Tell your healthcare professional if you have a family history of breast cancer or if you have had breast nodules or an abnormal mammogram. Women who currently have or have had breast cancer should not use oral contraceptives because breast cancer is usually a hormone-sensitive tumor.

Some studies have found an increase in the incidence of cancer of the cervix in women who use oral contraceptives. However, this finding may be related to factors other than the use of oral contraceptives. There is insufficient evidence to rule out the possibility that the pill may cause such cancers.

ESTIMATED RISK OF DEATH FROM A BIRTH CONTROL METHOD OR PREGNANCY

All methods of birth control and pregnancy are associated with a risk of developing certain diseases which may lead to disability or death. An estimate of the number of deaths associated with different methods of birth control and pregnancy has been calculated and is shown in the following table.

Table 4: Annual Number of Birth-Related or Method-Related Deaths Associated with Control of Fertility Per 100,000 Nonsterile Women, By Fertility-Control Method and According to Age

Method of control and outcome	15-19	20-24	25-29	30-34	35-39	40-44
No fertility-control methods*	7.0	7.4	9.1	14.8	25.7	28.2
Oral contraceptives nonsmoker**	0.3	0.5	0.9	1.9	13.8	31.6
Oral contraceptives smoker**	2.2	3.4	6.6	13.5	51.1	117.2
IUD**	0.8	0.8	1.0	1.0	1.4	1.4
Condom*	1.1	1.6	0.7	0.2	0.3	0.4
Diaphragm/spermicide*	1.9	1.2	1.2	1.3	2.2	2.8
Periodic abstinence*	2.5	1.6	1.6	1.7	2.9	3.6

* Deaths are birth-related

** Deaths are method-related

Adapted from H.W. Ory, Family Planning Perspectives, Ref. #35.

In the above table, the risk of death from any birth control method is less than the risk of childbirth, except for oral contraceptive users over the age of 35 who smoke and pill users over the age of 40 even if they do not smoke. It can be seen in the table that for women aged 15 to 39, the risk of death was highest with pregnancy (7-26 deaths per 100,000 women, depending on age). Among pill users who do not smoke, the risk of death was always lower than that associated with pregnancy for any age group, although over the age of 40, the risk increases to 32 deaths per 100,000 women, compared to 28 associated with pregnancy at that age. However, for pill users who smoke and are over the age of 35, the estimated number of deaths exceeds those for other methods of birth control. If a woman is over the age of 40 and smokes, her estimated risk of death is four times higher (117/100,000 women) than the estimated risk associated with pregnancy (28/100,000 women) in that age group.

The suggestion that women over 40 who do not smoke should not take oral contraceptives is based on information from older, higher-dose pills. An Advisory Committee of the FDA discussed this issue in 1989 and recommended that the benefits of low-dose oral contraceptive use by healthy, non-smoking women over 40 years of age may outweigh the possible risks. Older women, as all women, who take oral contraceptives, should take an oral contraceptive which contains the least amount of estrogen and progestogen that is compatible with the individual patient needs.

WARNING SIGNALS

If any of these adverse effects occur while you are taking oral contraceptives, call your healthcare professional immediately:

- Sharp chest pain, coughing of blood, or sudden shortness of breath (indicating a possible clot in the lung)
- Pain in the calf (indicating a possible clot in the leg)
- Crushing chest pain or heaviness in the chest (indicating a possible heart attack)
- Sudden severe headache or vomiting, dizziness or fainting, disturbances of vision or speech, weakness, or numbness in an arm or leg (indicating a possible stroke)
- Sudden partial or complete loss of vision (indicating a possible clot in the eye)
- Breast lumps (indicating possible breast cancer or fibrocystic disease of the breast; ask your healthcare professional to show you how to examine your breasts)
- Severe pain or tenderness in the stomach area (indicating a possibly ruptured liver tumor)
- Difficulty in sleeping, weakness, lack of energy, fatigue, or change in mood (possibly indicating severe depression)
- Jaundice or a yellowing of the skin or eyeballs, accompanied frequently by fever, fatigue, loss of appetite, dark colored urine, or light colored bowel movements (indicating possible liver problems)

SIDE EFFECTS OF ORAL CONTRACEPTIVES

1. Vaginal Bleeding

Irregular vaginal bleeding or spotting may occur while you are taking the pills. Irregular bleeding may vary from slight staining between menstrual periods to breakthrough bleeding which is a flow much like a regular period. Irregular bleeding occurs most often during the first few months of oral contraceptive use, but may also occur after you have been taking the pill for some time. Such bleeding may be temporary

and usually does not indicate any serious problems. It is important to continue taking your pills on schedule. If the bleeding occurs in more than one cycle or lasts for more than a few days, talk to your healthcare professional.

2. Contact Lenses

If you wear contact lenses and notice a change in vision or an inability to wear your lenses, contact your healthcare professional.

3. Fluid Retention

Oral contraceptives may cause edema (fluid retention) with swelling of the fingers or ankles and may raise your blood pressure. If you experience fluid retention, contact your healthcare professional.

4. Melasma

A spotty darkening of the skin is possible, particularly of the face, which may persist.

5. Other Side Effects

Other side effects may include nausea and vomiting, change in appetite, headache, nervousness, depression, dizziness, loss of scalp hair, rash, vaginal infections, and allergic reactions.

If any of these side effects bother you, call your healthcare professional.

GENERAL PRECAUTIONS

1. Missed Periods and Use of Oral Contraceptives Before or During Early Pregnancy

There may be times when you may not menstruate regularly after you have completed taking a cycle of pills. If you have taken your pills regularly and miss one menstrual period, continue taking your pills for the next cycle but be sure to inform your healthcare professional. If you have not taken the pills daily as instructed and missed a menstrual period, you may be pregnant. If you missed two consecutive menstrual periods, you may be pregnant. Check with your healthcare professional immediately to determine whether you are pregnant. Stop taking oral contraceptives if pregnancy is confirmed.

There is no conclusive evidence that oral contraceptive use is associated with an increase in birth defects, when taken inadvertently during early pregnancy. Previously, a few studies had reported that oral contraceptives might be associated with birth defects, but these findings have not been seen in more recent studies. Nevertheless, oral contraceptives should not be used during pregnancy. You should check with your healthcare professional about risks to your unborn child of any medication taken during pregnancy.

2. While Breastfeeding

If you are breastfeeding, consult your healthcare professional before starting oral contraceptives. Some of the drug will be passed on to the child in the milk. A few adverse effects on the child have been reported, including yellowing of the skin (jaundice) and breast enlargement. In addition, oral contraceptives may decrease the amount and quality of your milk. If possible, do not use oral contraceptives while breastfeeding. You should use another method of contraception since breastfeeding provides only partial protection from becoming pregnant and this partial protection decreases significantly as you breastfeed for longer periods of time. You should consider starting oral contraceptives only after you have weaned your child completely.

3. Laboratory Tests

If you are scheduled for any laboratory tests, tell your healthcare professional you are taking birth control pills. Certain blood tests may be affected by birth control pills.

4. Drug Interactions

Certain drugs may interact with birth control pills to make them less effective in preventing pregnancy or cause an increase in breakthrough bleeding. Such drugs include rifampin, drugs used for epilepsy such as barbiturates (for example, phenobarbital), topiramate (TOPAMAX®), carbamazepine (Tegretol® is one brand of this drug), or phenytoin (Dilantin® is one brand of this drug); phenylbutazone (Butazolidin® is one brand); certain drugs used in the treatment of HIV or AIDS; and possibly certain antibiotics. Medicine for pulmonary hypertension, such as bosentan (Tracleer®). Pregnancies and breakthrough bleeding have been reported by women who also used some form of the herbal supplement St. John's Wort while using combined hormonal contraceptives. Hormonal contraceptives may interact with lamotrigine (LAMICTAL®), an anticonvulsant used for epilepsy. This may increase the risk of seizures so your healthcare professional may need to adjust the dose of lamotrigine. You may need to use additional contraception when you take other products which can make oral contraceptives less effective. Be sure to tell your healthcare professional if you are taking or start taking any medications while taking birth control pills.

5. Sexually Transmitted Diseases

ORTHO TRI-CYCLEN® Lo (like all oral contraceptives) is intended to prevent pregnancy. Oral contraceptives do not protect against transmission of HIV (AIDS) and other sexually transmitted diseases such as chlamydia, genital herpes, genital warts, gonorrhea, hepatitis B, and syphilis.

HOW TO TAKE THE PILL

IMPORTANT POINTS TO REMEMBER

BEFORE YOU START TAKING YOUR PILLS:

1. **BE SURE TO READ THESE DIRECTIONS:**
Before you start taking your pills.
Anytime you are not sure what to do.
2. **THE RIGHT WAY TO TAKE THE PILL IS TO TAKE ONE PILL EVERY DAY AT THE SAME TIME.**
If you miss pills you could get pregnant. This includes starting the pack late.
The more pills you miss, the more likely you are to get pregnant.
3. **MANY WOMEN HAVE SPOTTING OR LIGHT BLEEDING, OR MAY FEEL SICK TO THEIR STOMACH DURING THE FIRST 1-3 PACKS OF PILLS.** If you feel sick to your stomach, do not stop taking the pill. The problem will usually go away. If it doesn't go away, check with your healthcare professional.
4. **MISSING PILLS CAN ALSO CAUSE SPOTTING OR LIGHT BLEEDING,** even when you make up these missed pills.
On the days you take 2 pills to make up for missed pills, you could also feel a little sick to your stomach.
5. **IF YOU HAVE VOMITING OR DIARRHEA, or IF YOU TAKE SOME MEDICINES,** including some antibiotics, your pills may not work as well.
Use a back-up method (such as condoms or spermicides) until you check with your healthcare professional.
6. **IF YOU HAVE TROUBLE REMEMBERING TO TAKE THE PILL,** talk to your healthcare professional about how to make pill-taking easier or about using another method of birth control.
7. **IF YOU HAVE ANY QUESTIONS OR ARE UNSURE ABOUT THE INFORMATION IN THIS LEAFLET,** call your healthcare professional.

BEFORE YOU START TAKING YOUR PILLS

1. **DECIDE WHAT TIME OF DAY YOU WANT TO TAKE YOUR PILL.**
It is important to take it at about the same time every day.
2. The pill pack has 21 white, light blue, and dark blue "active" pills (with hormones) to take for 3 weeks. This is followed by 1 week of dark green "reminder" pills (without hormones).
3. **ALSO FIND:**
 - 1) where on the pack to start taking pills,
 - 2) in what order to take the pills.

CHECK PICTURE OF PILL PACK AND ADDITIONAL INSTRUCTIONS FOR USING THIS PACKAGE IN THE BRIEF SUMMARY PATIENT PACKAGE INSERT.

4. **BE SURE YOU HAVE READY AT ALL TIMES:**
ANOTHER KIND OF BIRTH CONTROL (such as condoms or spermicide) to use as a back-up method in case you miss pills.
AN EXTRA, FULL PILL PACK.

WHEN TO START THE FIRST PACK OF PILLS

You have a choice of which day to start taking your first pack of pills. ORTHO TRI-CYCLEN® Lo is available in the DIALPAK® Tablet Dispenser which is preset for a Sunday Start. Day 1 Start is also provided. Decide with your healthcare professional which is the best day for you. Pick a time of day which will be easy to remember.

Sunday Start:

Take the first white "active" pill of the first pack on the Sunday after your period starts, even if you are still bleeding. If your period begins on Sunday, start the pack that same day.

Use another method of birth control (such as condoms or spermicide) as a back-up method if you have sex anytime from the Sunday you start your first pack until the next Sunday (7 days).

Day 1 Start:

Take the first white "active" pill of the first pack during the first 24 hours of your period.

You will not need to use a back-up method of birth control, since you are starting the pill at the beginning of your period.

WHAT TO DO DURING THE MONTH

1. **Take One Pill At The Same Time Every Day Until The Pack Is Empty**
Do not skip pills even if you are spotting or bleeding between monthly periods or feel sick to your stomach (nausea).
Do not skip pills even if you do not have sex very often.
2. **When You Finish A Pack Or Switch Your Brand Of Pills**
Start the next pack on the day after your last dark green "reminder" pill. Do not wait any days between packs.

WHAT TO DO IF YOU MISS PILLS

If you **MISS 1** white, light blue or dark blue "active" pill:

1. Take it as soon as you remember. Take the next pill at your regular time. This means you may take 2 pills in 1 day.

2. You do not need to use a back-up birth control method if you have sex.

If you **MISS 2** white or light blue “active” pills in a row in **WEEK 1 OR WEEK 2** of your pack:

1. Take 2 pills on the day you remember and 2 pills the next day.
2. Then take 1 pill a day until you finish the pack.
3. You **COULD BECOME PREGNANT** if you have sex in the 7 days after you miss pills. You **MUST** use another birth control method (such as condoms or spermicide) as a back-up method for those 7 days.

If you **MISS 2** dark blue “active” pills in a row in **THE 3RD WEEK**:

1. If you are a Sunday Starter:

Keep taking 1 pill every day until Sunday. On Sunday, **THROW OUT** the rest of the pack and start a new pack of pills that same day.

If you are a Day 1 Starter:

THROW OUT the rest of the pill pack and start a new pack that same day.

2. You may not have your period this month but this is expected. However, if you miss your period 2 months in a row, call your healthcare professional because you might be pregnant.
3. You **COULD BECOME PREGNANT** if you have sex in the 7 days after you miss pills. You **MUST** use another birth control method (such as condoms or spermicide) as a back-up method for those 7 days.

If you **MISS 3 OR MORE** white, light blue or dark blue “active” pills in a row (during the first 3 weeks):

1. If you are a Sunday Starter:

Keep taking 1 pill every day until Sunday. On Sunday, **THROW OUT** the rest of the pack and start a new pack of pills that same day.

If you are a Day 1 Starter:

THROW OUT the rest of the pill pack and start a new pack that same day.

2. You may not have your period this month but this is expected. However, if you miss your period 2 months in a row, call your healthcare professional because you might be pregnant.
3. You **COULD BECOME PREGNANT** if you have sex in the 7 days after you miss pills. You **MUST** use another birth control method (such as condoms or spermicide) as a back-up method for those 7 days.

If you forget any of the 7 dark green “reminder” pills in Week 4: **THROW AWAY** the pills you missed.

Keep taking 1 pill each day until the pack is empty.

You do not need a back-up method.

FINALLY, IF YOU ARE STILL NOT SURE WHAT TO DO ABOUT THE PILLS YOU HAVE MISSED:

Use a **BACK-UP METHOD** anytime you have sex.

KEEP TAKING ONE “ACTIVE” PILL EACH DAY until you can reach your healthcare professional.

PREGNANCY DUE TO PILL FAILURE

When taken correctly without missing any pills, oral contraceptives are highly effective; however the typical failure rate of large numbers of pill users is 5% per year when women who miss pills are included. If failure does occur, the risk to the fetus is minimal.

PREGNANCY AFTER STOPPING THE PILL

There may be some delay in becoming pregnant after you stop using oral contraceptives, especially if you had irregular menstrual cycles before you used oral contraceptives. It may be advisable to postpone conception until you begin menstruating regularly once you have stopped taking the pill and desire pregnancy.

There does not appear to be any increase in birth defects in newborn babies when pregnancy occurs soon after stopping the pill.

OVERDOSAGE

Serious ill effects have not been reported following ingestion of large doses of oral contraceptives by young children. Overdosage may cause nausea and withdrawal bleeding in females. In case of overdosage, contact your healthcare professional or pharmacist.

OTHER INFORMATION

Your healthcare professional will take a medical and family history before prescribing oral contraceptives and will examine you. The physical examination may be delayed to another time if you request

it and the healthcare professional believes that it is a good medical practice to postpone it. You should be reexamined at least once a year. Be sure to inform your healthcare professional if there is a family history of any of the conditions listed previously in this leaflet. Be sure to keep all appointments with your healthcare professional, because this is a time to determine if there are early signs of side effects of oral contraceptive use.

Do not use the drug for any condition other than the one for which it was prescribed. This drug has been prescribed specifically for you; do not give it to others who may want birth control pills.

HEALTH BENEFITS FROM ORAL CONTRACEPTIVES

In addition to preventing pregnancy, use of combination oral contraceptives may provide certain benefits. They are:

- menstrual cycles may become more regular
- blood flow during menstruation may be lighter and less iron may be lost. Therefore, anemia due to iron deficiency is less likely to occur
- pain or other symptoms during menstruation may be encountered less frequently
- ectopic (tubal) pregnancy may occur less frequently
- noncancerous cysts or lumps in the breast may occur less frequently
- acute pelvic inflammatory disease may occur less frequently
- oral contraceptive use may provide some protection against developing two forms of cancer: cancer of the ovaries and cancer of the lining of the uterus

If you want more information about birth control pills, ask your healthcare professional or pharmacist. They have a more technical leaflet called the Professional Labeling, which you may wish to read. The professional labeling is also published in a book entitled *Physicians' Desk Reference*, available in many bookstores and public libraries.



Mfd. for:
Ortho Women's Health & Urology,
Division of Ortho-McNeil-Janssen Pharmaceuticals, Inc.
Raritan, New Jersey 08869

Mfd. by:
Janssen Ortho, LLC
Manati, Puerto Rico 00674
© Ortho-McNeil-Janssen Pharmaceuticals, Inc. 1998
Revised August 2008

635-50-951-5

SUPPLEMENT TO
OBG
MANAGEMENT

