

Problems in Family Practice

Joint Pain of Recent Onset

Cecil O. Samuelson, Jr, MD
John R. Ward, MD
Salt Lake City, Utah

A joint pain of recent onset is a common and often challenging problem to the family physician. The many etiologies of acute arthritis require a systematic approach to a careful definition of the problem which is essential to safe and effective therapy. The proper evaluation of joint pain of recent onset requires a detailed history and physical examination as well as a rationally selected battery of laboratory studies. This paper provides the busy physician with a specific approach to the diagnosis of acute joint pain as the basis for precise management.

For the busy family physician, acute joint pain, like fever, is a frequently encountered and often challenging problem. Because joint pain is a symptom of underlying pathology or disease, the diagnosis of "arthritis" is no more precise than "fever" or "anemia." While it is often possible to find an immediate cause and thus correct the problem, the many etiologies of acute arthritis must include a systematic approach to a precise definition of the problem. The purpose of this paper is to assist the physician in developing a method to successfully and expeditiously deal with the constellation of problems presenting as acute joint pain.

Like "fever" and "anemia," arthritis may imply different pathological conditions to individual physicians. While precise definitions are difficult, the term arthralgia is a subjective symptom which denotes joint pain, discomfort, or tenderness, without objective findings upon examination. Arthritis describes the presence of objective joint inflammation such as soft tissue swelling, joint effusion, heat, or redness. Acute arthritis is a confusing term and may denote the severity or course of arthritis. Thus, a more precise description might be arthritis of recent onset. This definition should be reserved for inflammation of the joints alone, and is distinct from involvement of the periarticular tissues as is found in bursitis or tendinitis.

While many disease processes may present as, or be associated with, arthritis of recent onset, the approach to the problem is similar whatever the etiology. Usually, the most important information is historical. By knowing the setting of the problem, the pattern of joint involvement, and the associated manifestations, the etiology can often be narrowed to the point where

the physical examination and laboratory data are necessary only to confirm the diagnosis. In other instances, only the most thorough and thoughtful evaluation will reveal the cause. Some patients will defy early diagnosis because of an inability to muster sufficient data to establish or exclude disease. Thus, the problem may remain unresolved and demand careful and periodic re-evaluation.

In many patients with joint pain of recent onset, the physician will immediately identify the nature of the problem. For example, the preschool child may refuse to use an arm which he protects by holding the elbow flexed and the forearm pronated. Painful limitation of forearm supination is found and a diagnosis of "pulled" or dislocated elbow is established. A patient may present with a "sprain of the wrist." There is marked local tenderness in the region of the anatomic "snuff box" with pain on dorsiflexion and radial deviation. If x-rays show a fracture of the scaphoid (navicular) bone, the diagnosis is clear. If not, repeat x-rays in one to two weeks will usually reveal the fracture. Similarly, stress fractures in the metatarsals may present as local pain and tenderness, often with swelling. Early x-rays may or may not reveal the fracture in the neck of a metatarsal. Older or adolescent children may sustain lateral dislocations of the patella. The history is characteristic with sudden sharp pain, the knee gives way and the patient falls. Physical examination reveals a grossly swollen knee due to hemarthrosis. If the patella is found to be laterally displaced, the diagnosis is evident.

The preceding examples are cited to emphasize the importance of traumatic lesions in monarticular joint disease. However, we have chosen to exclude injuries in this discussion because the cause is usually obvious. Thus, non-traumatic diseases presenting with arthritis of recent onset are usually systemic in nature and the history and physical examination should be general and thorough. To restrict the evaluation to the patient's chief complaints incurs the risk of misdiagnosis. Subsequent laboratory and x-ray evaluations should be dictated by a well-conceived "game plan" which clearly identifies the priorities for action and overrides the "shotgun" approach. Finally, therapeutic trials

From the Department of Internal Medicine, Division of Arthritis, University of Utah College of Medicine, Salt Lake City, Utah. Reprinted with updating where appropriate from *The Journal of Family Practice*, Volume 1 Number 3/4, 1974. Requests for reprints should be addressed to Dr. C. O. Samuelson, Jr., Division of Arthritis, University of Utah Medical Center, 50 North Medical Drive, Salt Lake City, UT 84132.

are rarely definitive and should not be used in an attempt to identify an etiology.

Subjective Data

Success or failure in making the correct diagnosis is often determined by the quality of the historical data. It is essential to elicit a clear pattern of symptoms. Is the joint pain confined to a single or to multiple joints? Which joints are involved? Is the arthritis migratory? Are there systemic features to the illness? The diagnosis is frequently revealed by understanding associated diseases, epidemiological patterns, life-styles, and familial disease processes.

When a young person presents with migratory polyarthritis, a history of sore throat, fever, and perhaps choreiform movements, the diagnosis of rheumatic fever seems probable.¹ It must be remembered, however, that while the large joints of the extremities are most frequently affected and effusions, if present, are usually transient, multiple joints may be affected simultaneously. Also, the arthritis may be persistent and non-migratory. In the unusual presentation, it may be even more important to obtain the appropriate history with regard to the occurrence of carditis, skin rash, and nodules as well as the more common systemic features of acute rheumatic fever.

Subtle historical data may be helpful in identifying the child in the same age group who has juvenile rheumatoid arthritis.² It is important to remember that juvenile rheumatoid arthritis may present in varied ways and differs significantly in many respects from adult rheumatoid arthritis. Among the most important differences are the high frequency of severe systemic disease, persistent pauciarticular manifestations, and ocular involvement. Although it can occur at any age, the most frequent ages of onset are between 2 to 5 and 9 to 12 years. While usually polyarticular at the onset, 30 percent of cases are monoarticular with the knee being the most common joint involved. Frequently, young children will deny pain even when the joint is obviously involved. On the other hand, arthralgias, transient joint swelling, and constitutional symptoms may be the sequelae of rubella immunization or an otherwise non-consequential viral infection.³ Only a

careful history may provide the clues to the proper diagnosis.

If a patient presenting with the sudden onset of arthritis is a promiscuous "go-go dancer" with perhaps associated fever and skin rash, the historical probability of gonococcal arthritis may be established,⁴ but is enhanced if urethral or vaginal discharge, tenosynovitis, and multiple joint involvement are also present. The arthritis of Australian-antigen-associated hepatitis⁵ should not be overlooked in those possibly exposed, including drug abusers, especially when they also have fever and urticarial skin rash. The same historical features are of course equally compatible with systemic lupus erythematosus,⁶ but it is helpful to know if the rash is photosensitive and accompanied by such features as Raynaud phenomenon, pleuritic chest pain, and even cardiac or neurological symptoms. While it is not practical to list all diseases associated with acute joint pain and arthritis, Table 1 includes most pathological processes in which acute joint pain may occur.

Abnormalities in only the soft tissues approximating the joints may be reported by the patient as joint pain. Thus, inflammation of the bicipital tendon may be referred to as shoulder pain, synovitis in the olecranon bursa as elbow "arthritis," and fluid in the prepatellar bursa as "housemaid's knee." It must also be remembered that nerve entrapment syndromes frequently present with pain or numbness referable to the adjacent bones and joints and may also occur suddenly, at least historically. Thus, the carpal tunnel syndrome should be considered when the patient reports a painful hand that "goes to sleep," is weak, and drops objects. Careful gathering and evaluation of the historical clues presenting with each individual problem are most often the key to proper diagnosis.

While the particular joint involved may be most helpful in the differential diagnosis, it should be remembered that no joints are immune from less common afflictions. Even though the first metatarsophalangeal joint is the classical location for podagra or acute gouty arthritis, it is by no means diagnostic in any patient. Certainly, if the patient is a menstruating female with a negative family history for gout, it is highly unlikely that she has

gouty arthritis, whatever joint is involved.

Objective Data

A complete physical examination is essential as multiple diagnostic clues are often available if sought carefully. Also, the systemic features of an illness may be more specific than the articular manifestations alone. As stated above, skin rashes are common in some entities but rare in others, such as in gouty or rheumatoid arthritis where tophi or nodules may be found. Similarly, conjunctivitis, retinal changes, hepatosplenomegaly, heart murmurs, peripheral neuropathy, mononeuritis multiplex, petechial hemorrhages, and lesions consistent with vasculitis, as well as others, may narrow the diagnostic spectrum when present. As the joints are examined, it is useful to remember that synovial membranes and periarticular tissues are primarily involved in such processes as rheumatoid, infectious, and gouty arthritis, while the bones and cartilage are most often affected by hereditary, neoplastic, and degenerative processes.

A detailed description of the technique for the examination of each joint is beyond the scope of this paper. Several points have general applicability and should be kept in mind in the examination of the patient with joint pain of recent onset. The range of motion of each joint should be assessed both actively and passively and compared with normal. If the examiner is aware of the range of motion of his/her own joints, a useful comparison is at hand. Inspection of each joint should include the identification of anatomical landmarks, skin lesions and color, swelling, deformities, periarticular muscle atrophy, tenosynovitis, and abnormal positioning or posture. Palpation should reveal the degree of tenderness, temperature, type of swelling, and localized swellings such as ganglia, nodules, or tophi, as well as crepitus or grating. Synovial proliferation presents as a doughy enlargement in the areas where the joint capsule and synovial reflections are near the surface. A joint effusion may be detected by a producible fluid wave or a ballotable patella in an accessible joint such as the knee, but may be impossible to demonstrate in the hip or shoulder short of arthrocentesis. These types of

Table 1. Causes of Acute Joint Pain

1. **Infection:** Seen most often in the very young, the very old, and the debilitated. Also frequently seen in rheumatoid arthritis, collagen vascular disease, and immunosuppressed patients.
 - a. Bacterial — gonococcal, staphylococcal, streptococcal, others
 - b. Viral — rubella, mumps, measles, infectious hepatitis, others
 - c. Mycobacteria, fungi, rickettsiae, others
2. **Gout:** Abnormal purine metabolism resulting in hyperuricemia and urate crystal deposition, usually monarticular and in older males. "Secondary" gout may result from many causes such as hematopoietic or neoplastic diseases, drug therapy, renal failure, etc.
3. **Pseudogout:** Deposition of calcium pyrophosphate dihydrate crystals in joints of older patients. Acute "gouty" episodes are usually in larger joints, may become chronic. Chondrocalcinosis is usually present, but not invariably. May also mimic rheumatoid arthritis and degenerative joint disease.
4. **Rheumatoid Arthritis:** Usually a polyarthritis (may be monarticular initially) of symmetrical peripheral joints, more frequently found in females.
5. **Juvenile Rheumatoid Arthritis:** May present as monarticular, oligoarticular, or polyarticular disease which may be associated with severe systemic disease and is found in both boys and girls.
6. **Acute Rheumatic Fever:** An inflammatory disease occurring as a sequel to Group A streptococcal infection, usually between ages 5 and 15.
7. **Systemic Lupus Erythematosus:** A chronic inflammatory disease of unknown etiology that may affect many organ systems and is usually seen in young females.
8. **Reiter's syndrome:** Characterized by urethritis, arthritis, conjunctivitis, circinate balanitis, and keratoderma blennorrhagica in young adult males. Incomplete syndrome is possible with proper laboratory confirmation.
9. **Ankylosing Spondylitis:** Chronic, progressive arthritis of the sacroiliac joints and axial skeleton, usually found in males in their second or third decade. May also involve peripheral joints.
10. **Sarcoidosis:** Usually associated with erythema nodosum and is often migratory, polyarticular, and involves large joints.
11. **Palindromic Rheumatism:** Recurrent, acute, and self-limiting arthritis attacks.
12. **Intermittent Hydrarthrosis:** Recurrent joint effusions, usually of the knee, which occur at regular intervals and are idiopathic.
13. **Inflammatory Bowel Disease:** Regional Enteritis (RE), Ulcerative Colitis (UC): Found in 5 percent of RE and 20 percent of UC cases and presents as either spondylitis or as an acute peripheral polyarthritis involving large joints.
14. **Erythema Nodosum:** An acute, self-limiting condition of multiple red, tender nodules in the skin of the lower extremities, most common in females and of multiple etiologies.
15. **Hyperparathyroidism:** Excessive parathormone production has been associated with arthritis of multiple types and is most common in adult females.
16. **Hypothyroidism:** Thyroid hormone deficiency is associated with arthralgias and myalgias and occasionally arthritis. Most frequently in middle-aged females and many have carpal tunnel syndrome.
17. **Leukemia:** Hematologic neoplasia with arthritis due to joint hemorrhage or infiltration and found most frequently in lower extremities with severe pain.
18. **Polyarteritis Nodosa:** Arthralgias associated with multiple systems involvement secondary to diffuse, but focal, panarteritis and found most frequently in males. True arthritis is rare.
19. **Progressive Systemic Sclerosis (Scleroderma):** Peripheral arthritis most common in females with involved joints usually related to affected skin.
20. **Psoriasis:** Ten percent of patients with this skin disease develop a peripheral small joint arthritis.
21. **Pulmonary Disease (Hypertrophic Pulmonary Osteoarthropathy):** The syndrome of digital clubbing, synovitis, and periostitis most often seen in middle-aged men with bronchogenic carcinoma.
22. **Shoulder-Hand Syndrome:** Painful disability of the shoulder associated with pain, swelling, and vasomotor changes in the hand, most frequently seen in adults with myocardial infarction, cervical spondylosis, trauma, or no apparent cause.
23. **Agammaglobulinemia:** Thirty percent of patients with this rare disorder develop a polyarticular, peripheral arthritis, which is most common in children.
24. **Amyloidosis:** A polyarthritis resembling rheumatoid may occur secondary to amyloid infiltration of the synovium, usually in older patients.
25. **Mixed Connective Tissue Disease:** An overlap syndrome with variable features, consistent with scleroderma, rheumatoid arthritis, and systemic lupus erythematosus.
26. **Henoch-Schönlein Purpura:** Arthritis associated with purpura in children, usually affecting knees and ankles, and often following upper respiratory infection.
27. **Hyperlipidemia:** Periarticular symptoms and arthritis associated with xanthomas may be associated with the various hyperlipoproteinemias and follow primary incidence and prevalence patterns.
28. **Hemophilia:** Hemarthrosis occurs in 80 to 90 percent of patients with hemophilia and affects the knees, elbows, and ankles most frequently.
29. **Hemochromatosis:** Most frequent in males over 40 with synovitis, and bony enlargement most common in the small joints of the hands.
30. **Sickle Cell Anemia:** Crises may be accompanied by intensely painful polyarthralgias and arthritis.
31. **Ochronosis (Alkaptonuria):** An accumulation of homogentisic acid secondary to the deficiency of homogentisic acid oxidase, resulting in deposition of a pigmented polymer in cartilage which leads to a degenerative arthritis.
32. **Whipple's Disease:** Diarrhea, malabsorption, and constitutional symptoms may be associated with migratory arthritis in middle-aged men.
33. **Behcet Syndrome:** The triad of aphthous stomatitis, genital ulceration, and iritis may be associated with polyarthritis, usually involving the knee and central nervous system abnormalities and is more common in males.
34. **Multicentric Reticulohistiocytosis:** Destructive polyarthritis and nodular skin lesions most common in females with histologic examination revealing infiltration of tissues by giant cells with foamy cytoplasm.
35. **Familial Mediterranean Fever:** Recurrent attacks of fever and monarticular arthritis in young people of Mediterranean extraction.
36. **Neoplasms of Joints:** Pigmented villonodular synovitis, others: rare, usually present in a single joint with swelling and/or pain and may require surgical exploration for diagnosis.
37. **Serum Sickness (Drug Reactions):** Acute onset of arthralgias, myalgias, urticaria, gastrointestinal, and constitutional symptoms, 7 to 12 days after injection of foreign serum or drug, usually penicillin, but occurs with many drugs.

Table 2. Laboratory Studies Helpful in Arthritis of Recent Onset

A. Complete blood count

1. Anemia
 - a. Anemia of chronic disease in rheumatoid arthritis, infection
 - b. Hemolysis — systemic lupus erythematosus
2. Erythrocytosis — gout
3. Elevated white cell count — infection, leukemia
4. Depressed white cell count — systemic lupus erythematosus, Felty syndrome
5. Thrombocytopenia — systemic lupus erythematosus

B. Westergren erythrocyte sedimentation rate

1. Elevated — any inflammatory process
2. Normal — degenerative arthritis, trauma

C. Serological tests

1. Serological tests for syphilis — may be false positive in systemic lupus erythematosus, rheumatoid arthritis
2. Antinuclear antibodies — systemic lupus erythematosus, other connective tissue diseases
 - a. LE cell test
 - b. Fluorescent antibody tests — patterns may be helpful
 1. Speckled — progressive systemic sclerosis
 2. Peripheral (shaggy) — systemic lupus erythematosus
 3. Nucleolar — progressive systemic sclerosis
 4. Homogenous — systemic lupus erythematosus
 - c. Anti-DNA — systemic lupus erythematosus
 - d. Extractable nuclear antigen — mixed connective tissue disease
3. Rheumatoid factor — rheumatoid arthritis, frequently negative early
4. Streptococcal antibody tests (anti-streptolysin-O, anti-hyaluronidase, anti-streptokinase, others) — rheumatic fever
5. Gonococcal antibodies — gonococcal arthritis

D. Cultures in addition to synovial fluid — infectious arthritis

1. Blood
2. Skin lesions
3. Urethra, rectum, pharynx, cervix, vagina

E. Urinalysis — systemic lupus erythematosus, Henoch-Schönlein purpura, alkaptonuria

F. Uric acid — elevated in gouty arthritis, also may be elevated on a familial basis and secondary to drugs, renal disease, others

G. Chest x-ray — sarcoidosis, systemic lupus erythematosus

H. Electrocardiogram — acute rheumatic fever, systemic lupus erythematosus, pericarditis, ankylosing spondylitis

I. Serum complement

1. Elevated — rheumatoid arthritis
2. Depressed — systemic LE

J. Liver function tests, Australian-antigen — hepatitis, sarcoidosis

K. Joint x-rays — trauma, soft tissue swelling

L. Serum iron and total iron binding capacity, serum ferritin, bone marrow, Coombs — differentiate anemia of chronic disease from others

M. Sickle cell prep, Hemoglobin electrophoresis — sickle cell anemia, others

N. Synovial fluid

1. Gram stain — infection
2. Culture - infection
3. WBC and differential — inflammatory vs noninflammatory
4. Crystals — gout, pseudogout, steroid injections
5. RBC — hemorrhage, hemophilia, pigmented villinodular synovitis

O. Biopsy

Table 3. Synovial Fluid Findings

| | Crystals (polarized light exam) | WBC/mm ³ | PMN % | Culture |
|------------------------------|------------------------------------|--|---|---|
| Normal | None | Less than 200 | Less than 25% | Negative |
| Trauma | None | Approximates blood | Less than 5% | Negative |
| Infectious Arthritis | None | Usually greater than 100,000 but may be much lower | Usually greater than 75%, but may be less | Usually positive (except in gonococcal arthritis) |
| Gouty Arthritis | Positive monosodium urate | 200-100,000 | Usually greater than 50% | Negative |
| Pseudogout | Positive calcium pyrophosphate | 200-100,000 | Usually greater than 50% | Negative |
| Rheumatoid Arthritis | None | Variable usually 10,000-100,000 | Variable usually 40-80% | Negative |
| Acute Rheumatic Fever | None | Usually 10,000-30,000 | 10-50% | Negative |
| Systemic Lupus Erythematosus | None | Variable usually 10,000-100,000 | Variable usually 25-75% | Negative |

swelling should be distinguished from the firm, generally non-tender spurs or lipping of bony origin common in degenerative disease. Testing for instability of an affected joint is particularly important in trauma and progressive disease. Initially, it is perhaps of the most importance to look carefully for signs that suggest infection as this is the problem which usually requires the most rapid attention.

Laboratory studies are helpful and necessary but should not be ordered indiscriminately. Those that are helpful most often are found in Table 2. Initially, a complete blood count and a Westergren sedimentation rate should be obtained on each patient presenting with arthritis of recent onset. Those studies which will help answer questions raised in the history and physical examination should also be done. Rarely, if ever, are all of the tests mentioned in Table 2 indicated in the

same patient. The "shotgun" approach is not only expensive for the patient, but suggests faulty reasoning by the physician. Rarely is a uric acid determination helpful in children or young women and lupus erythematosus (LE) preps usually contribute nothing in the older man presenting with classical gouty arthritis on clinical grounds. A few carefully chosen studies considered in the light of a complete history and physical examination are more helpful than a battery of tests used alone.

A properly obtained and correctly examined synovial fluid specimen can be diagnostic for hemarthrosis, septic arthritis, gout, and pseudogout as well as being helpful in other processes. Joint aspiration is safe and simple if properly done. It must be remembered, however, that a normal or negative synovial fluid does not exclude any etiology from consideration.

Also, many laboratories are not able to deal optimally with synovial fluid specimens, and as a result, improper inferences may be drawn from the faulty analysis of synovial fluid. Of great importance is the need to scrupulously avoid the introduction of infection into the joint space. This is accomplished by using sterile technique, including gloves and a mask when necessary. A common error frequently made in joint aspiration is the failure to use a needle with a large enough bore to allow viscous pus, crystals, or cellular debris to pass into the syringe. This usually requires an 18-gauge or larger needle when a diagnostic arthrocentesis is performed. This needle size is not unduly traumatic and is not particularly painful if the injection site is initially infiltrated with local anesthetic.

Useful and important laboratory examinations on synovial fluid are

Table 4. Infectious Arthritis*

| Organism | Incidence | | Drug of Choice | Dosage | Duration of Therapy | Alternates | Comments |
|---|-----------|----------|---------------------------------------|--|---|---|---|
| | Adults | Children | | | | | |
| Gram Positive Cocci identified by Gram stain prior to culture results | 45% | 75% | Nafcillin | 100 mg/kg/day in 4-6 doses IV | 2 weeks or until joint is normal, then 4-6 weeks oral therapy | Cephalosporins Penicillins Erythromycin | Penicillin G resistant. Staphylococcal must be considered present until disproved |
| Staphylococcus Aureus Penicillin G resistant (culture proven) | 25% | 40% | Nafcillin | 100 mg/kg/day in 4-6 doses IV | 2 weeks or until joint is normal, then 4-6 weeks oral therapy | " | |
| Staphylococcus Aureus Penicillin G sensitive (culture proven) | | | Penicillin G | 2-3x10 ⁵ u/kg/day in 4-6 doses IV | " | " | |
| Diplococcus Pneumonia | 10% | 10% | Penicillin G | " | 10-14 days | " | Lower doses may be appropriate |
| Streptococcus | 10% | 25% | Penicillin G | " | " | " | |
| Gram Negative Cocci identified by Gram stain prior to culture results | 50% | 10% | Penicillin G (Ampicillin in children) | 200 mg/kg/day in 4-6 doses IV or IM | 10-14 days | Tetracycline Chloramphenicol | Avoid Tetracycline due to labeling of teeth |

*Repeated, frequent joint aspiration is necessary in all infected joints.

Table 5. Acute Gouty Arthritis

| History | Physical Examination | Laboratory Data | Treatment Modality |
|--|--|---|---|
| <p>1. Usually occurs in males over 30 years.</p> <p>2. Attack is sudden in onset.</p> <p>3. First attack is usually monarticular and affects the big toe, instep, ankle, and knee most commonly.</p> <p>4. The pain is excruciating and may be described as burning, crushing, or throbbing.</p> <p>5. Periods between attacks are usually symptom-free.</p> | <p>1. Joint is usually swollen, hot, dusky red, and tender.</p> <p>2. The big toe joint is the first affected in 50% of patients (podagra).</p> <p>3. Upper extremity joints are involved less frequently than those of the lower extremities.</p> | <p>1. The diagnosis is firmly made by the demonstration of intracellular uric acid crystals in the synovial fluid of the affected joint. These crystals are negatively birefringent by polarized microscopy.</p> <p>2. The serum uric acid level is almost always greater than 7 mg% by the uricase method.</p> <p>3. Sedimentation rate may be elevated.</p> <p>4. X-rays are usually negative with early disease.</p> <p>5. There may be decreased enzyme levels such as hypoxanthine-guanine phosphoribosyl transferase in certain subgroups of patients with gouty arthritis.</p> | <p>1. Oral Colchicine 0.6 mg qh until pain is relieved or toxic symptoms occur such as nausea, vomiting, or diarrhea. Rarely are more than 10 to 12 doses necessary. The acute arthritis of sarcoidosis has been reported to be responsive to this regimen.</p> <p>2. Intravenous Colchicine 2 to 4 mg IV may be repeated once in 4 hours if the response is questionable or incomplete. Gastrointestinal toxicity is minimized with this regimen.</p> <p>3. Indomethacin 150 to 200 mg by mouth in divided doses on the first day with a daily dose of 100 to 150 mg to follow for the next several days. This is considered to be as effective as Colchicine by many. Occasional toxicity is usually manifest as gastrointestinal upset, headache, or dizziness.</p> <p>4. Phenylbutazone or oxyphenbutazone 400 to 800 mg by mouth in divided doses on the first day with a dose of 100 mg 4 times a day for the next several days. Probably as effective as Indomethacin and Colchicine, but undesirable for chronic use because of toxicity.</p> <p>5. Steroids are undesirable because they are not as effective as the above agents and incur greater risk.</p> <p>6. Allopurinol and uricosuric agents have no place in the treatment of acute gouty arthritis and may cause exacerbation in the acute phase.</p> |

Table 6. Rheumatoid Arthritis with Acute Onset

| History | Physical Examination | Laboratory Data | Treatment |
|---|---|--|--|
| <p>1. Usually occurs in females in middle years, but can occur in any age group or sex.</p> <p>2. Attack is usually insidious but may be of rapid onset.</p> <p>3. Usually attacks symmetrically with multiple joint involvement.</p> <p>4. Generalized stiffness, most severe in the morning, is frequently a major complaint.</p> <p>5. Systemic complaints of fatigue, weakness, and malaise are frequent. Temperature higher than 103 F may occur but is rare and requires the exclusion of other etiologies.</p> | <p>1. Usually swelling, tenderness, and decreased motion of affected joint(s).</p> <p>2. Peripheral joints are usually affected. The proximal interphalangeal, metacarpophalangeal, and metatarsophalangeal joints are most often affected. The more proximal limb joints are frequently involved but the distal interphalangeal joints are rarely inflamed.</p> <p>3. Joint swelling consists of soft tissue and/or fluid, not bony enlargement.</p> <p>4. Nodules, bone, and joint destruction and extra-articular manifestations are not usually seen in new cases of rapid onset.</p> | <p>1. Westergren erythrocyte sedimentation rate is usually elevated.</p> <p>2. Rheumatoid factor may be absent early.</p> <p>3. Synovial fluid analysis usually shows less than 50,000 WBCs with 75% polys (PMNs) or less. The mucin clot may be poor and frequently the rheumatoid factor titer is significantly higher than that found in the serum while the complement level is lower.</p> <p>4. X-rays rarely reveal anything but soft tissue swelling early.</p> | <p>1. Rest and avoidance of trauma of affected joint(s).</p> <p>2. Range of motion exercises should begin early but should be very gentle in the severely inflamed joint.</p> <p>3. Early attention to proper position and posture is important. Night splints, etc, may be required to prevent flexion contractures.</p> <p>4. Virtually all patients should be tried on full salicylates (3.6 to 4.8 gm/day) before adding additional agents. (Serum salicylate level 20 to 30 gm/100 ml.)</p> <p>5. Non-steroidal anti-inflammatory agents, gold, and other agents should not be added until the diagnosis is secure and a fair trial for a conservative program has been accomplished.</p> |

Table 7. Juvenile Rheumatoid Arthritis

| History | Physical Examination | Laboratory Data | Treatment |
|--|--|---|--|
| <p>1. Onset is very acute in approximately 20% of cases. Systemic features may include high fever, skin rash, iridocyclitis, splenomegaly, and pericarditis, and arthritis may be absent.</p> <p>2. Fifty percent present with polyarticular disease.</p> <p>3. Thirty percent may present with monoarticular disease with no systemic features.</p> <p>4. Children may deny joint pain, but present with guarding, swelling, limited motion, or a limp.</p> <p>5. Transient temporomandibular joint complaints are common.</p> <p>6. Often difficult to distinguish from infection, malignancy, or rheumatic fever early.</p> | <p>1. Knees, wrists, ankles, and neck are common initially involved sites.</p> <p>2. Radial deviation is more common than ulnar deviation in children.</p> <p>3. Nodules are rarely found.</p> <p>4. Fever is frequent.</p> <p>5. Evanescent, trunkal, salmon-colored macular rash is found in 30%.</p> <p>6. Lymphadenopathy is frequent.</p> <p>7. Cardiorespiratory findings are rare.</p> <p>8. Iridocyclitis occurs in 10 to 15%, usually those with monoarticular onset. Requires slit lamp to detect.</p> | <p>1. No specific diagnostic tests available.</p> <p>2. Leukocytosis is frequent.</p> <p>3. Often mild normocytic, normochromic anemia.</p> <p>4. Sedimentation rate is usually elevated but need not be so.</p> <p>5. ASO titer may be elevated due to unrelated prior infection.</p> <p>6. Rheumatoid latex fixation is usually negative (10 to 20% positive).</p> <p>7. Synovial fluid is nonspecific, "inflammatory" in nature.</p> | <p>1. Adequate rest, including daytime rest periods.</p> <p>2. Avoidance of trauma to involved joints.</p> <p>3. Slit lamp examinations are necessary acutely and every three to six months.</p> <p>4. Active and passive exercise daily.</p> <p>5. Splints may help protect badly inflamed joints or avoid flexion deformities.</p> <p>6. Drugs should be restricted to salicylates, reserving steroids for the most severe cases. Gold and other anti-inflammatory agents may not be appropriate because of increased and potentially fatal toxicity.</p> <p>a. Aspirin 50 to 130 mg/kg/day. Toxicity is difficult to detect. Tinnitus is rare in children.</p> <p>b. Steroids should be held to lowest dosages possible and used only by physicians acquainted with potential problems.</p> |

aerobic, anaerobic, mycobacterial, and fungal cultures, gram stain, cell count with differential, and examination for crystals. In order to diagnose gouty arthritis or pseudogout, it is necessary to examine the synovial fluid for crystals by polarized light microscopy. While a standard microscope may identify crystals on occasion, it is impossible to characterize their birefringence. Most importantly, the crystals are frequently missed when standard light microscopy is used. It should be noted that on occasion cholesterol crystals may be present and, if there have been previous joint injections, steroid crystals may also be seen. The diagnosis is most secure when the crystals are detected intracellularly in the leukocytes.

Rheumatoid factor and synovial fluid complement are occasionally helpful, but rarely do serology, glucose, or enzyme levels add significantly to the diagnostic process in acute cases. Evaluation of the mucin clot has historical interest, but as it is friable in all forms of inflammation, it usually is not helpful in diagnosing arthritis of recent onset once trauma has been excluded. In evaluating recent onset joint pain when trauma has been excluded, x-rays are usually not helpful diagnostically. Synovial biopsy also contributes little if any specific information early. Typical synovial fluid parameters are found in Table 3.

Assessment and Plan

After an adequate data base related to the acute joint pain has been obtained, it must be integrated into an appropriate assessment of the problem. It should always be kept in mind that to make a definite diagnosis, specific diagnostic criteria must be met. The individual positive pieces of data are necessary to make the diagnosis, but by themselves are not sufficient. For example, many of Jones criteria⁷ may be present in a given patient and yet the diagnosis is clearly not acute rheumatic fever. Likewise, gouty arthritis⁸ should not be equated

with hyperuricemia nor do fever and leukocytosis exist only in pyogenic arthritis.⁹ Not only should the physician not be fooled in the assessment process, he/she must be able to act expeditiously and economically. In order to do so, the establishment of priorities is essential. The misdiagnosis or neglect of infectious arthritis or acute rheumatic fever may mean permanent crippling or even be life-threatening, and gonococcal arthritis or gouty arthritis are completely correctable if diagnosed early and properly treated. In the majority of the remaining disease processes presenting with acute arthritis, appropriate diagnosis and management can significantly alter the course of the illness in a positive fashion, but definitive diagnosis and treatment are not as urgent. While firm diagnostic criteria are not available for all diseases presenting with arthritis of recent onset, definite attempts are being made in this area.^{10,11}

For the physician to deal realistically with the problem of joint pain of recent onset, he/she must understand the natural history and course of the specific disease with which he/she is dealing. This is necessary in order to formulate appropriate objectives for the specific patient and establish a plan of action whereby they can be accomplished.

Once the most emergent conditions are excluded from the differential, the wisest approach is to observe the patient without treatment until the diagnosis is evident or the patient improves. Therapeutic trials in general should be avoided. They usually do not contribute insight to a particular problem and may mask a condition requiring specific intervention.

It is not possible to give specific details on all the possible disease processes responsible for joint pain of recent onset. However, the causes of acute onset arthritis in a few of the most common, representative diseases found in a busy primary care practice are summarized in Tables 4 through 7. Infectious arthritis (Table 4), including the articular manifestations of gonococcemia, is included because this is a true medical emergency when present and thus requires early and appropriate therapy. Gout (Table 5) is included not only because it is common, but because it is frequently overdiagnosed due to the failure of the

physician to use rigid diagnostic criteria.⁹ In addition, proper therapy is usually very effective and is thus most satisfying to both patient and physician. Rheumatoid arthritis (Table 6) is included as a reminder that chronic, systemic disease involving many joints often begins in a solitary joint and may appear suddenly. The diagnosis of juvenile rheumatoid arthritis (Table 7) is occasionally overlooked when it appears as other than typical Still's disease.

In summary, a systematic approach to the patient presenting with the problem of acute joint pain usually provides the correct diagnosis which in turn leads to appropriate therapy. The proper evaluation of this problem requires a detailed history and physical examination, as well as a rationally selected battery of laboratory studies. A less thorough approach to the problem frequently proves to be unsatisfactory for patient and physician alike.

References

1. Taranta A, Lehr JW: Post-streptococcal disease. In Samter M (ed): *Immunological Diseases*, ed 2. Boston, Little, Brown and Co, 1971, p 601
2. Schaller J, Wedgewood RJ: Juvenile rheumatoid arthritis: A review. *Pediatrics* 50:940, 1972
3. Thompson GR, Ferreira A, Brackett RG: Acute arthritis complicating rubella vaccination. *Arthritis Rheum* 14:19, 1971
4. Keiser H, Ruben FL, Wolinsky E, et al: Clinical forms of gonococcal arthritis. *N Engl J Med* 279:234, 1968
5. Alpert E, Isselbacher KJ, Schur PH: The pathogenesis of arthritis associated with viral hepatitis. *N Engl J Med* 285:185, 1971
6. Dubois EL (ed): *Lupus Erythematosus*, ed 2. New York, Blakiston Division, McGraw-Hill, 1973
7. Stollerman GH, Markowitz M, Taranta A, et al: Jones criteria (revised) for guidance in the diagnosis of rheumatic fever. *Circulation* 32:664, 1965
8. McCarty DJ: Pathogenesis and treatment of the acute attack of gout. *Clin Orthop* 71:28, 1971
9. Hollander JL, McCarty AJ, Jr (eds): *Arthritis and Allied Conditions*, ed 8. Philadelphia, Lea and Febiger, 1972
10. Rodnan GP (ed): *Primer on the Rheumatic Diseases*, ed 7. *JAMA* 224(5) (Suppl):661, 1973
11. Huskisson EC, Hart FD: *Joint Disease: All the Arthropathies*. Baltimore, Williams and Wilkins, 1973