

## EDITORIALS

## Ultrabrief Delirium Assessments—Are They Ready for Primetime?

Jin H. Han, MD, MSc<sup>1\*</sup>, Eduard E. Vasilevskis, MD, MPH<sup>2,3</sup>

<sup>1</sup>Department of Emergency Medicine, Center for Quality Aging, Vanderbilt University Medical Center, Nashville, Tennessee; <sup>2</sup>Department of Medicine, Center for Quality Aging, Vanderbilt University Medical Center, Nashville, Tennessee; <sup>3</sup>VA Tennessee Valley Healthcare System, Geriatric Research, Education and Clinical Center, Nashville, Tennessee.

Delirium is a form of acute brain failure that affects up to 64% of older hospitalized patients and is associated with a multitude of adverse outcomes.<sup>1</sup> Healthcare providers, regardless of clinical setting, do not identify delirium in approximately 75% of cases.<sup>2,3</sup> The paucity of brief and simple delirium assessment tools has been a barrier to improving delirium recognition.

To address this unmet need, several ultrabrief (<30 seconds) delirium assessment tools have been recently studied. In this issue of the *Journal of Hospital Medicine*, Fick et al. evaluated 20 individual components of the 3-minute diagnostic interview for delirium using the Confusion Assessment Method (3D-CAM), which was recently validated in older hospitalized patients.<sup>4,5</sup> They observed that the best-performing single-item delirium assessment was the months of the year backward (MOTYB) task from December to January. This task assesses for inattention, a cardinal feature of delirium. Using a cutoff of 1 or more errors, the MOTYB was 83% sensitive and 69% specific for delirium.<sup>5</sup> By adding “name the day of the week,” the sensitivity increased to 93% with similar specificity (64%). This supports research by O’Regan et al., who examined MOTYB, but defined a positive screen if they could not recite the months backward from December to July perfectly. They observed a sensitivity and specificity of 84% and 90%, respectively, in older hospitalized patients.<sup>6</sup>

The assessment of arousal, another feature of delirium, has also garnered significant interest as another ultrabrief delirium screening method. Arousal is the patient’s responsiveness to the environment and can be assessed during routine clinical care. Fick et al. observed that impaired arousal using the 3D-CAM was 19% sensitive for delirium. This is in contrast to others who have reported sensitivities of 64% to 84%.<sup>7–9</sup> The difference in sensitivity may in part be explained by the method used to detect arousal. The 3D-CAM asks, “Was the patient sleep/stuporous?” or “Was the patient hypervigilant?” Previous studies used the Rich-

mond Agitation Sedation Scale (RASS), an arousal scale based on eye contact and physical behaviors to assess patients from –5 (coma) to +4 (combative).<sup>10</sup> Therefore, it is important to consider the method of arousal assessment if using this feature for delirium screening.

These ultrabrief delirium assessments would be even more clinically useful if they identified patients at high risk for adverse outcomes. In this same journal issue, 2 studies evaluated the prognostic ability of several ultrabrief delirium assessments. Zdravec et al. observed that an abnormal RASS was a moderately good predictor of 24-hour mortality, with an area under the receiver operating characteristic curve of 0.82.<sup>11</sup> Yevchak et al. observed that an abnormal RASS or MOTYB was associated with longer hospital length of stays, increased in-hospital mortality, and need for skilled nursing.<sup>12</sup>

Viewed as a whole, these studies represent a significant advancement in delirium measurement and have the potential to improve this quality-of-care issue. However, uncertainties still exist. (1) Can these ultrabrief delirium assessments be used as standalone assessments? Based upon current data, these assessments have a significant proportion of false negative and positive rates. The effect on such misclassification on patient outcomes and healthcare utilization needs to be clarified. Because of this concern, Fick et al. recommended performing a more specific delirium assessment in those who have a positive MOTYB screen.<sup>5</sup> (2) What is the optimal cutoff of the MOTYB task and does this cutoff vary in different patient populations? The optimal cutoff will depend on whether or not a more sensitive test (lower error threshold) or specific test (higher error threshold) is desired. The optimal cutoff may also depend on the patient population (eg, demented versus nondemented). (3) Most important to practicing hospitalist and patients, will introducing these ultrabrief delirium assessments improve delirium recognition and improve patient outcomes? The impetus for widespread implementation of these assessments would be strengthened if healthcare providers successfully applied these assessments in clinical practice and subsequently improved outcomes.

In conclusion, the MOTYB and the assessment of arousal may be reasonable alternatives to more conventional delirium screening, especially in clinical environments with significant time constraints. However, additional research is needed to better refine

\*Address for correspondence and reprint requests: Jin H. Han, MD, Department of Emergency Medicine, Vanderbilt University Medical Center, 703 Oxford House, Nashville, TN 37232-4700; Telephone: 615-936-0253; Fax: 615-936-1316; E-mail: jin.h.han@vanderbilt.edu

Received: August 17, 2015; Accepted: August 18, 2015  
2015 Society of Hospital Medicine DOI 10.1002/jhm.2478  
Published online in Wiley Online Library (Wileyonlinelibrary.com).

these instruments to the clinical environment they will be used and determine how they impact clinical care and patient outcomes.

Disclosures: Dr. Han is supported the National Heart, Lung, and Blood Institute (K12HL109019). Dr. Vasilevskis is supported by the National Institutes of Health (K23AG040157) and the Geriatric Research, Education and Clinical Center (GRECC). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the Department of Veterans Affairs. The authors report no conflicts of interest.

## References

1. Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. *Lancet*. 2014;383(9920):911–922.
2. Collins N, Blanchard MR, Tookman A, Sampson EL. Detection of delirium in the acute hospital. *Age Ageing*. 2010;39(1):131–135.
3. Han JH, Zimmerman EE, Cutler N, et al. Delirium in older emergency department patients: recognition, risk factors, and psychomotor subtypes. *Acad Emerg Med*. 2009;16(3):193–200.
4. Marcantonio ER, Ngo LH, O'Connor M, et al. 3D-CAM: derivation and validation of a 3-minute diagnostic interview for CAM-defined delirium: a cross-sectional diagnostic test study. *Ann Intern Med*. 2014;161(8):554–561.
5. Fick DM, Inouye SK, Guess J, et al. Preliminary development of an ultrasbrief two-item bedside test for delirium. *J Hosp Med*. 2015;10(00):000–000.
6. O'Regan NA, Ryan DJ, Boland E, et al. Attention! A good bedside test for delirium? *J Neurol Neurosurg Psychiatry*. 2014;85(10):1122–1131.
7. Chester JG, Beth Harrington M, Rudolph JL, VA Delirium Working Group. Serial administration of a modified Richmond Agitation and Sedation Scale for delirium screening. *J Hosp Med*. 2012;7(5):450–453.
8. Tieges Z, McGrath A, Hall RJ, Maclullich AM. Abnormal level of arousal as a predictor of delirium and inattention: an exploratory study. *Am J Geriatr Psychiatry*. 2013;21(12):1244–1253.
9. Han JH, Vasilevskis EE, Schnelle JF, et al. The diagnostic performance of the Richmond Agitation Sedation Scale for detecting delirium in older emergency department patients. *Acad Emerg Med*. 2015;22(7):878–882.
10. Sessler CN, Gosnell MS, Grap MJ, et al. The Richmond Agitation-Sedation Scale: validity and reliability in adult intensive care unit patients. *Am J Respir Crit Care Med*. 2002;166(10):1338–1344.
11. Zdravec F, Tien L, Robertson-Dick B, et al. Comparison of mental status scales for predicting mortality on the general wards. *J Hosp Med*. 2015;10(10):658–663.
12. Yevchak AM, Doherty K, Archambault E, Kelley KW, Fonda JR, Rudolph JL. The association between an ultrasbrief cognitive screening in older adults and hospital outcomes. *J Hosp Med*. 2015;10(10):651–657.